



Effect of Thyroid Hormones on the Redox Balance of Broiler Chickens

Lin H.^{1,2}, E. Decuypere² and J. Buyse^{2,*}

¹ Department of Animal Science, Shandong Agricultural University, Taian, Shandong 271018, China

ABSTRACT : In the present study, two trials were conducted to evaluate the effects of hyper- and hypothyroid status on the redox balance of broiler chickens. In Trial 1, 3 groups of broiler chickens were randomly subjected to one of the three treatments: subcutaneous administration of triiodothyronine (T_3 , 150 $\mu\text{g}/\text{kg}$ BW), methimazole (MMI, 150 mg/kg BW), or saline. The blood, liver and heart were sampled at 3 h after injection. In Trial 2, three groups of 20 broiler chickens were randomly fed with one of the three diets: control, dietary supplementation of T_3 (1.5 mg/kg diet) or MMI (1 g/kg diet) for 7 days. In trial 1, the plasma concentrations of T_3 and T_3 to thyronine ratio (T_3/T_4) were significantly increased by T_3 injection. Plasma levels of thiobarbituric acid reacting substances (TBARS) tended to be increased ($p = 0.067$) by both T_3 and MMI treatments while the ferric reduced/antioxidant capacity (FRAP) was increased only by MMI treatment. Acute T_3 treatment had no significant effect on the activities of superoxide dismutase (SOD) and the concentrations of FRAP and TBARS in either liver or heart tissue. In contrast, the hepatic activities of SOD were decreased ($p < 0.05$) while the cardiac levels of FRAP were significantly increased ($p < 0.0001$) by MMI treatment. In chronic treatments, the rectal temperature of chickens was significantly decreased ($p < 0.05$) by MMI treatment. The circulating T_3 levels were significantly increased ($p < 0.05$) by long-term T_3 treatment, and showed a trend to decrease in MMI treatment. The plasma concentrations of TBARS were significantly ($p < 0.05$) increased by MMI treatment. All the redox parameters measured in either liver or heart were not significantly altered by either long-term T_3 or MMI treatment except that the hepatic SOD activities were significantly augmented by T_3 treatment. The result showed that neither acute nor long-term elevation of circulating T_3 levels induced lipid peroxidation in broiler chickens. The enhanced enzymatic antioxidant system (SOD in cardiac tissue) may be involved in the protection of the bird to increased oxidative challenge. The responses of redox balance to changed thyroid state seem to be tissue specific. (**Key Words :** Thyroid Hormone, Broiler Chickens, Lipid Peroxidation)

INTRODUCTION

Thyroid hormones are known to regulate the energy metabolism of most tissues including liver, kidney, heart, and skeletal muscles. It is well established that thyroid hormones accelerate the basal metabolic rate and oxidative metabolism by causing an increase in the mitochondria mass, mitochondrial cytochrome content and respiratory rate. In chickens, the plasma levels of thyroid hormones are changed with ambient temperature, age, feeding status and pathophysiologic status (Decuypere and Kühn, 1984, 1988; Lin et al., 2000; Stojevic et al., 2000; Luger et al., 2002). One the other hand, the reactive oxygen species (ROS) are generated as by-products of oxidative metabolism in

mitochondria.

In chickens, there is limited information about the association of thyroid status with the induction of oxidative stress. In mammals, it has been suggested that the susceptibility of mitochondria to oxidative stress is increased by hyperthyroidism (Das and Chainy, 2001; Venditti et al., 2003) and decreased by hypothyroidism (Venditti et al., 2003). The administration of triiodothyronine (T_3) to hypothyroid rats results in a significant augmentation of oxidative stress parameters (Das and Chainy, 2001). In hyperthyroidism rats, enhanced oxidative stress parameters are observed in the heart (Asayama et al., 1987; Zation et al., 1993; Civelek et al., 2001), soleus muscles (Asayama et al., 1987; Zation et al., 1993), and liver (Fernandez and Videla, 1996; Venditti et al., 1997). Hypothyroidism does not modify or even decrease lipid peroxidation and oxidative damages (Venditti et al., 1997; Gredilla et al., 2001a). However, hypothyroid female patients were found to have significantly higher level of plasma TBARS, compared to healthy ones (Komukoglu et

* Corresponding Author: Johan Buyse. Tel: +32-16-328525, Fax: +32-16-321994, E-mail: Johan.Buyse@biw.kuleuven.ac.be

² Lab of Physiology and Immunology of Domestic Animals, Department of Biosystems, Catholic University Leuven, Kasteelpark Arenberg 30, 3001 Leuven, Belgium.

Received June 1, 2007; Accepted October 22, 2007

al., 2002). The enzymatic and not-enzymatic antioxidant systems are involving in the enhanced oxidative stress in hypothyroid patients (Adali et al., 1999; Resch et al., 2002).

Broiler chickens are more susceptible to environmental challenge and the lipid peroxidation is one of the undesired results. For example, exogenous glucocorticoid administration (Lin et al., 2004 a, b) and heat stress (Lin et al., 2000; Shim et al., 2006) resulted in lipid peroxidation in broiler chickens, which should be at least partially responsible for the reduced zootechnical performance. Moreover, the maintenance of redox balance has been found to be important for the health (Bottje et al., 1998; Iqbal et al., 2002) and meat quality of broiler chickens (Jiang et al., 2007). In modern strains of broiler chickens, the selection for high feed conversion ratio could result in functional hypothyroidism by either decreased thyroid hormone production or peripheral deiodination of thyroxine, which is believed to be linked with the susceptible of broiler chickens to ascites (Decuypere et al., 2003). The impaired redox balance is involved in the development of ascites (Iqbal et al., 2002; Han et al., 2005). Hence, we hypothesize that the changed thyroid status may play a role in redox balance of broiler chickens. Moreover, as chickens have quite different metabolic characteristics such as high body temperature and metabolic rates and enhanced antioxidant system comparing to mammals, we speculated that chickens may response to hypo- and hyperthyroid status in a different way as that of mammals.

In the present study, two trials were conducted to evaluate the involvement of changed thyroid status in the redox balance in broiler chickens. The hyperthyroidism or hypothyroidism status was respectively mimicked by dietary and subcutaneous injection of T_3 or methimazole (MMI). MMI can block thyroid hormone biosynthesis by inhibiting thyroid peroxidase, a key enzyme of thyroid hormone biosynthesis, and has been used as an antithyroid drug (Cooper, 1984).

MATERIALS AND METHODS

Chickens and diets

Commercial broiler chickens (Cobb) were obtained from local hatchery at one day of age and reared in an environmentally controlled room. The brooding temperature was maintained at 35°C for the first 2 d, and then decreased gradually to 21°C (45% RH) at 28 d and maintained as such thereafter. The broilers received a commercial starter diet (12.10 MJ ME/kg and 22.2% CP) until 7 d of age, after which a commercial grower diet (12.68 MJ/kg, 21.4% crude protein) was provided until the end of the experiment. For a detailed description of both diets, see Buyse et al. (2001). The light regime was 23 L:1 D. The birds had free access to feed and water during rearing period.

Trial 1: Acute subcutaneous administration

Eighteen 32-d-old broiler chickens of both sexes with similar body weight were randomly divided into three groups of 6 chickens and subjected randomly to the three treatments: subcutaneous administration of T_3 (150 μ g/kg BW, Sigma Chemical), MMI (150 mg/kg BW, Sigma Chemical) or saline (serving as control). A blood sample of each chicken was respectively obtained before and at 3 h after injection from a wing vein with a heparinized syringe, and collected in iced tubes. At the same time, the body weight (BW) loss was recorded and rectal temperature was determined with a digital thermometer ($\pm 0.1^\circ\text{C}$, Huger Electronics GmbH, Germany). During the 3 h experimental period, feed was withdrawn and experimental chickens had free access to water.

Immediately after the second blood sample was obtained, all chickens were sacrificed by cervical dislocation and the liver and heart were dissected immediately. After washing with cold saline (4°C, 0.95%) and removed of blood, the tissues were frozen in liquid nitrogen and stored at -76°C until analysis. Plasma was obtained after centrifugation at 3,000 rpm for 10 min at 4°C and was stored below -20°C for further analysis.

Trial 2: Dietary treatment

Sixty chickens of 28-d of age were randomly assigned to three groups of 20 chickens and fed with one of the three diets: control, diet supplemented with T_3 (1.5 mg/kg diet) or MMI (1 g/kg diet), from 31 to 38 days of age. At 38 days of age, rectal temperature was determined with a digital thermometer for each chicken. Thereafter, 7 birds of each group were sacrificed by cervical dislocation after a blood sample was drawn from a wing vein with heparinized syringe and the liver and heart were dissected immediately. After washing with cold saline (4°C, 0.95%) and removing blood, the tissues were frozen in liquid nitrogen and stored at -76°C until analysis. During experimental period, all the chickens had free access to feed and water.

Plasma parameters

Commercial colorimetric diagnostic kits were used to measure plasma glucose (IL TestTM kit, No. 182508-00), urate (IL TestTM kit, No. 181685-00) and creatine kinase (CK) (IL TestTM kit, No. 181605-90), using the MonarchTM 2000 Chemistry system Model 760 (Monarch Chemistry System, Instrumentation Laboratories, B-1930, Zaventem, Belgium). Corticosterone (CORT) was measured using a sensitive and highly specific radioimmunoassay kit (IDS, Inc., Boldon, NE35 9PD, UK), with a sensitivity of 0.39 ng/ml, and low cross reactions with aldosterone (0.20%), cortisol (0.40%) and deoxycorticosterone (3.30%). The intra assay variability was 3.8%. Before assayed, plasma samples were heated at 80°C for 10 minutes to inactivate

Table 1. Effects of acute (3 h) subcutaneous injection of triiodothyronine (T₃, 150 µg T₃/kg BW) or methimazole (MMI, 150 mg/kg BW) or saline (control treatment) on plasma parameters of broiler chickens in Trial 1

	T ₃	MMI	Control
T ₃ (ng/ml)	150.1±20.9 ^x	4.84±1.29 ^{xy}	3.52±0.88 ^y
T ₄ (ng/ml)	16.7±3.5	10.5±2.4	12.1±1.6
T ₃ /T ₄ ratio	12.8±4.3 ^x	0.65±0.3 ^y	0.37±0.1 ^y
CORT (ng/ml)	16.2±4.0	26.6±3.4	24.7±8.3
TBARS (nmol/ml)	0.53±0.05	0.55±0.05	0.38±0.05
FRAP (µmol/ml)	807±45 ^y	2,353±117 ^x	715±54 ^y
CK (IU/L)	5,927±1,443	7,219±1,665	4,684±795
Urate (mg/d)	2.3±0.3	2.7±0.3	2.1±0.4
Glucose (mg/dl)	235±5	230±9	238±6

Values are means±SEM (n = 6).

^{xy} Means with different superscript within the same row differ significantly (p<0.05).

CK = Creatine kinase; CORT = Corticosterone; FRAP = Ferric reducing/antioxidant power.

T₃ = Triiodothyronine; T₄ = Thyroxine; TBARS = Thiobarbituric acid reacting substances.

corticosterone-binding proteins.

Thyroid hormones were measured by radioimmunoassay according to the method described by Darras et al. (1996). Briefly, T₃ measurements were performed using a commercial available T₃ antiserum from Byk-Sangtec (Byc-Sangtec Diagnostica GmbH, Dietzenbach, Germany), in combination with a specific tracer (Amersham International, Slough, England). The intra-assay coefficient of variation was 4.5%. T₄ concentrations were assayed using tracer from Amersham (Byc-Sangtec Diagnostica GmbH, Dietzenbach, Germany) and a rabbit T₄ antiserum (Mallinckrodt Diagnostica, Dietzenbach, Germany). This T₄ antiserum had 0.16% cross-reactivity with T₃. The intra-assay coefficient of variation was 3.2%.

Plasma lipid peroxidation was estimated by spectrophotometric determination of thiobarbituric acid reacting substances (TBARS) with a method of Lin et al (2004a). TBARS were expressed as nmoles of malondialdehyde (MDA) per ml plasma. Total plasma antioxidant activity was determined by the ferric reducing/antioxidant power (FRAP) assay as described by Benzie and Strain (1996, 1999). The measurement was conducted at room temperature and a 5-min time window was used.

Tissue parameters

Tissue samples were homogenised in 9 volumes of 10 mM sodium phosphate buffer (pH 7.4) containing 1.15% potassium chloride. TBARS of tissue homogenates was determined spectrophotometrically according to Lin et al. (2004a, b). TBARS were expressed as nmoles of MDA per gram of wet tissue. After the homogenates were centrifuged (4,000 rpm, for 10 min at 4°C), the supernatant was used for the measurement of superoxide dismutase (SOD) and FRAP. The SOD activity was measured with the commercial kit (Dojindo Inc.) using a microplate reader (Titertek Multiskan[®], MCC/340).

All samples were run in the same assay to avoid inter-assay variability.

Statistical analysis

Data in Trial 1 were subjected to Repeated Measures ANOVA by SAS for windows (version 8e, SAS Institute Inc.) to evaluate the effects of treatment and time, as well as the interaction between treatment and time. In Trial 2, a one-way model of ANOVA was used to estimate the main effects of treatment. Multiple comparisons were performed between treatments means using Duncan test. Means were considered significantly different at p<0.05.

RESULTS

In trial 1, at 3 h after treatment, the BW and RT of experimental chickens were all decreased in all treatments but there was no significant difference in BW loss (T₃, 17.2±3.0 g; MMI, 8.3±13.1 g; Control, 20.2±5.1 g) and RT reduction (T₃, -0.4±0.1°C; MMI, -0.4±0.1°C; Control, -0.3±0.1°C) between treatments (p>0.05). The effects of treatment were significant (p<0.05) for the plasma concentrations of FRAP and T₃ and T₃/T₄ ratio, but not for plasma levels of T₄, TBARS, glucose, urate and CORT, and activity of CK (p>0.05) (Table 1). Compared to control group, the plasma concentrations of T₃ and T₃/T₄ ratio were significantly increased by T₃ treatment but were not affected by MMI treatment. The plasma concentrations of TBARS tended to be increased (p = 0.067) by both T₃ and MMI treatments, while the plasma levels of FRAP were significantly increased (p<0.05) by MMI treatment. In liver, the activities of SOD were significantly lower (p<0.05) in MMI chickens, compared to control or T₃ chickens, while the concentrations of TBARS and FRAP were not significantly (p>0.05) changed (Table 2). Neither the activities of SOD nor the concentrations of TBARS in heart were significantly affected by treatments (p>0.05). In

Table 2. Effects of acute (3 h) subcutaneous injection of triiodothyronine (T₃, 150 µg T₃/kg BW) or methimazole (MMI, 150 mg/kg BW) or saline (control treatment) on the biomarkers of oxidative stress in liver and heart tissues of broiler chickens in Trial 1

	T ₃	MMI	Control
Liver			
SOD (U/mg protein)	13.33±1.17 ^{xy}	11.47±0.26 ^y	15.10±0.61 ^x
TBARS (nmol/mg protein)	0.28±0.04	0.25±0.00	0.28±0.01
FRAP (µmol/mg protein)	131.4±6.7	126.2±4.0	129.5±6.4
Heart			
SOD (U/mg protein)	3.53±0.28	3.08±0.19	2.71±0.40
TBARS (nmol/mg protein)	0.27±0.01	0.23±0.01	0.25±0.02
FRAP (µmol/mg protein)	50.8±2.4 ^y	68.2±1.5 ^x	51.5±2.9 ^y

Values are means±SEM (n = 6).

^{xy} Means with different superscript within the same row differ significantly (p<0.05).

SOD = Superoxide dismutase; FRAP = Ferric reducing/antioxidant power; TBARS = Thiobarbituric acid reacting substances.

Table 3. Effects of dietary supplementation of triiodothyronine (T₃, 1.5 mg T₃/kg diet) or methimazole (MMI, 1 g/kg diet) on rectal temperature and plasma parameters of broiler chickens in Trial 2

	T ₃	MMI	Control
Rectal temperature (°C)	42.5±0.1 ^x	42.0±0.1 ^y	42.5±0.1 ^x
T ₃ (ng/ml)	9.64±4.57 ^x	1.72±0.42 ^y	2.08±0.33 ^y
T ₄ (ng/ml)	4.47±0.55	5.53±0.94	7.67±1.30
T ₃ /T ₄	2.65±1.46	0.37±0.09	0.30±0.09
CORT (ng/ml)	14.2±4.2	18.20±4.48	15.11±3.33
TBARS (nmol/ml)	0.63±0.06 ^y	0.84±0.09 ^x	0.61±0.06 ^y
FRAP (µmol/ml)	817±55	961±31	829±49
CK (IU/L)	4,677±843	3,337±281	5,283±1,624
Urate (mg/dl)	2.96±0.34	3.3±0.2	3.2±0.4
Glucose (mg/dl)	255±7	264±5	262±6

Values are means±SEM (n = 7).

^{xy} Means with different superscript within the same row differ significantly (p<0.05).

CK = Creatine kinase; CORT = Corticosterone; FRAP = Ferric reducing/antioxidant power.

T₃ = Triiodothyronine; T₄ = Thyroxine; TBARS = Thiobarbituric acid reacting substances.

contrast, the cardiac FRAP levels were significantly increased (p<0.0001) by MMI treatment compared to T₃ and control treatments.

After the chronic dietary treatment, the RT was significantly decreased (p<0.05) by MMI treatment (Table 3). The circulating T₃ levels were significantly increased (p<0.05) by dietary supplementation of T₃, and showed a trend to decrease in MMI treatment. The plasma concentrations of TBARS were significantly increased by MMI treatment (p<0.05) compared to other two treatments. The plasma levels of FRAP showed a trend to be increased by dietary supplementation of MMI (p = 0.073) compared to other treatments. The plasma levels of T₄, CK, glucose, urate, CORT and T₃/T₄ ratio were not significantly changed (p>0.05) by dietary treatment (Table 3). Dietary treatments had a significant effect on SOD activity in liver (p<0.02) but not in heart tissue (p>0.05). Hepatic SOD activities were significantly (p<0.05) increased by dietary T₃ supplementation (Table 4). The concentrations of TBARS and FRAP in both heart and liver tissues were not significantly changed by different dietary treatments (p>0.05).

DISCUSSION

Hyperthyroidism has been proven to link with the occurrence of oxidative stress in mammals (Fernandez et al., 2005; Venditti and Di Meo, 2006). In the present study, T₃ administration was used to induce hyperthyroidism of broiler chickens (Decuypere et al., 1987) and the elevated circulating levels of T₃ were observed in Trial 1 of the present study. In Trial 1, the plasma concentrations of TBARS, as a biomarker of oxidative injury, showed a trend to be increased by T₃ administration at 3-h later, with no obvious changes in all the other blood parameters, such as the levels of total antioxidant capacity (FRAP), urate and CORT. Moreover, the unobviously changed levels of TBARS and FRAP and activity of SOD in liver and heart indicated the unaffected redox balance in these tissues, suggesting that acute upregulation of thyroid hormones may not induce oxidative injury within a short period (<3 h).

The long-term effect of hyperthyroidism was further explored in Trial 2 of this study. In line with the previous report (Decuypere et al., 1987), dietary T₃ supplementation significantly increased the plasma levels of T₃. In mammals,

Table 4. Effects of dietary supplementation of triiodothyronine (T₃, 1.5 mg T₃/kg diet) methimazole (MMI, 1 g/kg diet) on the biomarkers of oxidative stress in liver and heart tissues of broiler chickens in Trial 2

	T ₃	MMI	Control
Heart			
SOD (U/mg protein)	4.48±0.21	4.57±0.18	4.13±0.35
TBARS (nmol/mg protein)	0.28±0.01	0.28±0.03	0.31±0.02
FRAP (µmol/mg protein)	48.0±1.3	52.1±1.8	51.8±2.0
Liver			
SOD (U/mg protein)	15.0±1.34 ^x	12.9±0.87 ^{xy}	9.95±0.96 ^y
TBARS (nmol/mg protein)	0.27±0.02	0.39±0.07	0.30±0.03
FRAP (µmol/mg protein)	112±7	115±8	125±5

Values are means±SEM (n = 7).

^{xy} Means with different superscript within the same row differ significantly (p<0.05).

SOD = Superoxide dismutase; TBARS = Thiobarbituric acid reacting substances; FRAP = Ferric reducing/antioxidant power.

it has been suggested that the susceptibility of tissues to oxidative stress is increased by hyperthyroidism. The augmented lipid peroxidation and changed levels of antioxidant substances and activities of enzymes were observed in hyperthyroid rats (Asayama et al., 1987; Zation et al., 1993; Fernandez and Videla, 1996; Venditti et al., 1997; Civelek et al., 2001). In contrast with these studies in mammals, lipid peroxidation was not obviously altered in hyperthyroid chickens in the present study. The oxidative damage, however, is not necessarily linked to increased reactive oxygen species (ROS) due to the altered thyroid state, as the antioxidant systems could respond to changes in the production of ROS (Mano et al., 1995b; Civelek et al., 2001). The significantly increased hepatic SOD activities together with the lack of obvious change of TBARS implies that the enhanced enzymatic scavenge capacity may compensate for the changes induced by hyperthyroidism. In addition, as these phenomena were not observed in heart, this influence seems to be tissue specific, which was found in mammals as well (Venditti et al., 1997).

On the other hand, the induction of lipid peroxidation by hyperthyroidism depends not only on the production of ROS, but also on the susceptibility of fatty acids to oxidative attack. The endogenous lipid peroxidation, sensitivity to lipid peroxidation and fatty acid unsaturation in liver were decreased in mouse treatment with T₄, indicating that the lipid changes observed in hyperthyroid animals can protect them against an increased oxidative attack to tissue lipids (Guerrero et al., 1999). Compared to mammals of similar body weight, birds have significantly lower degree of fatty acid unsaturation, which in turn leads to a lower sensitivity to lipid peroxidation (Pamplona et al., 1999a, b). The low rates of mitochondrial free radical production and relative lower degree of fatty acid unsaturation of avian tissue cellular membranes may protect them better from oxidative stress compared to mammals (Barja, 2002). Therefore, these results may infer that lower susceptibility of avian species to hyperthyroid state-induced oxidative stress.

Methimazole (MMI) can block thyroid hormone

biosynthesis by inhibiting thyroid peroxidase (Cooper, 1984). In the present study, plasma levels of thyroid hormones were not significantly affected by a single injection of MMI (150 mg/kg BW), indicating the unsuccessful induction of hypothyroidism in chickens. The increased total antioxidant capacities in MMI chickens may be related to the inherent antioxidant characteristics of MMI, which has already been proven *in vitro* (Asmus et al., 1996). Methimazole could inhibit the biosynthesis of thyroid hormone by impairing the formation of H₂O₂ or scavenging H₂O₂ directly as an antioxidant *in vivo*, which is a limiting step in thyroid hormone biosynthesis (Kim et al., 2001; Ferreira et al., 2003).

In trial 2 of this study, MMI was supplemented in the diet for one week to induce hypothyroidism. The relatively lower plasma levels of T₃ and the significantly decreased rectal temperature in MMI supplemental chickens are indication of their hypothyroid status. In line with the observation in hypothyroid patients (Konukoglu et al., 2002), the significantly augmented plasma lipid peroxidation in MMI chickens indicated the oxidative stress is induced by MMI treatment. In other studies with mammals, however, hypothyroidism does not modify or even decreases lipid peroxidation and oxidative damage (Venditti et al., 1997; Gredilla et al., 2001a), which may be related to the altered defence systems of antioxidants (Mano et al., 1995b; Gredilla et al., 2001b; Resch et al., 2002; Konukoglu et al., 2002; Brzezinska-Slebodzinska, 2003; Choudhury et al., 2003). However, the lack of obvious changes of TBARS levels, SOD activity and total antioxidant capacity in the liver and heart of MMI chickens compared to control group, indicate that redox status in these tissues were not affected by hypothyroidism. Therefore, whether the augmented lipid peroxidation should be ascribed to the hypothyroidism or MMI per se should be investigated further.

In conclusion, neither acute nor long-term elevation of circulating T₃ induces lipid peroxidation in broiler chickens. The enhanced enzymatic antioxidant system (e.g. in cardiac tissue) may be involved in the protection of birds from the

increased oxidative challenge induced by altered thyroid status.

ACKNOWLEDGMENT

The diligent technical assistance of C. Borgers and G. Nackaerts is gratefully appreciated. Hai Lin was supported by grant from National Basic Research Program of China (2004CB117507), grant from Program for New Century Excellent Talents in University and Grant F/00/89 from the Research Fund - K. U. Leuven.

REFERENCES

- Adali, M., M. Inal-Erden, A. Akalin and B. Efe. 1999. Effects of propylthiouracil, propranolol, and vitamin E on lipid peroxidation and antioxidant status in hyperthyroid patients. *Clin. Biochem.* 32:363-367.
- Asayama, K., K. Dobashi, H. Hayashibe, Y. Megata and K. Kato. 1987. Lipid peroxidation and free radical scavengers in thyroid dysfunction in the rat: a possible mechanism of injury to heart and skeletal muscle in hyperthyroidism. *Endocrinol.* 121: 2112-2118.
- Asmus, K. D., R. V. Bensasson, J. L. Bernier, R. Houssin and E. J. Land. 1996. One-electron oxidation of ergothioneine and analogues investigated by pulse radiolysis: redox reaction involving ergothioneine and vitamin C. *Biochem. J.* 315:625-629.
- Barja, G. 2002. Rate of generation of oxidative stress-related damage and animal longevity. *Free Radic. Biol. Med.* 33: 1167-1172.
- Benzie, I. F. and J. J. Strain. 1996. The ferric reducing ability of plasma (FRAP) as a measure of "antioxidant power": the FRAP assay. *Anal. Biochem.* 239:70-76.
- Benzie, I. F. and J. J. Strain. 1999. Ferric reducing/antioxidant power assay: direct measure of total antioxidant activity of biological fluids and modified version for simultaneous measurement of total antioxidant power and ascorbic acid concentration. *Methods Enzymol.* 299:15-27.
- Bottje, W. G., S. Wang, F. J. Kelly, C. Dunster, A. Williams and I. Mudway. 1998. Antioxidant defenses in lung lining fluid of broilers: impact of poor ventilation conditions. *Poult. Sci.* 77: 516-522.
- Brzezinska-Slebodzinska, E. 2003. Influence of hypothyroidism on lipid peroxidation, erythrocyte resistance and antioxidant plasma properties in rabbits. *Acta Vet. Hung.* 51:343-351.
- Buyse, J., G. P. J. Janssens and E. Decuypere. 2001. The effect of dietary L-carnitine supplementation on the performance, organ weights and circulating hormone and metabolite concentrations of broiler chickens reared under a normal or low temperature schedule. *Br. Poult. Sci.* 42:230-241.
- Choudhury, S., G. B. Chainy and M. M. Mishro. 2003. Experimentally induced hypo- and hyper-thyroidism influence on the antioxidant defence systems in adult rat testis. *Andrologia.* 35:131-140.
- Civelek, S., O. Seymen, A. Seven, G. Yigit, H. Hatemi and G. Burcak. 2001. Oxidative stress in heart tissue of hyperthyroid and iron supplemented rats. *J. Toxicol. Environ. Health* 23: 499-506.
- Cooper, D. S. 1984. Antithyroid drugs. *N. Engl. J. Med.* 311: 1353-1362.
- Darras, V. M., S. P. Kotanen, K. L. Geris, L. R. Berghman and E. R. Hühn. 1996. Plasma thyroid hormone levels and iodothyronine deiodinase activity following an acute glucocorticoid challenge in embryonic compared with posthatch chickens. *Gen. Comp. Endocrinol.* 104:203-212.
- Das, K. and G. B. Chainy. 2001. Modulation of rat liver mitochondrial antioxidant defence system by thyroid hormone. *Biochem. Biophys. Acta* 1537:1-13.
- Decuypere, E. and E. R. Kühn. 1984. Effect of fasting and feeding time on circadian rhythms of serum thyroid hormone concentrations, glucose, liver monodeiodinase activity and rectal temperature in growing chickens. *Dom. Anim. Endocrinol.* 1:251-262.
- Decuypere, E. and E. R. Kühn. 1984. Alterations in thyroid hormone physiology induced by temperature and feeding in newly hatched chickens. *Acta Physiol.* 39:380-391.
- Decuypere, E., J. Buyse, C. G. Scanes, L. Huybrechts and E. R. Kühn. 1987. Effects of hyper- or hypothyroid status on growth, adipose and levels of growth hormone, somatomedin C and thyroid metabolism in broiler chickens. *Reprod. Nutr. Develop.* 27:71-81.
- Decuypere, E., V. Bruggeman, G. F. Barbato and J. Buyse. 2003. Growth and reproduction problems associated with selection for increased broiler meat production. In: *Poultry Genetics, Breeding and Biotechnology* (Ed. W. M. Muir and S. E. Aggrey). CAB International, pp. 13-27.
- Fernández, V. and L. A. Videla. 1996. Effect of hyperthyroidism on the biliary release of thiobarbituric acid reactants in the rat. *Toxicol. Lett.* 84:149-153.
- Fernández, V., G. Tapia, P. Varela, P. Romanque, D. Cartier-Ugarte and L. A. Videla. 2005. Thyroid hormone-induced oxidative stress in rodents and humans, a comparative view and relation to redox regulation of gene expression. *Comp. Biochem. Physiol. C* 142:231-239.
- Ferreira, A. C., L. de Carvalho Cardoso, D. Rosenthal and D. P. de Carvalho. 2003. Thyroid Ca^{2+} /NADPH-dependent H_2O_2 generation is partially inhibited by propylthiouracil and methimazole. *Eur. J. Biochem.* 270:2363-2368.
- Gredilla, R., G. Barja and M. Lopez-Torres. 2001a. Thyroid hormone-induced oxidative damage on lipids, glutathione and DNA in the mouse heart. *Free Radic. Res.* 35:417-425.
- Gredilla, R., M. Lopez Torres, M. Portero-Otin, R. Pamplona and G. Barja. 2001b. Influence of hyper- and hypothyroidism on lipid peroxidation, unsaturation of phospholipids, glutathione system and oxidative damage to nuclear and mitochondria DNA in mice skeletal muscle. *Mol. Cell. Biochem.* 221:41-48.
- Guerrero, A., R. Pamplona, M. Portero-Otin, G. Barja and M. Lopez-Torres. 1999. Effect of thyroid status on lipid composition and peroxidation in the mouse liver. *Free Radic. Biol. Med.* 26:73-80.
- Han, B., S. S. Yoon, H. R. Han, W. J. Qu and F. Nigussie. 2005. Effect of low ambient temperature on the concentration of free radicals related to ascites in broiler chickens. *Asian-Aust. J. Anim. Sci.* 18:1182-1187.

- Iqbal, M., D. Cawthon, K. Beers, R. F. Jr Wideman and W. G. Bottje. 2002. Antioxidant enzyme activities and mitochondrial fatty acids in pulmonary hypertension syndrome (PHS) in broilers. *Poult. Sci.* 81:252-260.
- Jiang, S. Q., Z. Y. Jiang, Y. C. Lin, P. B. Xi and X. Y. Ma. 2007. Effects of soy isoflavone on performance, meat quality and antioxidative property of male broilers fed oxidized fish oil. *Asian-Aust. J. Anim. Sci.* 20:1252-1257.
- Lin, H., R. Du, X. H. Gu, F. C. Li and Z. Y. Zhang. 2000. A study on the plasma biochemical indices of heat stressed broilers. *Asian-Aust. J. Anim. Sci.* 13:1210-1218.
- Lin, H., R. Du and Z. Y. Zhang. 2000. Peroxide status in tissues of heat-stressed broilers. *Asian-Aust. J. Anim. Sci.* 13:1373-1376.
- Lin, H., E. Decuypere and J. Buyse. 2004a. Oxidative stress induced by corticosterone administration in broiler chickens (*Gallus gallus domesticus*) 1. Chronic exposure. *Comp. Biochem. Physiol. B* 139:737-744.
- Lin, H., E. Decuypere and J. Buyse. 2004b. Oxidative stress induced by corticosterone administration in broiler chickens (*Gallus gallus domesticus*) 2. Short-term effect. *Comp. Biochem. Physiol. B* 139:745-751.
- Lin, H., E. Decuypere and J. Buyse. 2006. Acute heat stress induces oxidative stress in broiler chickens. *Comp. Biochem. Physiol. A* 144:11-17.
- Luger, D., D. Shinder and S. Yahav. 2002. Hyper- or hypothyroidism: its association with the development of ascites syndrome in fast-growing chickens. *Gen. Comp. Endocrinol.* 127:293-299.
- Kim, H., T. Lee, Y. S. Hwang, M. A. Bang, K. H. Kim, J. M. Suh, H. K. Chung, D. Yu, Y. Lee, O. Kwon, H. K. Ro and M. Shong. 2001. Methimazole as an antioxidant and immunomodulator in thyroid cells: mechanisms involving interferon- γ signalling and H₂O₂ scavenging. *Mol. Pharmacol.* 60:972-980.
- Konukoglu, D., M. Ercan and H. Hatemi. 2002. Plasma viscosity in female patients with hypothyroidism: effects of oxidative stress and cholesterol. *Clin. Hemorheol. Microcirc.* 27:107-113.
- Mano, T., R. Sinohara, Y. Sawai, N. Oda, Y. Nishida, T. Mokuno, K. Asano, Y. Ito, M. Kotake and M. Hamada. 1995. Changes in lipid peroxidation and free radical scavengers in the brain of hyper- and hypothyroid aged rats. *J. Endocrinol.* 147:361-365.
- Pamplona, R., M. Portero-Otín, D. Riba, F. Ledo, R. Gredilla, A. Herrero and G. Barja. 1999 a. Heart fatty acid unsaturation and lipid peroxidation, and aging rate, are lower in the canary and the parakeet than in the mouse. *Aging (Milano)* 11:44-49.
- Pamplona, R., M. Portero-Otín, J. R. Requena, S. R. Thorpe, A. Herrero and G. Barja. 1999b. A low degree of fatty acid unsaturation leads to lower lipid peroxidation and lipoxidation-derived protein modification in heart mitochondria of the longevous pigeon than in the short-lived rat. *Mech. Ag. Develop.* 106:283-296.
- Resch, U., G. Helsel, F. Tatzber and H. Sinzinger. 2002. Antioxidant status in thyroid dysfunction. *Clin. Chem. Lab. Med.* 40:1132-1134.
- SAS Institute. 1998. User's Guide: Statistic. SAS Institute Inc., Cary, NC.
- Shim, K. S., K. T. Hwang, M. W. Son and G. H. Park. 2006. Lipid Metabolism and Peroxidation in Broiler Chicks under Chronic Heat Stress. *Asian-Aust. J. Anim. Sci.* 19:1206-1211.
- Stojevic, Z., S. Milinkovic-Tur and K. Curcija. 2000. Changes in thyroid hormones concentrations in chicken blood plasma during fattening. *Vet. Arhiv* 70:31-37.
- Venditti, P., K. Balestrieri, S. Di Meo and T. De Leo. 1997. Effect of thyroid state on lipid peroxidation, antioxidant defences, and susceptibility to oxidative stress in rat tissues. *J. Endocrinol.* 155:151-157.
- Venditti, P., R. De Rosa and S. Di Meo. 2003. Effect of thyroid state on susceptibility to oxidants and swelling of mitochondria from rat tissues. *Free Radic. Biol. Med.* 35:485-494.
- Venditti, P. and S. Di Meo. 2006. Thyroid hormone-induced oxidative stress. *Cell Mol. Life Sci.* 63:414-434.
- Zation, Z., Z. Merican, B. A. Khalid, J. B. Mohamed and S. Baharom. 1993. The effects of propranolol on skeletal muscle contraction, lipid peroxidation products and antioxidant activity in experimental hyperthyroidism. *Gen. Pharmacol.* 24: 195-199.