

## RESEARCH ARTICLE

# Differentiated Thyroid Carcinoma Risk Factors in French Polynesia

Constance Xhaard<sup>1,2,3</sup>, Yan Ren<sup>1,2,3</sup>, Enora Clero<sup>1,2,3</sup>, Stephane Maillard<sup>1,2,3</sup>, Pauline Brindel<sup>1,2,3</sup>, Frederique Rachedi<sup>4</sup>, Jean-Louis Boissin<sup>5</sup>, Joseph Sebbag<sup>5</sup>, Larrys Shan<sup>5</sup>, Frederique Bost-Bezeaud<sup>4</sup>, Patrick Petitdidier<sup>6</sup>, Vladimir Drozdovitch<sup>7,8</sup>, Françoise Doyon<sup>1,2,3</sup>, Carole Rubino<sup>1,2,3</sup>, Florent de Vathaire<sup>1,2,3\*</sup>

### Abstract

**Background:** To investigate differentiated thyroid cancer risk factors in natives of French Polynesia is of interest because of the very high incidence of this cancer in the archipelago. **Materials and Methods:** To assess the role of various potential risk factors of thyroid cancer in the natives of French Polynesia we performed a case-control study. The study included almost all the French Polynesians diagnosed with differentiated thyroid carcinoma between 1981 and 2003 (n=229) and 373 French Polynesian control individuals from the general population without cancer. **Results:** Thyroid radiation dose received from nuclear fallout before the age of 15, a personal history of neck or/and head medical irradiation, obesity, tallness, large number of children, an artificial menopause, a familial history of thyroid cancer, a low dietary iodine intake, and having a spring as the main source of drinking water were found to be significant risk factors. No roles of smoking habits, alcohol consumption, iodine containing drugs, and exposure to pesticides were evidenced. **Conclusions:** Except for smoking, differentiated thyroid carcinoma risk factors in natives of French Polynesia are similar to those in other populations. Our finding on the role of having a spring as a drinking water origin is coherent with some other studies and could be due to geological factors.

**Keywords:** Differentiated thyroid carcinoma - nuclear test - French Polynesia - radiation induced cancer

*Asian Pac J Cancer Prev*, 15 (6), 2675-2680

### Introduction

Substantial variations in thyroid cancer incidence in the world strongly implicate environmental factors in the etiology of this cancer. The highest thyroid cancer incidence rates are observed in the Pacific populations (Henderson et al., 1982; Curado et al., 2007), but only few publications have investigated the reasons for these high rates.

Thyroid cancer mortality (de Vathaire and Le Vu, 1996) and incidence (Le Vu et al., 2000; de Vathaire et al., 2000) in French Polynesia was reported to be the highest in the world, in particular among natives of this set of archipelago (Gleize et al., 2000). To investigate the reasons for this level of incidence and the potential role of atmospheric fallout from nuclear tests performed by France between 1966 and 1974, we conducted between 2002 and 2005 a population-based case-control study among natives of French Polynesia including most of the thyroid cancer patients diagnosed between 1981 and 2003,

who were aged 55 years or less and still alive. Previously, we reported that a low level of education (Brindel et al., 2008), tallness and a high body mass index (Brindel et al., 2009), a family history of thyroid cancer (Brindel et al., 2010), and, in women, a large number of pregnancies (Brindel et al., 2008) were associated with an increased risk of thyroid cancer in this population. We also estimated the radiation doses received by the study subjects during the atmospheric nuclear tests (Drozdovitch et al., 2008) and concluded that nuclear radiation fallout played a small but significant role in the high incidence of thyroid cancer in French Polynesia (de Vathaire et al., 2010). Lastly, we estimated that traditional Polynesian dietary pattern was associated with a lower thyroid cancer incidence than western pattern (Clero et al., 2012a). French Polynesia is a mild iodine deficiency area in which a higher consumption of food from the sea and a higher dietary iodine intake are significantly associated with a decreased risk of thyroid cancer (Clero et al., 2012b).

In this report we investigated the potential risk factors

<sup>1</sup>Inserm, Centre for Research in Epidemiology and Population Health (CESP), U1018, Radiation Epidemiology Group, <sup>2</sup>Gustave Roussy, <sup>3</sup>University Paris XI, Villejuif, <sup>7</sup>International Agency for Research on Cancer, Lyon, France, <sup>4</sup>Territorial Hospital Taaone, <sup>5</sup>Endocrinologist, <sup>6</sup>Laboratoire Boz, Papeete, French Polynesia, <sup>8</sup>Division of Cancer Epidemiology and Genetics, National Cancer Institute, NIH, DHHS, Bethesda, USA \*For correspondence: [florent.devathaire@gustaveroussy.fr](mailto:florent.devathaire@gustaveroussy.fr)

that have not been previously reported.

## Materials and Methods

### Study population

Between 1980 and 1984 two partial registries maintained by clinicians and pathologists were in operation. The cancer incidence registry of French Polynesia was officially created in 1984. Between 1996 and 1998, we investigated the exhaustiveness of the registry and identified 539 new cancers after having examined all the medical transport files, insurance records, and hospital and pathological laboratory files available in French Polynesia. An additional survey limited to thyroid cancer was also performed in 2002–2005, to complete registry data with incident cases diagnosed up to 2002 in data from medical insurance files, and records from the four endocrinologists in Tahiti.

The methodology and practical realization of the case-control study can be found elsewhere (de Vathaire et al., 2010). Briefly, 229 of the of 255 patients diagnosed with differentiated thyroid carcinoma between 1981 and 2003 at the age of 55 or less, and born and residing in French Polynesia at time of diagnosis, as well as 373 population controls matched on sex and birth date extracted from the French Polynesian birth registry were included in the study. Due to errors in inclusion, some patients aged between 56 and 62 were included and kept in this analysis. The study subjects were interviewed face to face by trained Maori interviewers. Information on ethnic group, education, occupation, places of residence, weight history, personal and familial history of thyroid disease and cancer, gynecologic and reproductive history, medical X-ray exposure, and diet at the time of the interview and in childhood was collected during the personal interview.

**Table 1. Description of Thyroid Cancer Cases, French Polynesia**

Characteristics	Cases (N=229)	
	N	%
Sex	Men	26 11.4
	Women	203 88.6
Age at time of first test (years)	15+	61 26.6
	0-14	124 54.2
	Born during tests	33 14.4
	Born after tests	11 4.8
Age at diagnosis (years)	10-19	8 3.5
	20-29	39 17
	30-39	76 33.2
	40-49	63 27.5
	50-62	43 18.8
Histology	Papillary	176 76.9
	Follicular	53 23.1
Tumour size (mm)	≤ 10	106 52.2
	11-40	75 37.0
	>40	22 10.8
	Unknown	26 -
Multifocal tumour	No	128 61
	Yes	82 39
	Unknown	19 -
Extra-thyroidal invasion	No	191 83.4
	Yes	38 16.6

Table 1 describes the 229 cases included in the study, and Table 2 summarizes the main findings previously published.

### Estimation of radiation thyroid doses

Radiation doses to the thyroids of the study subjects were assessed based on available annual reports published by France on the radiological situation in French Polynesia, including measurements of the radioactivity in the air, vegetables, milk, and fish, and on meteorological data including daily precipitation and wind speed and direction. Dietary consumption data concerning childhood and information on the source of drinking water collected during the personal interviews were also used to estimate individual radiation doses to the thyroid. Dose reconstruction was conducted without knowledge of the case or control status of the subject. The methodology used to estimate thyroid doses to the study subjects and results can be found elsewhere (Drozdovitch et al., 2008).

### Statistical analysis

Data were analyzed using conditional logistic regression (Breslow and Day, 1987) with the Epicure epidemiological software (Preston et al., 1993) and SAS® V9.2. To investigate the relationship between the radiation dose to the thyroid and the risk of thyroid cancer, we compared nested models using likelihood-ratio tests (Moolgavkar and Venzon, 1987). Tests for linear trend (Breslow and Day, 1987) were also performed.

## Results

Only 17 study subjects reported use of traditional Polynesian medicine, a practice that did not significantly modify the risk of thyroid cancer. The use of Colchimax or Amiodarone, which are drugs containing iodine, did not significantly modify the risk of thyroid carcinoma. Only 11 cases and 18 controls were or had been farmers. This

**Table 3. Risk factors for Thyroid Cancer in French Polynesia**

	Cases (n=229)		Controls (n=373)		OR (95%CI)*	p value**
	No.	%	No.	%		
Traditional Polynesian medicine (Raau Tahiti).						
No	222	97	363	97	1*	
Yes	7	3	10	3	0.6 (0.2-1.9)	0.4
Colchimax or Amiodarone (drug containing iodine)						
No	212	93	342	92	1*	
Yes	17	7	31	8	0.9 (0.4-1.8)	0.8
Farmer as occupation						
No	218	95	355	95	1*	
Yes	11	5	18	5	1.0 (0.4-2.3)	1
Domestic or professional use of pesticides						
No	127	55	163	44	1*	
Yes	102	45	210	56	1.0 (0.6-1.5)	1
Report of regular alcohol consumption						
No	222	97	361	97	1*	
Yes	7	3	10	3	1.2 (0.3-4.5)	0.8
Current smoker						
No	101	44	197	53	1*	
Yes	128	56	176	47	1.1 (0.8-1.7)	0.8

\*OR: odds ratio, CI: confidence interval; \*\*Multivariate conditional logistic regression, taking into account all other factors in Table 2

**Table 2. Risk Factors for Thyroid Cancer in French Polynesia**

		Cases (n=229)		Controls (n=373)		OR (95%CI)*	p value**
		No.	%	No.	%		
Maximal Body Mass Index (kg/m <sup>2</sup> )	≤ 22.5	12	5	49	13	1*	<0.0005
	22.6-26.3	36	16	91	24	1.1 (0.5-2.5)	
	26.4-31.6	62	27	103	28	1.8 (0.8-3.9)	
	≥ 31.7	119	52	130	35	2.8 (1.3-5.8)	
Height (m) (Female) (Males)	(<1.60) (<1.70)	32	14	80	22	1*	<0.05
	(1.60-1.64) (1.70-1.74)	64	28	101	27	1.6 (0.9-2.8)	
	(1.65-1.69) (1.75-1.79)	72	31	103	28	1.7 (1.0-3.1)	
	(≥1.70) (≥1.80)	61	27	89	24	1.9 (1.0-3.4)	
Women: number of pregnancies	0	25	12	49	15	1*	<0.05
	1-Feb	38	19	84	26	1.2 (0.5-2.7)	
	3-Apr	53	26	84	26	1.5 (0.6-3.3)	
	5-Jun	35	17	48	15	1.9 (0.8-4.5)	
	7 or more	52	26	61	19	2.3 (1.0-5.2)	
Women: menopausal status	Premenstrual	1	1	2	1	1*	0.02
	Still menstruated	161	81	277	87	1*	
	Natural menopause	17	8	26	8	1.9 (0.7-5.0)	
	Artificial menopause	19	10	11	4	4.5 (1.5-12.0)	
Nuclear worker during atmospheric test	No	219	96	361	97	1*	0.9
	Yes	10	4	12	3	1.0 (0.3-3.3)	
Atmospheric tests nuclear radiation fallout : thyroid dose before age 15	<1 mGy	156	68	250	67	1*	<0.05
	1-9.9 mGy	61	27	111	30	1.0 (0.5-2.0)	
	10-19.9 mGy	7	3	9	2	3.9 (0.9-17)	
	20-39 mGy	5	2	3	1	7.0 (0.9-53)	
First degree familial history of thyroid cancer	No	205	90	362	97	1*	0.01
	Yes	24	10	11	3	4.5 (1.9-10.6)	
Total food from sea consumption (g/day)	≤45	112	49	136	37	1*	0.0002
	46-90	58	25	115	31	0.5 (0.3-0.9)	
	≥91	59	26	120	32	0.4 (0.3-0.8)	
Parents ethnic group	Both Polynesian	122	53	206	55	1*	0.6 0.9 0.08
	Poly-Asian	37	16	63	17	1.3 (0.8-2.3)	
	Poly-European	46	20	75	20	1.3 (0.7-2.2)	
	Mixed	23	10	29	8	2.2 (1.1-4.5)	
	Missing	1					
Educational level	No diploma	101	44	138	37	1*	<0.02
	Primary leaving certificate	53	23	84	23	0.9 (0.5-1.4)	
	Middle school degree	24	11	31	8	1.5 (0.8-3.3)	
	Vocational training certificate	32	14	49	13	0.8 (0.4-1.5)	
	Technical school certificate	10	4	18	5	0.7 (0.3-1.8)	
	High school diploma	9	4	53	14	0.2 (0.1-0.6)	
Medical or therapeutic irradiation of the neck	No	191	83	331	89	1*	0.02
	Yes	38	17	42	11	1.9 (1.1-3.4)	

\*OR: odds ratio, CI: confidence interval; \*\*Multivariate conditional logistic regression, taking into account all other factors in the Table

**Table 4. Origin of Drinking Water and Thyroid Cancer in French Polynesia**

		Cases (n=229)		Controls (n=373)		OR (95%CI)*	p value**
		No.	%	No.	%		
Spring as main drinking water origin	No	179	78	321	86	1*	0.03 0.9 0.1 0.03 0.6 0.3
	Yes	50	22	52	14	1.7 (1.1-2.9)	
	Yes in Tahiti or Moorea	4	2	2	2	1.1 (0.3-4.5)	
	Yes in other Société islands	5	2	3	1	3.8 (0.3-4.5)	
	Yes in Australes	16	7	13	3	2.7 (1.1-6.6)	
	Yes in Marquises	23	10	29	8	1.2 (0.6-2.7)	
	Yes in atolls of Tuamotu-Gambier	4	1	2	2	2.4 (0.4-13.1)	
Spring as main drinking water origin before the age of 15	No	192	84	339	91	1*	0.003
	Yes	37	16	34	9	2.5 (1.4-4.7)	
Personal cistern as main drinking water origin	No	188	82	315	84	1*	0.9
	Yes	41	18	58	16	1.0 (0.6-1.7)	
Public cistern as main drinking water origin	No	215	94	354	95	1*	0.9
	Yes	14	6	19	5	1.0 (0.5-2.1)	

\*OR: odds ratio, CI: confidence interval; \*\*Multivariate conditional logistic regression, taking into account all other factors listed in Table 2

occupation did not significantly modify the risk of thyroid carcinoma. Reported domestic occupational exposure to pesticides was not found to be significantly linked to thyroid carcinoma risk (Table 3).

Only 7 cases and 10 controls reported regularly

drinking alcohol. We did not find evidence of any significant relation between reported alcohol consumption or smoking habits and thyroid carcinoma risk (Table 3).

Fifty cases and 52 controls declared having or having had a spring as main source of drinking water in at least

one of their places of residence, mainly in Australes and Marquises Islands. Having or having had a spring as main origin of drinking water was associated with a significantly increased risk of thyroid cancer, in particular at age less than 15 years old. Among 102 subjects who have had a spring as a main source of drinking water, 3 (2.9%) had received 10 mGy or more to the thyroid during atmospheric tests, as compared with 21 (4.2%) of the 500 subjects who never had such source of drinking water. Adjusting for spring as origin of drinking water did not modify the relation between thyroid radiation dose estimate and thyroid cancer risk. No relation between other sources of water, public or private rainwater cisterns, and thyroid cancer risk was found (Table 4).

## Discussion

This case-control study aimed to identify lifestyle and environmental risk factors, specific to French Polynesia, which could explain the high differentiated thyroid carcinoma incidence observed in these archipelagos. The study identifies having a spring as origin of drinking water as a risk factor, but the low frequency of this risk factor (14% in controls and 22% in cases) does not permit it to play an important role, its population attributable fraction being about 9%. On the other hand, we failed to evidence the role of smoking habits, alcohol consumption, iodine containing drugs, and exposure to pesticides.

The description of general strengths and weaknesses of this case-control study can be found elsewhere (Brindel et al., 2008; de Vathaire et al., 2010; Clero et al., 2012b). In brief, the weaknesses of this study are the traditional ones of case-control studies in general population in which exposure assessment is based on self-reported data, i.e. potential bias of anamnesis and uncertainties in exposure assessment. Its strengths are those arising from the small size of the French Polynesian population (it is still possible to localize the current living place of any person when knowing only name, first name, and birth date) and its geographic isolation up to the 1990s (traditional Polynesian food was the rule in all Polynesian islands except Tahiti, until very recently).

As regards the potential risk factors specifically investigated in this article, the most important weakness concerns the weight of religious prohibitions and is illustrated by the unrealistically low (3%) proportion of cases and controls who declared drinking alcohol, as compared with what is known about alcoholic epidemics in French Polynesia (Bertrand and Berry, 2013). This strong limitation means that our study is unable to investigate alcohol role, which in fact may be inhibitory (de Menezes et al., 2013). Similar limitation probably does not exist for smoking habits, the self-reported current smoker proportion in controls (47%) being similar to what is expected in a French Polynesian population of the same age (Rasanathan and Tukuitonga, 2007), and an over-reporting of tobacco in cases being improbable. Nevertheless, smoking has been seen as negatively associated with thyroid carcinoma incidence in several studies (Hershman, 2012; Kabat et al., 2012; Kitahara et al., 2012; Zivaljevic et al., 2013), and additional data are

needed to interpret the lack of association in our study (OR=1.1, 95%CI: 0.8 to 1.7) as a specificity of the French Polynesia population, rather than being due to hazard or to low power.

An interesting result of this study is the association between the effect of having or having had a private spring, rather than private or public cisterns collecting rainwater or public drinking water distribution networks, as main drinking water origin, and the risk of developing thyroid cancer. This result is coherent with recent results concerning fertilizers and incidence of cancer and exposure to fertilizer (Ward, 2009), and in particular with those of a cohort study which evidenced an increased risk of developing thyroid cancer in subjects having artesian well as drinking water origin (Ward et al., 2010). Nitrate concentration could be higher in water from artesian wells and personal untreated springs than in water from public water networks. Another explanation could be the occurrence of metals or other elements in water from artesian wells and personal springs. Indeed, the high incidence of thyroid cancer observed in islands, in particular in volcanic areas of some islands such as Sicily (Pellegriti et al., 2009), has been associated with a high concentration of metals in the drinking water (Malandrino et al., 2013). If such elements are in higher concentrations in artesian wells and personal spring water than in the public water networks, these findings could explain our results.

Thyroid cancer risk factors in French Polynesia were similar to those observed in other populations of the Asia and Pacific area. Some discrepancies, nevertheless, have to be noted.

The most controverted putative differentiated thyroid cancer risk factor is parity. Despite two case-control studies performed in China (Preston-Martin et al., 1993) and New-Caledonia (Truong et al., 2005) evidenced, as we did, an increased risk in women who were ever-pregnant, two cohort studies, performed in China (Wong et al., 2006) and Japan (Pham et al., 2009), did not evidence such an increased risk, and another performed in Thailand, although based on 17 cases showed a decreased risk (Sungwalee et al., 2013). Results in other populations are heterogeneous (Peterson et al., 2012) and this heterogeneity does not seem to be attributable only to differences in number of children between studied populations. As an example, no relation between number of pregnancies and risk of thyroid cancer was found in a large scale cohort study in which each woman had on average more than three children (Schonfeld et al., 2011). This heterogeneity remains to be explored and was observed both between recent studies (Schonfeld et al., 2011; Peterson et al., 2012) and between studies performed more than 20 years ago (La Vecchia et al., 1999). An explanation of this heterogeneity could come from cofactors. Indeed, the need for additional thyroid hormones during pregnancy leads to a multiplication of thyroid cells (Glinoe et al., 1990; Manole et al., 2001; Sack, 2013), and this multiplication could act by promoting carcinogens involved in cancer initiation. In our study, a positive significant ( $p=0.03$ ) interaction was evidenced between nuclear fallout exposure and

pregnancy number: the increase in thyroid cancer risk when increasing thyroid radiation dose was limited to women having had four pregnancies or more (de Vathaire et al., 2010).

In comparison, body size, i.e., BMI (Renehan et al., 2008) and tallness (Engeland et al., 2006; Rinaldi et al., 2012), are established for thyroid cancer, and have been confirmed in Asian populations (Dal Maso et al., 2000; Suzuki et al., 2008; Kim et al., 2012).

Except for smoking, differentiated thyroid carcinoma risk factors in natives of French Polynesia are similar to those in other populations. Our finding on the role of having a spring as a drinking water origin is coherent with some other studies and could be due to geological factors.

## Acknowledgements

This study was supported by the Association pour la Recherche contre le Cancer, the Ligue Nationale Contre le Cancer, the Direction Generale de la Sante, the Comite de radioprotection de Electricite de France, Agence Française de Securite Sanitaire et Environnementale et du Travail and CHILD-THYR EEC programme. The authors thank J Paoafaite, J Teuri, J Iltis from the Institute of Research for Development (IRD) and Drs Ph Morales, P Giraud, P Didiergeorge, M Brisard, G Soubiran, A Merceron, ML Vanizette, P Dupire, M Berges, J Ienfa, G de Clermont, N Cerf, B Oddo, M Bambridge, C Baron, A Mouchard-Rachet, O Simonet, D Lamarque, A Vabret, J Delacre, MP Darquier, and J Leninger for their help.

## References

- Bertrand S, Berry AL (2013). Health survey 2010 in French Polynesia: surveillance of risk factors for non-communicable diseases. *Bulletin Epidemiologique Hebdomadaire*, **28-29**, 326-32.
- Breslow NE, Day NE (1987). Fitting models to grouped data. In: *Statistical Methods in Cancer Research: The Design and Analysis of Cohort Studies*. IARC scientific publication No. 82, IARC: Lyon, France, 119-76.
- Brindel P, Doyon F, Rachedi F, et al (2008). Menstrual and reproductive factors in the risk of differentiated thyroid carcinoma in native women in French Polynesia: a population-based case-control study. *Am J Epidemiol*, **167**, 219-29.
- Brindel P, Doyon F, Rachedi F, et al (2009). Anthropometric factors in differentiated thyroid cancer in French Polynesia: a case-control study. *Cancer Causes Control*, **20**, 581-90.
- Brindel P, Doyon F, Bourgain C, et al (2010). Family history of thyroid cancer and the risk of differentiated thyroid cancer in French Polynesia. *Thyroid*, **20**, 393-400.
- Clero E, Doyon F, Chungue V, et al (2012a). Dietary patterns, goitrogenic food, and thyroid cancer: A case-control study in French Polynesia. *Nutr Cancer*, **64**, 929-36.
- Clero E, Doyon F, Chungue V, et al (2012b). Dietary iodine and thyroid cancer risk in French Polynesia: A case-control study. *Thyroid*, **22**, 422-9
- Curado MP, Edwards B, Shin HR, et al (2007). Cancer Incidence in Five continents. Volume XI. IARC Scientific Publication No. 160, IARC: Lyon, France.
- Dal Maso L, La Vecchia C, Franceschi S, et al (2000). A pooled analysis of thyroid cancer studies. V. Anthropometric factors. *Cancer Causes Control*, **11**, 137-44.
- de Menezes RF, Bergmann A, Thuler LC (2013). Alcohol consumption and risk of cancer: a systematic literature review. *Asian Pac J Cancer Prev*, **14**, 4965-72.
- de Vathaire F, Le Vu B (1996). Cancer mortality in French Polynesia from 1984 to 1992. *Br J Cancer*, **74**, 1680-1.
- de Vathaire F, Le Vu B, Challeton de Vathaire C (2000). Thyroid cancer in French Polynesia between 1985 and 1995: influence of atmospheric nuclear bomb tests performed at Mururoa and Fangataufa between 1966 and 1974. *Cancer Causes Control*, **11**, 59-63.
- de Vathaire F, Drozdovitch V, Brindel P, et al (2010). Thyroid cancer following nuclear tests in French Polynesia. *Br J Cancer*, **103**, 1115-21.
- Drozdovitch V, Bouville A, Doyon F, et al (2008). Reconstruction of individual radiation doses for a case-control study of thyroid cancer in French Polynesia. *Health Physics*, **94**, 418-33.
- Engeland A, Tretli S, Akslen LA, et al (2006). Body size and thyroid cancer in two million Norwegian men and women. *Br J Cancer*, **95**, 366-70.
- Gleize L, Laudon F, Yen Kai Sun L, et al (2000). Cancer registry in French Polynesia: results for the 1990-1995 period for native and immigrant populations. *Eur J Epidemiol*, **16**, 661-7.
- Glinoeur D, de Nayer P, Bourdoux P, et al (1990). Regulation of maternal thyroid during pregnancy. *J Clin Endocrinol Metab*, **71**, 276-87.
- Henderson BE, Forter F, Kolonel LN (1982). Cancer in Polynesians. *NCI Monogr*, **62**, 73-8.
- Hershman JM (2012). Smoking may decrease the incidence of thyroid cancer in postmenopausal women. *Clin Thyroidol*, **24**, 8-9.
- Kabat GC, Kim MY, Wactawski-Wende J, et al (2012). Smoking and alcohol consumption in relation to risk of thyroid cancer in postmenopausal women. *Cancer Epidemiol*, **36**, 335-40.
- Kim JY, Jung EJ, Park ST, et al (2012). Body size and thyroid nodules in healthy Korean population. *J Korean Surg Soc*, **82**, 13-7.
- Kitahara CM, Linet MS, Beane LE, et al (2012). Cigarette smoking, alcohol intake, and thyroid cancer risk: a pooled analysis of five prospective studies in the United States. *Cancer Causes Control*, **23**, 1615-24.
- La Vecchia C, Ron E, Franceschi S, et al (1999). A pooled analysis of case-control studies of thyroid cancer. III. Oral contraceptives, menopausal replacement therapy and other female hormones. *Cancer Causes Control*, **10**, 157-66.
- Le Vu B, de Vathaire F, Challeton de Vathaire C, et al (2000). Cancer incidence in French Polynesia between 1985 and 1995. *Trop Med Int Health*, **5**, 722-31.
- Malandrino P, Scollo C, Marturano I, et al (2013). Front Endocrinol (Lausanne). Descriptive epidemiology of human thyroid cancer: experience from a regional registry and the "volcanic factor". *Frontiers Endocr*, **4**, 1-7.
- Manole D, Schildknecht B, Gosnell B, et al (2001). Estrogen promotes growth of human thyroid tumor cells by different molecular mechanisms. *J Clin Endocrinol Metab*, **86**, 1072-7.
- Moolgavkar SH, Venzon DJ (1987). A method for computing profile likelihood based confidence bounds. *Ann Stat*, **15**, 346-59.
- Pellegriti G, De Vathaire F, Scollo C, et al (2009). Papillary thyroid cancer incidence in the volcanic area of Sicily. *J Natl Cancer Inst*, **101**, 1575-83.
- Peterson E, De P, Nuttall R (2012). BMI, diet and female reproductive factors as risks for thyroid cancer: a systematic review. *PLoS One*, **7**, 29177.
- Pham TM, Fujino Y, Mikami H, et al (2009). Japan Collaborative

- Cohort Study Group. Reproductive and menstrual factors and thyroid cancer among Japanese women: the Japan Collaborative Cohort Study. *J Womens Health*, **18**, 331-5.
- Preston DL, Lubin JH, Pierce DA, et al (1993). EPICURE User's Guide. Hirosoft International Corp: Seattle, Washington.
- Preston-Martin S, Jin F, Duda MJ, et al (1993). A case-control study of thyroid cancer in women under age 55 in Shanghai (People's Republic of China). *Cancer Causes Control*, **4**, 431-40.
- Rasanathan K, Tukuitonga CF (2007). Tobacco smoking prevalence in Pacific Island countries and territories: a review. *N Z Med J*, **120**, 2742.
- Rehman AG, Tyson M, Egger M, et al (2008). Body-mass index and incidence of cancer: a systematic review and meta-analysis of prospective observational studies. *Lancet*, **371**, 569-78.
- Rinaldi S, Lise M, Clavel-Chapelon F (2012). Body size and risk of differentiated thyroid carcinomas: findings from the EPIC study. *Int J Cancer*, **131**, 1004-14.
- Sack J (2013). Thyroid function in pregnancy - maternal-fetal relationship in health and disease. *Pediatr Endocrinol Rev*, **1**, 170-6.
- Schonfeld SJ, Ron E, Kitahara CM, et al (2011). Hormonal and reproductive factors and risk of postmenopausal thyroid cancer in the NIH-AARP Diet and Health Study. *Cancer Epidemiol*, **35**, 85-90.
- Sungwalee W, Vatanasapt P, Kamsa-Ard S, et al (2013). Reproductive risk factors for thyroid cancer: a prospective cohort study in Khon Kaen, Thailand. *Asian Pac J Cancer Prev*, **14**, 5153-5.
- Suzuki T, Matsuo K, Hasegawa Y, et al (2008). Anthropometric factors at age 20 years and risk of thyroid cancer. *Cancer Causes Control*, **19**, 1233-42.
- Truong T, Orsi L, Dubourdieu D, et al (2005). Role of goiter and of menstrual and reproductive factors in thyroid cancer: a population-based case-control study in New Caledonia (South Pacific), a very high incidence area. *Am J Epidemiol*, **161**, 1056-65.
- Ward MH (2009). Too much of a good thing? Nitrate from nitrogen fertilizers and cancer. *Rev Environ Health*, **24**, 357-63.
- Ward MH, Kilfoy BA, Weyer PJ, Anderson KE, et al (2010). Nitrate intake and the risk of thyroid cancer and thyroid disease. *Epidemiology*, **21**, 389-95.
- Wong EY, Ray R, Gao DL, et al (2006). Reproductive history, occupational exposures, and thyroid cancer risk among women textile workers in Shanghai, China. *Int Arch Occup Environ Health*, **79**, 251-8.
- Zivaljevic V, Slijepcevic N, Sipetic S, et al (2013). Risk factors for well-differentiated thyroid cancer in men. *Tumori*, **99**, 458-62.