

CCD reduction package, we obtained Color-magnitude Diagrams(CMDs) with well defined main-sequences extending to $V=23(M71)$ and $V=25(M13)$, respectively. In order to investigate the age spread among globular cluster, we will compare these CMDs with those of 47 Tuc and M3.

Washington CCD Photometry of Globular Cluster in NGC 4472

Myung Gyoon Lee, Eun Hyeuk Kim (Seoul National Univ.)M

Doug Geisler(Kitt Peak National Observatory)

Globular cluster in giant elliptical galaxies provide us with important clues for understanding the formation and evolution of globular cluster as well as their parent galaxies. NGC 4472 (M49) is the brightest elliptical galaxy in the Virgo cluster and has a rich globular cluster system.

We present a study of the metallicity of a large sample of globular cluster in NGC 4472. Deep Washington CT_1 photometry of a wide ($16' \times 16'$) field of NGC 4472 was obtained using Tek 2048 \times 2048 CCD at the KPNO 4m telescope.

The color-magnitude diagram (T_1 vs $(C-T_1)$) of $\sim 9,500$ measured point sources show 1) two strong vertical structures in the color range of $1.0 < (C-T_1) < 2.3$ which consist mostly of globular cluster, and 2) a dominant horizontal structure fainter than $T_1=23$ mag most of which are unresolved faint background galaxies.

We have estimated the metallicity of $\sim 1,400$ globular cluster brighter than $T_1=22.5$ mag from the $(C-T_1)$ colors. The metallicity distribution of the bright globular cluster shows two strong peaks at $[Fe/H]=-1.3$ dex, and possibly two peaks at $[Fe/H]=+0.3$ and $+0.9$ dex. The metal-rich globular cluster are spatially more concentrated than the metal-poor globular cluster. The mean metallicity of the globular cluster is decreasing as the galactocentric radius increase, similarly to that of M87. These results are consistent with the merger hypothesis for the formation of giant elliptical galaxies.

The luminosity function of giant of the globular cluster shows clearly a peak at $T_1=23.3 \pm 0.1$, from which we derive a distance modulus of $(m-M)_0=31.2 \pm 0.2$ mag ($d=17.4 \pm 1.6$ Mpc). Using this value for the distance, we estimate the Hubble constant to be $H_0=66 \pm 14$ km/s Mpc for the mean velocity of the Virgo $v=1179 \pm 17$ km/s (Sandage & Tammann 1993) and $H_0=79 \pm 17$ km/s Mpc for the velocity $v=1404 \pm 80$ km/s (Huchra 1988).