

SI-4-1

Topology of the Regulatory Protein-Protein Interaction for *Escherichia coli* Diauxism

Yeong-Jae Seok

Department of Microbiology, Seoul National University

Sugar transport by some permeases in *E. coli* is allosterically regulated by the phosphorylation state of enzyme IIA^{glc} (EIIA^{glc}) of the phosphoenolpyruvate:sugar phosphotransferase system. The binding is stimulated by transportable substrates (lactose, melibiose, and raffinose), but not by sugars that are not transported (maltose and sucrose). Treatment of lactose permease with *N*-ethylmaleimide (NEM), which blocks ligand binding and transport by alkylating Cys-148, also blocks ligand binding. Preincubation with the substrate analog β -D-galactopyranosyl 1-thio- β -D-galactopyranoside protects both lactose transport and EIIA^{glc} binding against inhibition by NEM. A collection of lactose permease replacement mutants at Cys-148 showed, with the exception of C148V, a good correlation of relative transport activity and EIIA^{glc} binding. The nature of the interaction of EIIA^{glc} with the cytoplasmic face of lactose permease was explored. Lactose permease mutants with polyhistidine insertions in cytoplasmic loops IV/V and VI/VII and periplasmic loop VII/VIII retain transport activity, but do not bind EIIA^{glc}, indicating that these regions of lactose permease may be involved in recognition of EIIA^{glc}. Taken together, these results suggest that interaction of lactose permease with substrate promotes a conformational change that brings several cytoplasmic loops into an arrangement optimal for interaction with the regulatory protein, EIIA^{glc}. A topological map of the proposed interaction is presented.

SI-4-2

Properties of the Flavoenzyme Pyranose Oxidase from *Oxyporus* sp.

Jaeyul Kwon and Sa-Ouk Kang *

Department of Microbiology, College of Natural Sciences, and
Research Center for Molecular Microbiology, Seoul National
University

Pyranose oxidase contained 4 mol of covalently bound FAD per mol of tetrameric protein. The FAD moiety was shown to be covalently linked via the $\delta\alpha$ -methylene group to the *N*(1) position of the imidazole ring of histidine. The flavin portion of pyranose oxidase underwent characteristic spectral changes upon changing the pH. The pK_a for the spectral change of the first absorption band was 5.66. Aromatic carboxylates such as 2-hydroxybenzoate and *m*-hydroxybenzoate showed a marked inhibition effect on the enzyme activity by the formation of complex with the active site. The pH dependence of the absorption spectral changes suggested that an ionizable group of a pK_a of 6.4 is involved in formation of complex with 2-hydroxybenzoate. The ionizable group having the same pK_a value was shown to be involved in formation of complex with acetate which had some inhibition effect. The reaction product for D-glucose, the most preferred substrate, was D-arabino-hexos-2-ulose. Pyranose oxidase oxidized certain aldonic acid lactones such as L-gulonono-1,4-lactone and D-arabinono-1,4-lactone to produce L-ascorbic acid and D-erythroascorbic acid respectively.