

## Angular Dependent Rebalance Factor Method for Solving the $S_N$ Transport Equations in X-Y Geometry

Ser Gi Hong and Nam Zin Cho

Korea Advanced Institute of Science and Technology  
373-1 Kusong-dong, Yusong-gu, Taejon, Korea 305-701

### Abstract

*The angular dependent rebalance (ADR) factor method is extended to the two-dimensional neutron transport problems and applied to acceleration of three nodal transport methods (i.e., constant-linear (C-L), simplified linear-discontinuous (SLD), and bilinear discontinuous (BLD)) and three non-nodal spatial differencing schemes (i.e., step scheme (SS), step characteristic (SC), and constant-constant (C-C)) where the spatial distributions of the angular flux are assumed to be flat in mesh interior and mesh edges. In ADR, the rebalance factor is defined as the ratio of the angular fluxes of the same phase point in two iterates and it is angular dependent. In this paper, the  $S_2$  approximation for the angular dependency of the rebalance factor is used. The resulting  $S_2$ -like lower-order equation is solved by three iterative methods (four-cyclic (FC), conjugate gradient (CG), and bi-conjugate gradient stabilized (Bi-CGSTAB) methods). The numerical tests show that the ADR method can be used effectively for all tested transport methods.*

---

Proceedings of the Korean Nuclear Society Autumn Meeting  
Seoul, Korea, October 1998

### Asymptotic Derivation of the Multigroup Modified Time-Dependent Simplified $P_2$ Equations

Uncheol Shin

Korea Atomic Energy Research Institute  
P. O. Box 105, Yusung, Taejon 305-600 Korea

### Abstract

*The multigroup modified time-dependent simplified  $P_2$  (SP2) equations are derived as the higher order asymptotic approximations to the multigroup time-dependent transport equation in a physical regime in which the conventional multigroup time-dependent diffusion equations are the leading-order asymptotic approximation. The derivation is performed in general three-dimensional geometry.*