

## 구두발표 초록

### O-01. Identification of *Bipolaris setariae*, *Curvularia ergrostidis*, *Phoma glomerata* and *Exserohilum inaequale* Isolated from the Seeds of *Arundinella hirta*. Soon Min Hong<sup>1</sup>, Seo Yeon Chang, Sang Jin

Han<sup>2</sup> and Hee Kyu Kim<sup>3</sup>. <sup>1</sup>Pusan Branch Office, National Plant Quarantine Service, 10-4, 6, Choongang Dong, Pusan, Korea, 600-016. <sup>2</sup>National Plant Quarantine Service, 433-1, Anyang 6 Dong, Anyang, Korea 430-016. <sup>3</sup>College of Agriculture, Gyeongsang National University, Chinju, Korea 660-701.

*Curvularia ergrostidis*, *Bipolaris setariae*, *Phoma glomerata* and *Exserohilum inaequale* were isolated from 400 seed samples of *Arundinella hirta* Tanaka from China. *C. ergrostidis* was the predominant species in the samples followed by *B. setariae*, *P. glomerata*, and *E. inaequale*. The morphological characteristics of the fungi were as follows; *C. ergrostidis* was 3-distoseptate, ellipsoidal or barrel-shaped, the middle septum almost median appearing as a black band, central cells brown to dark brown, end cell paler, rather smooth, and  $18-37 \times 11-20\mu\text{m}$  in size. The conidiophore was solitary or in group, simple or rarely branched, straight or curved, multi septate, and variable in length; The conidia of *B. setariae* was slightly curved, rarely straight, fusoid or navicular, pale to mid golden brown, smooth, 5-10 -distoseptate, and  $50-70 \times 10-15\mu\text{m}$  in size. The conidiophore was single or in small group, straight to flexuous, septate, smooth, pale to mid brown or olivaceous brown, paler towards the apex, up to  $200\mu\text{m}$  long, and  $5-9\mu\text{m}$  thick; *P. glomerata* produced dictyochlamydospores commonly arising in unbranched or branched chains of 2-20, resembling the conidia of *Alternaria*, usually  $18-20 \times 12-30\mu\text{m}$  in size. Pycnidia was mostly regularly globose and the conidia was ellipsoidal, mostly  $6-7 \times 3-4\mu\text{m}$  in size; The conidia of *E. inaequale* was straight to slightly curved, broadly ellipsoidal to almost cylindrical and widening in the middle, brown, 4-5-distoseptate, all the septa thickened and darkened, and central cell often longer and wider than the other cells,  $40-65 \times 15-22\mu\text{m}$  in size. The conidiophore was simple, septate, straight to flexuous, geniculate above, smooth, up to  $500\mu\text{m}$  long more, and  $4.5 \times 7.0\mu\text{m}$  wide. Optimum temperature was examined for conidial germination and mycelium growth on OA and PDA.

### O-02. Mycotoxins Produced by *Fusarium* Isolates from Barley Collected from Chungnam Province in Korea. Soh Young Oh<sup>1</sup>, Jin-Cheol Kim<sup>2</sup> and Seung Hun Yu<sup>1</sup>. <sup>1</sup>Department of Agricultural Biology, College of Agriculture, Chungnam National University Taejon 305-764, Korea. <sup>2</sup>Screening Division, Korea Research Institute of Chemical Technology P. O. Box 107, Yusong, Taejon, Korea.

Thirty-four isolates of *Fusarium* species were isolated from barley seeds collected from Chungnam province in Korea in 1998. The predominant *Fusarium* species isolated from barley were *F. graminearum* and *F. culmorum*. Each isolate was grown on 100g autoclaved wheat for 2 weeks at 25°C, followed by 2 weeks at 10°C for the production of crude cultures. Crude cultures were extracted with aqueous methanol and purified by Florisil column. The column elutes were concentrated and applied to TLC and GC-Mass to detect DON, NIV, 4-ANIV and ZEA. Of the thirty-four isolates of *Fusarium*, 10 isolates produced NIV, 4 isolates produced NIV and ZEA, 2 isolates produced NIV and 4-ANIV, 5 isolates produced NIV, ZEA and 4-ANIV, 2 isolates produced ZEA, and 1 isolate produced ZEA and DON. The mean concentration of DON, NIV and ZEA produced by positive isolates were 0.030, 79 and 198  $\mu\text{g/g}$ , respectively.