

Australia. We collected nine species considered as new species of the genus *Filippinodillo* from Philippines. Among them two new species are described with illustrations of diagnostic characters. The new species are distinguished from previously described species of the genus in the shape of cephalon, locking structures and appendages.

A714

**Six New Species of the Genus
Spherillo Dana, 1852 (Crustacea,
Isopoda, Armadillidae) from
Philippines**

Dae Soo Jeon* and Do Heon Kwon

Department of Biology, Inje University, Kimhae
621-749

The genus *Spherillo* had been a problematic genus due to the absence of type designation and the tradition of isopod taxonomists to neglect the rules of nomenclatures. The problem was recently solved by Lehtinen, Taiti and Ferrara (1998) to choose *S. vitiensis* Dana, 1853 as the type species. At present, the genus *Spherillo* comprises only three species including the type species with numerous junior synonyms. Among the specimens which we collected in Philippines, we found six new species of the genus *Spherillo*. They are described with illustrations of diagnostic characters. Each of them has unique color pattern and morphology.

A715

**Systematic Study of Roe Deer
(*Capreolus pygargus tianschanicus*)
Based on Sequence Analyses of
Mitochondrial DNA Control Region
and Cytochrome b Genes with
Specimens from Far East Asia**

**Hung Sun Koh¹, Beong Guk Yang¹, Goo
Hee Kwon¹, Jung Won Yoo¹, Kwang Sun
Kim¹ and Randi Effore²**

Dept. of Biology, Chungbuk University, Cheongju
361-763¹; Nazionale per la Fauna Selvatica, Ozzano
dell'Emilia, Italy²

We carried out the analysis of partial sequence of mtDNA control region and cytochrome b gene with roe deers from Cheju (Korea), Chenyang (China), and Vladivostok (Russia). The sequence analyses of mtDNA control region of roe deers from Eurasia were also conducted. In the sequence analysis of mtDNA control region, Korean roe deer (*Capreolus p. tianschanicus*), Kurgan roe deer (*C. p. pygargus*), and Amur roe deer (*C. p. pygargus*) appeared to be distinct with one another, but Korean roe deer was more closely related to roe deer from Kurgan region than roe deer from Amur region. In the sequence analysis of mtDNA cytochrome b genes with roe deers of *C. p. tianschanicus* from Cheju, Chenyang, and Vladivostok, Korean samples were different from Chinese and Russian samples. Therefore, it is confirmed that 1) Korean roe deer from Cheju island is a distinct subspecies of *C. c. ochracea*, as described by Barclay (1935), and 2) far east Asian roe deer from north east China, neasby Russia, and Amur region is classified into *C. p. bedfordi*, as noted by Sokolov & Gromov (1990).

A716

**Sequence Analysis of Mitochondria
DNA Control Region and
Cytochrome b Gene with Korean
Raccoon Dog (*Nyctereutes
procyonoides koreensis*) from
Goesan**

**Yong Chul Ahn*, Dong Sun Shin, Jeong
Gyu Park and Hung Sun Koh**

Dept. of Biology, Chungbuk University, Cheongju
361-763

We obtained partial sequences of

mitochondrial DNA control region and cytochrome b gene. In the sequence analysis of mtDNA control region, Korean racoon dog appeared to be distinct from Japanese racoon dog with average 90% sequence similarity and 0.110 pairwise distance of Kimura 2-parameter. In the sequence analysis of mtDNA cytochrome b gene, they showed average 98% similarity and 0.010 pairwise distance. It was revealed that Korean racoon dog is distinct in the mtDNA sequence. Therefore, it is concluded that Japanese racoon dog (*Nyctereutes procyonoides viverrinus*) is a subspecies which is different in chromosomal karyotype and morphometry from Chinese racoon dog (*Nyctereutes p. procyonoides*) and in mtDNA sequences from Korean racoon dog (*Nyctereutes p. koreensis*). However, in order to clarify the subspecific status of Korean racoon dog, samples of Chinese one is needed for further analyses.

A717

Taxonomic Status of Korean Hare based on Mitochondrial DNA Cytochrome B Gene Comparison

Tae Young Chun^{*}, Sun Ook Heo and Hung Sun Koh

Depart. Biology, Chungbuk University, Cheongju 371-763

After the analysis with partial sequence of mtDNA cytochrome b gene of Korean hare (*Lepus coreanus*) from Mt. Weolak, we compared this sequence with those of Chinese hare (*Lepus sinensis*) and Manchurian hare (*Lepus mandshurinus*) obtained from Genbank. It was revealed that Korean hare is more similar with Manchurian hare than Chinese hare in their mtDNA sequences, and it is confirmed that Korean hare is not a subspecies of Chinese hare but a distinct species of *L. coreanus*, as concluded by Jones and Johnson (1965). Moreover, it becomes necessary to carry out

further mtDNA sequence analysis with additional specimens of Manchurian and Korean hares in order to decide that Korean hare is a subspecies of Manchurian hare, as noted by Flux and Angermann (1990).

A718

Mitochondrial DNA Cytochrome b Sequence of Korean Red Squirrel (*Sciurus vulgaris coreae*)

Hye Sook Yoo^{*}, Bae Keun Lee and Hung Sun Koh

Dapt. of Biology, Chungbuk University, Cheongju 361-763

We compared the partial sequence of mtDNA cytochrome b of Korean red squirrel (*Sciurus vulgaris coreae*) resulted from this study with those of red squirrel from Korea (*S. v. coreae*), Hokkaido (*S. v. orientis*), Transbaikalia (*S. v. fusconigrans*), and Italy (*S. v. vulgaris*) obtained from Genbank. It was revealed that Korean red squirrel is more or less similar with Transbaikalia and Italy red squirrel, and that Hokkaido red squirrel is different from other three subspecies. Therefore, it is confirmed that *S. v. coreae* and *S. v. fusconigrans* are the synonym of *S. v. vulgaris*, as noted by Corbet (1978). Moreover, it is concluded that *S. v. orientis* is a distinct subspecies, as suggested by Corbet (1978).

A719

Isolation and Phylogeny of Endogenous Retroviral Elements Belonging to the HERV-K LTR in cDNA Library of Human Fetal Brain and Xq21.3 Region Linked to Psychosis

Joo-Young Choi, Seung-Heui Jeon, Joo-Mi Lee, Jun-Seop Kim, Won-Ho Lee and Heui-Soo Kim

Division of Biological Sciences, College of Natural