

absence of any metals.

333

Purification of the *E.coli* expressed p60 Protein from *Listeria welshimeri* by Amylose Resin Based Affinity Chromatography

Chil-U Lee*1, Hee-Young Lim1, Ki-Ho Han1, Yong-Soon Lee2, Yoon-Kyu Lim3, Byoung-Su Yoon1

Dept. of Biology, College of Natural Science, Kyonggi University*1 College of Veterinary Medicine, Seoul National University2 College of Veterinary Medicine, Cheju National University3

The *Listeria welshimeri* is an animal and human pathogen and its p60 protein is a major extracellular protein, which is encoded in iap (invasion associated protein) gene. These proteins are believed to be involved in the invasion of these bacteria into their host cells. To produce p60 in *E.coli*, the iap gene was recombinantly cloned and overexpressed. A purification protocol was developed for MBP (maltose binding protein)-p60 fusion protein by amylose-resin based affinity chromatography. The purified MBP-p60 was detected either as denatured or neutralized form using a specific p60 monoclonal antibody. The method might be an easy alternative to common purification protocols of p60 from *Listeria* spp. for antibody production.

334

Studies on Mechanism and Binding Proteins of Human HtrA2, Serine Protease

Kyoung Hoa Lee and Jung Woo Kim

Bio-Med RRC and Division of Life Sciences of Biochemistry, Pai Chai University, TaeJön 302-735

Human HtrA 2 (omi) is a recently

described novel member of the mammalian serine proteases family homologous to the *Escherichia coli* Htr A gene that are essential for bacterial survival at high temperatures. Although the physiological function of this new family is unclear, the current understanding is that as well as being involved in the degradation of aberrantly folded proteins during the conditions of cellular stress, they may possess a chaperon-like role under normal conditions. For understanding the mechanism of the HtrA 2 protein, we isolated HtrA 2 binding protein using the yeast two-hybrid assay. The pLex-HtrA 2, containing the gene for N-terminal region of the HtrA 2 inserted into PEG202 vector, was used as a bait plasmid to identify interacting protein gene from a human HeLa cDNA library. Final 16 X-gal positive colonies were selected from 1×10^7 colonies, isolated the plasmid containing the library cDNA and sequenced. The selected sequence represented the human karyopherin alpha 2, human embryonic ectoderm development protein (HEED), NADH dehydrogenase (ubiquinone) 1.

335

HIV-1 Nef Decreased the Transcription Activity of the CREB-2

Mi Seon Lee, Yong-Soo Bae¹ and Jung Woo Kim

Bio-Med RRC and Division of Life Sciences of Biochemistry, Pai Chai University, TaeJön 302-735; Dept. of Microbiology, Hannam University, TaeJön 300-791¹

Nef is a 27kDa myristoylated phosphoprotein expressed early in infection by HIV-1. To find the HIV-1 Nef interacting protein from T Jurkat cell, yeast two hybrid system was performed. The sequence determination and homology search of the isolated cDNA showed that one of the selected colonies encode the human cAMP

responsive element regulatory protein, CREB-2. Previously, we showed specific interaction between the Nef and CREB-2 by using the in vitro protein binding assay. We showed that the basic leucine zipper carboxy terminal domain of CREB-2 is required for the interaction with the HIV-1 Nef by using in liquid β -galactosidase assay. And, we demonstrated that CREB-2 cooperates with Nef to enhance viral transcription. Here, we report the construction of the CRE-SEAP reporter plasmid and the stable cell line in Jurkat cell. Using this plasmid investigated the activity of the transfected cells. CREB-2 has the effect on the transcription activity on the pGL2 promoter when we performed the transient transfection assay with Nef and CRE.

E301

Alteration of Glutathione-Ascorbate Cycle in Tomato Seedlings Exposed to Cadmium

Kyung-Hee Hwang*, Chong-Min Lee and Un-Haing Cho

Dept. of Biology, Changwon National University, Changwon 641-773

Developmental changes of glutathione-ascorbate cycle in seedlings of tomato (*Lycopersicon esculentum*) exposed to toxic cadmium concentrations (up to 100 mM for 9 days) were investigated. The contents of total protein and thiols in both leaves and roots and total glutathione in leaves increased but the content of total ascorbic acid in leaves was maintained unchanged with cadmium exposure. Further, the ratio of oxidized glutathione (GSSG)/reduced glutathione (GSH) in both leaves and roots increased but the ratio of dehydroascorbate/ascorbate and the activity of glutathione reductase in leaves decreased with cadmium exposure. Our results imply that tomato seedlings exposed to toxic levels

of cadmium experience oxidative stress which might be caused by the impaired glutathione-ascorbate cycle resulting from the altered activities of enzymes and the lowered production of antioxidant compounds involved in the oxidation-reduction cycle.

E301

Development of the Screening System for Antagonist of PPAR- γ by Using Yeast Two Hybrid System

Soo-Jin Kwon and Jung Woo Kim

Bio-Med RRC and Division of Life Sciences, Pai Chai University, Taejon, 302-735; Institute of IMAGENE, Suwon 440-746

Obesity, an excessive accumulation of adipose tissue, is characterized by an increase in the number or size of adipocyte, or both. Adipocyte is highly specialized cell that play a critical role in energy homeostasis. Adipocyte differentiation is characterized by a coordinate increase in adipocyte-specific gene expression. Recently PPAR- γ was shown to be expressed in an adipose-specific manner and its expression induced early during the course of differentiation of several preadipocyte cell line. PPAR- γ and the retinoid X receptor α (RXR α) form a heteromeric complex that function as a central regulator of adipocyte differentiation. We developed the screening method to search the candidates for the inhibitors of the differentiation of preadipocyte to adipocyte, by using the yeast two hybrid system. The bait and prey were RXR and PPAR- γ , respectively. Using this system, we found several candidates from the culture broth of *Streptomyces* sp. and some plant extract.

E301

Oxygen Defense Role of Xanthine Dehydrogenase