

P-47

## CELECOXIB ATTENUATES ET-18-O-CH<sub>3</sub>-INDUCED APOPTOSIS IN H-ras TRANSFORMED HUMAN BREAST EPITHELIAL CELLS

Hye-Kyung Na and Young-Joon Surh

College of Pharmacy, Seoul National University, Seoul 151-742, Korea

Cyclooxygenase-2 (COX-2) is an inducible enzyme expressed in response to a variety of proinflammatory agents and cytokines. COX-2 expression has been shown to be elevated in several different types of human cancer. The presence of oncogenic ras has been associated with constitutive induction of COX-2 in certain H-ras transformed cells, and COX-2 overexpression confers resistance to apoptosis. Contrary to the above notion, we have found that apoptotic death occurs while the COX-2 expression is induced in the H-ras transformed human breast epithelial cell line (MCF-10A-ras) treated with ET-18-O-CH<sub>3</sub> (1-O-octadecyl-2-O-methyl-glycero-3-phosphocholine), a specific inhibitor of phosphoinositide phospholipase C. Thus, while MCF-10A-ras cells were found to undergo apoptosis by treatment with ET-18-O-CH<sub>3</sub> as revealed by proteolytic cleavage of poly(ADP-ribose) polymerase (PARP) and positive terminal deoxynucleotidyl transferase-mediated dUTP nick-labeling (TUNEL), the cells exhibited an increased expression of COX-2 as well as the elevated production of prostaglandins (PGE<sub>2</sub>). To determine whether the expression of COX-2 by ET-18-O-CH<sub>3</sub> is associated with the induction of apoptosis, the effects of the selective COX-2 inhibitor celecoxib (SC-58635) on ET-18-O-CH<sub>3</sub>-induced apoptotic death were examined. Treatment of MCF-10A-ras cells with celecoxib (50  $\mu$ M) attenuated ET-18-O-CH<sub>3</sub>-induced apoptosis as well as COX-2 expression and production of PGE<sub>2</sub> in a concentration dependent manner. To delineate the upstream signaling pathways responsible for the apoptosis-inducing activity of ET-18-O-CH<sub>3</sub>, we examined the effect of this drug on activation of the anti-apoptotic protein Akt/protein kinase B and also mitogen-activated protein (MAP) kinases such as c-Jun NH<sub>2</sub>-terminal kinases (JNK), extracellular signal-regulated kinases (ERK) and p38 MAP kinase that play a role in regulation of *cox-2* gene expression. ET-18-O-CH<sub>3</sub> inhibited the activation of ERK1/2 and p38 MAPK but not JNK, and also blocked the phosphorylation

of Akt/protein kinase B. Inhibitory effects of ET-18-O-CH<sub>3</sub> on activation of p38 MAPK, ERK1/2, and Akt were attenuated by co-treatment with celecoxib. Taken together, the above findings suggest that COX-2 up-regulation does not necessarily confer the resistance to apoptosis in *ras*-transformed cells, but may rather sensitize the transformed cells to apoptotic death.