

니코틴 중독의 단서노출치료를 위한 가상환경의 제작 및 욕구 유발 실험

김광욱, 조원근, 구정훈, 김훈, 김병년, 이장환, 김인영, 이종민, 김선일
한양대학교 의공학 교실

Cue exposure system using Virtual Reality for nicotine craving

Kwanguk Kim, Wongeun Cho, Jeonghun Ku, Hun Kim, Byoungnyun Kim,
Jang Han Lee, In Y. Kim, Jong Min Lee, Sun I. Kim
Dept. of Biomedical Engineering, Hanyang University, Seoul, Korea

Abstract

Research has shown that many smokers experience an increase in the desire to smoke when exposed to smoking related cues. Cue exposure treatment (CET) refers to the manualized, repeated exposure to smoking related cues, aimed at the reducing cue reactivity by extinction. In this study, we constructed a virtual reality system for evoking a desire of nicotine, which was based on the results of a Questionnaire of Nicotine craving. And we investigated the effectiveness of the virtual reality system as compared to classical device (pictures). As a result, we reached the conclusion that virtual reality elicits more craving symptoms than the classical devices.

Keywords: Cue exposure, Virtual Reality (VR), Nicotine craving, Cue exposure therapy (CET)

1. Introduction

“Craving” is a term derived from popular psychology, and is used to describe a mental state, namely, the intense desire for a certain object or experience (e.g., alcohol, nicotine, or drug) [1]. The World Health Organization in 1955 made a distinction between nonsymbolic craving (i.e. physiological withdrawal) and symbolic craving (i.e. loss of control, which is associated with relapse). It has been suggested that the term ‘craving’ should be replaced by the terms ‘physical dependence’ and ‘pathological desire’, which refer respectively, to the two conditions described above [2,3].

Researches have shown that many smokers experience an increase in the desire to smoke when exposed to smoking-related cues [4,5]. A strong desire to smoke, in turn, seems to play an important role in the maintenance of cigarette smoking [5,6]. According to learning-based theories, cue-induced craving might partly reflect a conditioned response (CR) established by a learned association between the cue (CS) and nicotine intake (US) [5,7]. Cue exposure techniques, which try to extinguish this learned association, have been increasingly forwarded as a potential treatment of addictive behaviors, including cigarette smoking [5,8].

Conditioned reactivity to cues is an important factor in the addiction to alcohol, nicotine, opiate, and cocaine. Cue exposure treatment (CET) refers to the manualized, repeated exposure to drug-related cues, aimed at reduction cue reactivity by extinction. In current CET, various stimuli are presented, such as, slides, video tapes, pictures, or paraphernalia in nonrealistic, experimental settings [9].

Virtual Reality can make a person look, feel, hear and interact in a computer

-generated situation. One pilot study on addiction to cocaine showed that immersive virtual reality (IVR) is as good or even better at eliciting subjective and physiological craving symptoms as classical devices [9].

In this study, we designed a virtual reality system to create desire for nicotine, which was based the responses from a nicotine-craving questionnaire. The virtual environment is composed of craving environments, craving objects and smoking avatars. We compared this virtual reality system to Picture, and investigated the potential value of the use of a virtual reality system for cue exposure treatment (CET).

2. Preliminary study

2-1 Nicotine Craving Questionnaire

Before we design the virtual environment for eliciting a smoking desire, we collected data about which cue would elicits one’s craving by using a questionnaire (See Table 1).

Table 1 Questions

| | |
|-----------|--|
| Section 1 | Tendency of smoking. Amount of smoking. Whether or not the subject had attempted to quit smoking. |
| Section 2 | Which object does induce your craving? Which place (or situation) does induce your craving? Which is more attractive between the object and place? |
| Section 3 | Nicotine Addiction Test (Only used on experiment subjects). Nicotine dependence: low, high, very high |

2-2 Questionnaire Result

Data were collected from 64 men, aged 19 to 38 years. The result showed that a place

evokes a higher level of craving than an object, bars and tobacco were the places and objects most often cited by participants (See Figure 1,2 and 3).




Fig. 1 Which one is more likely to elicit craving—the object or the place?




Fig. 2 What place induces craving?




Fig. 3 Which objects induces craving most?

This result shows that the VR could be used to elicit more one's desire related smoking because the VR can provide a immersive situation. So, using the result from this survey, we designed a virtual bar environment. Objects in the virtual bar were also selected from the questionnaire results. They are an alcoholic drink, a pack of cigarettes, a lighter, an ashtray, and a glass of beer.

3. Main Experiment

3-1 Subject

22 male smokers participated in this experiment. All subjects were free from brain injury, glycosuria, high blood pressure or medicinal poisoning, and were not handicapped in mentally or neurologically. The average of amount of smoking per day was 12.64, and the average nicotine addiction test score was 4.14.

3-2 Instrument

a. System

The Virtual Reality System consisted of a Pentium IV PC, OpenGL Accelerator VGA Card, HMD (i-visor DH-4400VPD), and a 3DOF Position Sensor (Intertrax2). The PC with a 3D Accelerator VGA Card generated real-time virtual images for the subject to navigate. The position sensor (tracker) transferred a subject's head orientation to the computer.



Fig. 4 System Composition Diagram

The Classical Device was displayed using the same virtual reality system, but the tracker was not used in order to eliminate the difference of the display modality.

Therefore, the VR system could provide a dynamic situation while the classical device provided a still picture.

b. Virtual Environment

The Virtual Environment was designed based on the nicotine-craving questionnaire. The background environment present here is a bar, the 5 kinds of objects were an

alcoholic drink, a pack of cigarettes, a lighter, an ashtray, and a glass of beer and avatars smoking cigarette (See Fig. 5 and 6).




Fig. 5 Virtual bar & Craving Object



Fig. 6 Avatar animation in the virtual bar

c. Classical Device (Picture)

Classical Device (CD) was also based on a questionnaire of nicotine-craving. Therefore the CD's background was bar, with the same 5 object types, an alcoholic drink, a pack of cigarettes, a lighter, an ashtray, and a glass of beer. CD contained the same content as the VE.

3-3 Procedure

Before the experiment subjects were asked for their, ages, educational backgrounds and medical history, to complete a questionnaire on nicotine-craving (a mount of smoking per day and nicotine craving of individual person), and to complete a Nicotine Addiction Questionnaire (from 0 to 11) and a Questionnaire on Current Nicotine craving. For the main experiment, the subjects were split into two groups randomly, a Virtual Reality (VR) group and a Classical Device (Picture) group and subjects were experience the VR and CD respectively for 5 min

3-4 Measurement

A Visual Analogue Scale (VAS) was used to subjectively measure the craving to smoke. This VAS was defined to have a range of 0-100 mm, from 'none at all' to 'extreme,' and individual rating was decided by subjects in response to the question: How strong is your urge to smoke now?

4. Result

Table 2 shows Mean (M), Standard Deviation (SD) and t-test result, nicotine addiction test score and the difference of nicotine craving between before and after experiment in both the CD group and the VR group. And it shows also the group difference, which means that the subjects of the two groups are no different in terms of group. But the difference of nicotine craving (After-Before) was significantly different.

Table 2 Compare of CD Group and VR Group

| | Group | | | | t-test |
|--|-------|------|------|------|-----------------|
| | CD | | VR | | |
| | M | SD | M | SD | |
| A mount of smoking per day | 13.17 | 9.83 | 12.0 | 6.85 | 0.321 (n.s) |
| Nicotine Addiction Test | 4.33 | 2.06 | 3.90 | 1.97 | 0.501 (n.s) |
| Usual Nicotine craving | 6.08 | 2.23 | 6.40 | 1.85 | 0.371 (n.s) |
| Current Nicotine craving before experiment | 5.67 | 2.81 | 5.40 | 1.96 | 0.253 (n.s) |
| Current Nicotine craving after experiment | 5.08 | 2.97 | 6.40 | 2.22 | 1.157 (n.s) |
| The difference of Nicotine craving between before and after experiment | 0.58 | 1.82 | 1.00 | 1.25 | -2.524 (p<0.05) |

CD=Classical Device (Picture), VR=Virtual Reality, M=Mean, SD=Standard Deviation

Figure 7 shows craving change before and after the experiment for the Classical Device (Picture) group and the Virtual Reality group.




Fig. 7 Nicotine craving Before & After Experiment

Figure 7 shows that the Classical Device group's nicotine craving changed from 5.67 to 5.08, and the Virtual Reality group's nicotine craving is increased from 5.40 to 6.40. However, the difference between the before and after experiment results of the Classical Device group was not significant ($t=1.246$, n.s.), and this difference in the Virtual Reality group was significant ($t=-2.535$, $p<0.05$).

5. Discussion

In the present study, we tried to determine whether virtual reality can be used in cases of nicotine-craving as a cue exposure treatment tool. Therefore, we compared a virtual reality system with a classical device (Picture), and measured difference of nicotine-craving using VAS.

We found that virtual reality is better at eliciting than classical device (Picture) for CET. This result is accord with the results of KUNTZE's pilot test for opioid dependent subjects [9]. It is because a VR can present a spatial stimulus, which resembles that of the real world and the VR is more immersive than Classical CET device.

As a result, the nicotine craving of the VR group increased significantly from 5.40 to 6.40. However, the CD group's nicotine-craving only showed a tendency to decrease from 5.67 to 5.08. This result shows that the picture showing is not as suitable a

device for eliciting nicotine-craving in normal subject. Additional experimentation in 8 subjects reconfirmed that the observed CD group's decrease was non-significant. Nicotine-craving, including of additional experiment, is changed from 5.50(before exp.) to 5.45(after exp.).

The future study is needed to investigate whether VR is superior to other classical methods at evoking nicotine craving other sources of objective evidence by acquiring and analyzing objective evidence such as, bio signals (Heart Rate, Galvanic Skin Response), fMRI (functional Magnetic Resonance Imaging) or PET (Positron emission tomography) and the possibility of a VR system for the other types of abuse and to apply the VR system to treat such abuse.

6. Conclusion

Our study demonstrates that virtual reality is probably better at eliciting craving symptoms than classical picture watching and we propose that Virtual Reality be used in Cue exposure Therapy.

Acknowledgement

This study was funded by the National Research Laboratory (NRL) Program at Korea Institute of Science & Technology Evaluation and Planning (2000-N-NL-01-C-159).

7. Reference

- [1] Daniel W. Hommer, "Functional imaging of craving." Alcohol Research & Health, Vol23 No5, pp187-196, 1999
- [2] Christos Kouimtsidis, "Role of craving in substance misuse". Current Opinion in Psychiatry, 13, pp299-303, 2000
- [3] Sinha R, O'Malley SS, "Craving for alcohol: findings from the clinic and the laboratory", Alcohol Alcoholism, 34, pp223-230, 1999

- [4] Payne, T. J., Schare, M. L., Levis, D. J., & Colletti, G., "*Exposure to smoking-relevant cues: Effects on desire to smoke and topographical components of smoking behavior.*" *Addictive Behaviors*, 16, pp467-479, 1991
- [5] M. Dols, B. Willems, M. Van Den Hout, & R. Bittoun, "*Smokers can learn to influence their urge to smoke.*" *Addictive Behaviors*, Vol25 No1, pp103-108, 2000
- [6] Killen, J. D., & Fortmann, S. P. "*Craving is associated with smoking relapse: findings from three prospective studies*", *Experimental and Clinical Psychopharmacology*, Vol5 No3, pp137-142, 1997
- [7] Niaura, R., Rohsenow, D. J., Binnkoff, J. A., Monti, P. M., Pedrazza, M., & Abrams, D. B. "*Relevance of cue reactivity to understanding alcohol and smoking relapse.*" *Journal of Abnormal Psychology*, Vol 97 No 2, pp133-152, 1998
- [8] Marlatt, G. A. "*Cue exposure and relapse prevention in the treatment of addictive behaviors*", *Addictive Behaviors*, 15, pp395-399, 1990
- [9] Marcus F. Kuntze, Robert Stoermer, Ralph Mager, Andreas Roessler, Franz Mueller-Spahn, & Alex H. Bullinger. "*Immersive Virtual Environments in Cue Exposure*", *CyberPsychology & Behavior*, 4(4):497-501, 2001
- [10] George V. Popescu. *Multimodal Interaction Model, HANDBOOK OF VIRTUAL ENVIRONMENT Design, Implement, and Application*, Kay M. Stanney Eds., LEA Inc., New Jersey, 2002