

3-3-12. The *Antheraea pernyi* Arylphorin That is Immunologically Cross-Reacted with the Storage Proteins of the Wild Silkmths, *Antheraea yamamai*, *Samia cynthia pryeri* and *Actias gnoma*

Bo Youn Kim¹, Gae Jong Choi¹, Nam Sook Park², Byung Rae Jin¹,
Sang Bong Park², Hong Ja Kim³, Sook Jae Seo³ and Sang Mong Lee²

¹Department of Sericultural and Entomological Biology, Faculty of Agriculture, Miryang National University, Miryang 627-130, Korea; ²College of Natural Resources and Life Science, Dong-A University, Busan 640-714, Korea; ³Division of Life Science, Gyeongsang National University, Chinju 660-701, Korea

The synthesis or expression of proteins related to *Antheraea pernyi* arylphorin in haemolymph, fat body, integument, mid gut and silk gland of the wild silkmths, *A. yamamai*, *A. pernyi*, *Samia cynthia pryeri* and *Actias gnoma* in the 5th larval instar were investigated by immunoblot analysis and immunocytochemistry using mouse polyclonal antibody for *A. pernyi* arylphorin. In *A. yamamai*, *A. pernyi* and *A. gnoma*, the major immunoreactive proteins with a molecular weight of 80 kDa were observed in the haemolymph, integument, fat body and mid gut except silk gland. In *S. cynthia pryeri*, the immunoreactive proteins with a molecular weight of 80 kDa were also observed in the haemolymph, integument and fat body except mid gut and silk gland. These results revealed that the *A. pernyi* arylphorin has almost same identity with those of the wild silkmths, *A. yamamai*, *S. cynthia pryeri* and *A. gnoma* with respect to immunological reactivity. In addition, it can be considered that the *A. pernyi* arylphorin has been evolved from a common ancestor gene of the wild silkmths described as the above.