

# **Antioxidant Activity of *Cercis chinensis* and Its Protective Effect on Skin Aging**

**MinKyun Na<sup>1,2</sup>, KiHwan Bae<sup>2</sup>, NamDoo Hong<sup>1</sup>, JaeKuk Yoo<sup>1</sup>, Nobuhiko Miwa<sup>3</sup>**

<sup>1</sup>*Jakwang Research Institute, Hansaeng Cosmetics Co., Ltd., Nonsan 320-854, Korea,* <sup>2</sup>*College of Pharmacy, Chungnam National University, Daejeon 305-764, Korea,*

<sup>3</sup>*Department of Biotechnology, Hiroshima Prefectural University School of BioSciences, Shobara, Hiroshima 728-0023, Japan*

\* Correspondence to: MinKyun Na, Jakwang Research Institute, Hansaeng Cosmetics Co., Ltd.,

Nonsan 320-854, Korea

E-mail: [nmk1010@cnu.ac.kr](mailto:nmk1010@cnu.ac.kr)

Tel: +82-41-740-2405

Fax: +82-41-740-2429

**Keywords:** *Cercis chinensis*, antioxidant activity, oxidative damage, UVB irradiation, cellular aging

## **Summary**

Reactive oxygen species are capable of damaging biomolecules such as lipids, proteins, and DNA, which can not only lead to various diseases, but also oxidative damage resulting aging. In our previous study, *Cercis chinensis* (Leguminosae) showed a potent antioxidant activity. Nineteen compounds were isolated through antioxidant activity-guided fractionation. The *C. chinensis* extract and some of the constituents exhibited a potent antioxidant activity on the free radicals and lipid peroxidation and a notable protective effect on the *t*-BuOOH induced oxidative damage. *In vivo* test of skin damage induced by UVB irradiation, the extract of *C. chinensis* and a constituent, piceatannol, exhibited a significant protective effect. The life-span of the HEK-N/F cells were extended by 1.21—2.12 fold as a result of the continuous administration of 3  $\mu$ g/ml of the *C. chinensis* extract and the active constituents compared to that of the control. These observations were attributed to the inhibitory effect of the *C. chinensis* extract and its constituents on the age-dependent shortening of the telomere. Thus, *C. chinensis* was demonstrated to protect the skin cells against oxidative stress and inhibit thereby the cellular aging, followed by expectation as anti-aging cosmetic ingredient.

## Introduction

It is well known that reactive oxygen species (ROS) generated during the metabolic pathway damage biomolecules such as lipids, proteins, sugars and DNA, which can not only lead to various diseases [1], but also oxidative damage resulting aging [2],[3]. Indeed, oxidized biomolecules such as the 8-oxo-2'-deoxyguanosine residues in DNA, carbonyls and dityrosines in proteins and hydroperoxides in lipids have been reported to accumulate in the tissues of aged animals [4—6].

This oxidative stress theory of aging is supported by many studies, and it is now accepted one of the most important theories of aging. Accordingly, antioxidants, which can prevent oxidative damage from ROS, are expected to inhibit the aging process. Recently, another theory, the telomere hypothesis of aging, has also been accepted together with the oxidative stress theory.

Telomeres located at the end of linear chromosome shorten with each cell division, and DNA damage if the telomeric DNA reaches a critical length, which can ultimately, lead to cell cycle arrest in senescent cells [7],[8]. Shortening of the telomeric DNA has been observed in human cells during the aging processes [9]. A life span extension has been observed after introducing telomerase, which can maintain the telomere length, into normal human cells [10].

In our previous study, the *Cersis chinensis* extract exhibited potent antioxidant activity on DPPH radical and lipid peroxidation. *C. chinensis* which belongs to the Leguminosae is widely cultivated as an ornamental plant. The stem bark, root bark and stem of *C. chinensis* have been used for a

promotion of blood circulation, dysmenorrhea, edema, a bruise, and an injury [11].

Based on the two aging theories, this study investigated the antioxidant activity of the *C. chinensis* extract and its constituents, and the protective effect on the oxidative stress induced by tertiary butyl hydroperoxide (*t*-BuOOH) and ultraviolet (UV)-B. Furthermore, the inhibitory effect of the medicinal plant on the cellular aging of human epidermal keratinocytes was examined by measuring the telomere length.

## **Materials and Methods**

**Plant Material** - The leaves and stem of *C. chinensis* were collected at Daejeon, Korea in September 2001 and was identified by Prof. KiHwan Bae, College of Pharmacy, Chungnam National University. A voucher specimen (HK 1122) was deposited in the Jakwang Research Institute of the Hansaeng Cosmetics Co., Ltd.

**Extraction and Isolation** - The dried leaves and stem from *C. chinensis* (1 kg) were extracted with 60% EtOH at room temperature for 2 weeks. The EtOH extract (75 g) was suspended in water and then partitioned with hexane, EtOAc and BuOH, sequentially. The EtOAc and BuOH fraction (Fr.) exhibited a potent activity with an IC<sub>50</sub> value of 24.0 and 27.0 µg/ml, respectively, in the DPPH assay. YMC gel and silica gel column chromatography, and preparative HPLC of the active EtOAc-

and BuOH-soluble fraction led to the isolation of nineteen compounds (1—19). They were identified as isoliquiritigenin (1), liquiritigenin (2), 2',4'-dihydroxy-4-methoxychalcone (3), resveratrol (4), piceatannol (5), gallic acid (6), methyl gallate (7), ethyl gallate (8), myricetin (9), afzelin (10), quercitrin (11), myricitrin (12), myricetin-3-O-(2''-O-galloyl)- $\alpha$ -L-rhamnopyranoside (13), syringetin-3-O-rutinoside (14), (+)-catechin (15), (-)-epicatechin-3-O-gallate (16), (-)-epigallocatechin-3-O-gallate (17), (-)-lyoniresinol-3a-O- $\beta$ -D-xylopyranoside (18) and (+)-lyoniresinol-3a-O- $\beta$ -D-glucopyranoside (19) by comparing their spectral data with those previously reported [12—18].

**Antioxidant activity on the free radicals and lipid peroxidation** - The DPPH radical, superoxide radical scavenging activity was measured using a method described previously [19]. The lipid peroxidation inhibitory activity in a rat brain homogenate was evaluated using the thiobarbituric acid (TBA) method described previously [19].

#### **Oxidative stress induced cell damage**

**Cell Culture** - The human epidermal keratinocytes-Neonatal/Foreskin (HEK-N/F) cells were purchased from Modern Tissue Technologies, Inc. (MC1312, Korea). The HEK-N/F cells were cultured in a type IV collagen coated plate with KGM<sup>®</sup> Bulletkit medium (CC-3111, Clonetics) in humidified atmosphere of 5% CO<sub>2</sub>/95% air at 37°C, and cultured to 90% confluence.

**t-BuOOH induced oxidative stress** - The HEK-N/F cells ( $1 \times 10^4$  cells/100  $\mu$ l) were seeded on a 96

well microplate and precultured for 24 h. The cells were then treated with 1  $\mu$ l of the sample and 10  $\mu$ l of t-BuOOH (1.5 mM) dissolved in Hank's balanced salt solution (HBSS, Gibco. BRL) for 3 h in order to induce cellular peroxidation. The cell viability was measured using the MTT method. The inhibitory activity of lipid peroxidation was also determined using the thiobarbituric acid (TBA) method, as previously described [20].

### ***In vivo study***

Female SKH-1 hairless mice (5 weeks old) were housed under standard conditions (temperature  $24 \pm 2^\circ\text{C}$ ; relative humidity  $50 \pm 10\%$ ; 12 hr/day light/dark cycle) and given a commercial diet and water *ad libitum*. The mice were divided into four groups containing five mice each. Samples were administered topically at dose of 50, 30 and 10 mg/kg on their dorsal skin, and the dorsal skin of the mice was irradiated with UVB within 30 min of administration as described previously [21]. After 24 hr, the dorsal skin was collected from each mouse and frozen at  $-70^\circ\text{C}$  until used.

**Assay of lipid peroxidation** - The dorsal skin was homogenized in 10 volumes of a 50 mM phosphate buffer (pH 7.8) under  $4^\circ\text{C}$ . The lipid peroxidation level was measured using the TBA method. The TBARS concentration was presented as nmol malondialdehyde/mg protein. The proteins were determined using the dye binding method using a Bio-Rad protein assay kit (Bio-Rad Lab., USA).

### **Inhibitory activity of telomere length shortening**

**Cell culture** - The HEK-N/F cells were successively subcultured at a dilution rate of 1:8. The cells were grown to the population doubling level (PDL) of approximately 3 in the absence of the sample, and fed with or without the sample at a dose of 3  $\mu\text{g/ml}$  successively upon each culture passage.

The PDL was regarded to be zero for the culture starting immediately after the primary culture of the HEK-N/F cells, and was calculated to increase according to the following equation:  $\log_2$  [(the number of collected cells)/(the number of seeded cells)]

**Determination of telomere length by Southern blots** - The genomic DNA was extracted using a nucleic acid extraction kit IsoQuick (ORCA Research Inc.) from  $10^6$  cells at each passage. The extracted DNA was cleaved with the restriction enzyme, *Hinf* I (TaKaRa), to produce terminal restriction fragments (TRFs). The telomere length was determined by Southern blots as previously described [22].

### **Statistical analysis**

The Data is expressed as a mean  $\pm$  SD, and the differences between the control and test groups were analyzed using a two-tailed Student's t-test. Differences  $P < 0.05$  were considered to be statistically significant.

## Results and Discussion

### Antioxidant activity on the free radicals and lipid peroxidation

The antioxidant activity of the *C. chinensis* extract and its constituents (**1**—**19**) on the free radicals and lipid peroxidation is shown in Table I. Piceatannol (**5**) and myricetin (**9**) exhibited the potent radical scavenging activity and the lipid peroxidation inhibitory activity.

### Cytoprotective effect on the *t*-BuOOH induced oxidative stress

*t*-BuOOH is an hydroperoxidant that can be metabolized to free radical intermediates, which can subsequently initiate lipid peroxidation resulting in cell damage. The cytoprotective effect of the *C. chinensis* extract and its constituents induced by *t*-BuOOH is shown in Table II. The cell viability of the HEK-N/F cells was significantly decreased by  $11.2 \pm 1.2\%$  when they were treated with 1.5 mM *t*-BuOOH for 3 h, whereas the cell viability was increased by  $53.5 \pm 3.8\%$  when they were treated with the *C. chinensis* extract at a concentration of 50  $\mu\text{g/ml}$ . Some of the constituents exhibited potent cytoprotective activity (Table II). The cytoprotective activity was increased according to the number of hydroxyl groups on the aromatic ring, which suggests that their cytoprotective activity results in their electron donating activity, and in accordance with the radical scavenging activity [23]. Piceatannol (**5**) exhibited the most potent cytoprotective activity ( $84.7 \pm 6.9\%$ ) on the tested compounds. Glycosylation of a hydroxyl group is completely inactive on the cytoprotective activity.



The aglycone, myricetin (**9**), exhibited higher activity than the glycoside, myricitrin (**12**). These results are in accordance with those reported in other assay systems [23],[24]. Although, Myricetin-3-O-(2''-O-galloyl)- $\alpha$ -L-rhamnopyranoside (**13**), (-)-Epicatechin-3-O-gallate (**16**) and (-)-Epigallocatechin-3-O-gallate (**17**) exhibited strong activity on radical scavenging and lipid peroxidation due to the residual galloyl group, they did not show a high activity in this assay system.

The cytoprotective effect of the *C. chinensis* extract and its constituents (**1—19**) against the *t*-BuOOH induced oxidative stress was also confirmed by determining the level of lipid peroxidation (Table II). The TBA reactive substance (TBARS) of the cells treated with *t*-BuOOH (1.5 mM) for 3 h was increased 23.6-fold compared to the untreated control. As shown in Table II, the TBARS of the cells treated with the *C. chinensis* extract and some of the constituents, in particular, **5** were also significantly decreased, which means that the *C. chinensis* extract and its constituents could protect the cells from oxidative stress.

#### **Protective effect on UVB-induced skin lipid peroxidation (*In vivo*)**

The *C. chinensis* extract and piceatannol (**5**), which showed the most potent cytoprotective activity on oxidative stress, were further investigated to determine the protective effect against the UVB-induced skin lipid peroxidation *in vivo*. UVB radiation (in the range of 290-320 nm) is believed to be the major cause of skin cancer [25]. UVB is also known to induce excessive ROS generation, and lipid peroxidation is a marker of oxidative stress [25],[26]. As shown in Table III, the UVB exposed

mice skin had a significantly higher amount of TBARS ( $0.68 \pm 0.10$  nmol/mg protein) in 24 h after UVB irradiation. In contrast, UVB-induced skin lipid peroxidation was inhibited when the mice were topically administered with the *C. chinensis* extract at a dose of 50 and 30 mg/kg (Table III). The TBARS concentration was significantly reduced by  $0.07 \pm 0.03$  nmol/mg protein ( $P < 0.001$ ) when it was topically administered with **5** at a dose of 30 mg/kg. The amount of TBARS ( $0.16 \pm 0.03$  nmol/mg protein) was also significantly decreased even at dose as low as 10 mg/kg. As a result, it is demonstrated that the *C. chinensis* extract and **5** affords protection against the oxidative damage induced by UVB irradiation due to its potent antioxidant properties that efficiently reduce oxidative stress.

#### **Elongating effect on cellular life-span**

Most mammalian cells cultivated *in vitro* undergo a limited number of cell divisions and then arrest in a stage known as replicative senescence. In a report [22], it was found that cultivating the cells under enhanced oxidative stress accelerates the replicative senescence, and antioxidants could elongate the life span. To investigate the effect of the *C. chinensis* extract and its active constituents, **5** and **9**, on the cellular doubling potential, the HEK-N/F cells were serially subcultured until the spontaneous stoppage of cell division in the presence or absence of the samples. The untreated cells rapidly grew to a PDL of 14.1, whereas 3  $\mu$ g/ml of the *C. chinensis* extract, **5** and **9** added cells enhanced the maximum PDL to 17.0, 29.9 and 23.2, respectively. The cellular life was

extended 1.21-, 2.12- and 1.65-fold, respectively, in comparison to the control group (Fig. 1).

### **Inhibitory effect of age-dependent telomere shortening**

Based on the telomere theory of aging, the elongating effect of the cellular life span was evaluated by measuring the telomere length in each culture passage. The DNA extracted from the HEK-N/F cells of each culture passage was restricted by *Hinf* I to produce the TRFs, which were analyzed by Southern blot analysis. As shown in Fig. 2, the telomeres in the control cells gradually shortened from 12.42 kbp (PDL 2.8) to 7.90 kbp (PDL 14.1). In comparison to this, when the *C. chinensis* extract, **5** and **9** were continuously administered to the cells, the telomere length shortening was inhibited. It was observed that in the case of the cells administered to **5**, the telomere length had shortened by an average of 169 bp/PDL, which was slower than the 368 bp/PDL observed in the control group (Fig. 2). Cell division can be continuously promoted as far as the telomeric DNA is kept at more than a critical value (in this report, it is assumed to be approximately 7.9 kbp). From this viewpoint, it was found that the *C. chinensis* extract, **5** and **9** could be delay the shortening rate of the telomeric DNA and thereby, the rate to reach the critical value to be delayed and therefore it could extend a period for the termination of cell division.

### **Anti-aging effect of *T. chebula* based on the two aging theories**

The two aging theories, oxidative stress theory and telomere theory, have been established well by

many related evidences. Recently, it was recognized that oxidative damage is repaired less well in the telomeric DNA than elsewhere in the chromosome, and oxidative stress accelerates telomere loss [22],[27—29]. Measuring both antioxidant activity on oxidative stress and telomere shortening rate, we found that the *C. chinensis* extract and its constituents could slowdown the telomere shortening rate by inhibition of oxidative stress. Therefore, it is suggested that the *C. chinensis* and its constituents protect the skin cells against oxidative stress and prevent thereby the cellular aging, followed by expectation as anti-aging cosmetic ingredient.

## References

- [1] Vishwanath, M.S., Role of antioxidants in health maintenance, *Nutr. Clin. Pract.*, **10** (1995) 19-25.
- [2] Barja, G., Endogenous oxidative stress: relationship to aging, longevity and caloric restriction, *Ageing Res. Rev.*, **1** (2002) 397-411.
- [3] Sohal, R.S., Mockett, R.J., Orr, W.C., Mechanism of aging: an appraisal of the oxidative stress hypothesis, *Free Radic. Biol. Med.*, **33** (2002) 575-586.
- [4] Lezza, A.M.S., Mecocci, P., Cormio, A., Beal, M.F., Cherubini, A., Cantatore, P., Senin, U., Gadaleta, M.N., Mitochondrial DNA 4977 bp deletion and OH<sup>8</sup>dG levels correlated in the brain of aged subjects but not Alzheimer's disease patients, *FASEB J.*, **13** (1999) 1083-1088.
- [5] Berlett, B.S., Stadtman, E.R., Protein oxidation in aging, disease, and oxidative stress, *J. Biol. Chem.*, **272** (1997) 20313-20316.
- [6] Tahara, S., Matsuo, M., Kaneko, T., Age-related changes in oxidative damage to lipids and DNA in rat skin, *Mech. Ageing Dev.*, **122** (2001) 415-426.
- [7] Harley, C.B., Vaziri, H., Counter, C.M., Allsopp, R.C., The telomere hypothesis of cellular aging, *Exp. Gerontol.*, **27** (1992) 375-382.
- [8] Vaziri, H., Benchimol, S., From telomere loss to p53 induction and activation of a DNA-damage pathway at senescence: The telomere loss/DNA damage model of cell aging, *Exp. Gerontol.*, **31**

(1996) 295-301.

[9] Tsuji, A., Ishiko, A., Takasaki, T., Ikeda, N., Estimating age of humans based on telomere shortening, *Forensic Sci. Int.*, **126** (2002) 197-199.

[10] Bodnar, A.G., Ouellette, M., Frolkis, M., Holt, S.E., Chiu, C., Morin, G.B., Harley, C.B., Shay, J.W., Lichsteiner, S., Wright, W.E., Extension of life span by introduction of telomerase into normal human cells, *Science*, **279** (1998) 349-352.

[11] Bae, K.H., The medicinal plants of Korea, Kyo-Hak Publishing, Seoul, Korea, 2000.

[12] Hwang, J.O., Ahn, D.K., Woo, E.R., Kim, H.J., Seo, S.H., Park, H., Studies on the antibacterial constituents of Baenongtang, *Nat. Prod. Sci.*, **4** (1998) 130-135.

[13] Wang, M., Jin, Y., Ho, C.T., Evaluation of resveratrol derivatives as potential antioxidants and identification of a reaction product of resveratrol and 2,2-diphenyl-1-picrylhydrazyl radical, *J. Agric. Food Chem.*, **47** (1999) 3974-3977.

[14] Redwane, A., Lazrek, H.B., Bouallam, S., Markouk, M., Amarouch, H., Jana, M., Larvicidal activity of extracts from *Quercus lusitania* var. *infectoria* galls (Oliv.), *J. Ethnopharmacol.*, **79** (2002) 261-263.

[15] Markham, K.R., Ternai, B., Stanley, R., Geiger, H., Mabry, T.J., Carbon-13 NMR studies of flavonoids-III: Naturally occurring flavonoid glycosides and their acylated derivatives, *Tetrahedron*, **34** (1978) 1389-1397.

[16] Nicollier, G., Thompson, A.C., Flavonoids of *Desmanthus illinoensis*, *J. Nat. Prod.*, **46** (1983)

112-117.

[17] Davis, A.L., Cai, Y., Davies, A.P., Lewis, J.R., <sup>1</sup>H- and <sup>13</sup>C-NMR assignments of some green tea polyphenols, *Magn. Reson. Chem.*, **34** (1996) 887-890.

[18] Inoshiri, S., Sasaki, M., Kohda, H., Oztuka, H., Yamasaki, K. Aromatic glycosides from *Berchemia racemosa*, *Phytochemistry*, **26** (1987) 2811-2814.

[19] Na, M.K., An, R.B., Lee, S.M., Min, B.S., Kim, Y.H., Bae, K.H., Kang, S.S., Antioxidant compounds from the stem bark of *Sorbus commixta*, *Nat. Prod. Sci.*, **8** (2002) 26–29.

[20] Park, E.J., Nan, J.X., Kim, J.Y., Kang, H.C., Choi, J.H., Lee, S.J., Lee, B.H., Kim, S.J., Lee, J.H., Kim, Y.C., Sohn, D.H., 2000. The ethanol-soluble part of a hot-water extract from *Artemisia iwayomogi* inhibits liver fibrosis induced by carbon tetrachloride in rats, *J. Pharm. Pharmacol.*, **52** (2000) 875-881.

[21] Kobayashi, S., Takehana, M., Itoh, S., Ogata, E., Protective effect of magnesium-L-ascorbyl-2-phosphate against skin damage induced by UVB irradiation, *Photochem. Photobiol.*, **64** (1996) 224-228.

[22] Furumoto, K., Inoue, E., Nagao, N., Hiyama, E., Miwa, N., Age-dependent telomere shortening is slowed down by enrichment of intracellular vitamin C via suppression of oxidative stress, *Life Sci.*, **63** (1998) 935-948.

[23] Cos, P., Calomme, M., Sindambiwe, J.B., Bruyne, T.D., Cimanga, K., Pieters, L., Cytotoxicity and lipid peroxidation-inhibiting activity of flavonoids, *Planta Med.*, **67** (2001) 515-519.

- [24] Heim, K.E., Tagliaferro, A.R., Bobilya, D.J., Flavonoid antioxidants: chemistry, metabolism and structure-activity relationships, *J. Nutr. Biochem.*, **13** (2002) 572-584.
- [25] Black, H.S., deGrujil, F.R., Forbes, P.D., Cleaver, J.E., Ananthaswamy, H.N., deFabo, E.C., Ullrich, S.E., Tyrrell, R.M., Photocarcinogenesis: an overview, *J. Photochem. Photobiol.*, **B40** (1997) 29-47.
- [26] Valenzuela, A., The biological significance of malondialdehyde determination in the assessment of tissue oxidative stress, *Life Sci.*, **48** (1991) 301-309.
- [27] Lorenz, M., Saretzki, G., Sitte, N., Metzcow, S., von Zglinicki, T., BJ fibroblasts display high antioxidant capacity and slow telomere shortening independent of hTERT transfection, *Free Radic. Biol. Med.*, **31** (2001) 824-831.
- [28] Saretzki, G., von Zglinicki, T., Replicative aging, telomeres, and oxidative stress, *Ann. N. Y. Acad. Sci.*, **959** (2002) 24-29.
- [29] von Zglinicki, T., Oxidative stress shortens telomeres, *Trends Biochem. Sci.*, **27** (2002) 339-344.



## Legend of Figures

**Fig. 1.** Elongating effect of the *C. chinensis* extract and its active constituents, piceatannol (**5**) and myricetin (**9**), on the life-span in the HEK-N/F cells. Cellular life-spans were elongated from PDL 14.1 for control to PDL 17.0, 29.9 and 23.2, respectively, for the *C. chinensis* extract, **5** and **9**-added cells.

**Fig. 2.** Dependence of the telomeric DNA length on the cellular ages (PDL) of the HEK-N/F cells administered with or without the *C. chinensis* extract, piceatannol (**5**) and myricetin (**9**). Mean TRF length is estimated as a center of mass and expressed in kbp based on the following equation:  $\Sigma(MW_i \times ODi) / \Sigma(ODi)$ , where  $ODi$  is densitometric output and  $MW_i$  is the length of the DNA at position  $i$ .




Fig. 1




Fig. 2

**Table I.** Antioxidant activity of the *C. chinensis* extract and its constituents on the free radicals and lipid peroxidation

<b>Compounds</b>	<b>DPPH radical scavenging activity IC<sub>50</sub> (μg/ml)</b>	<b>Superoxide radical scavenging activity IC<sub>50</sub> (μg/ml)</b>	<b>Lipid peroxidation inhibitory activity IC<sub>50</sub> (μg/ml)</b>
EtOH extract	35.6 ± 2.1	39.1 ± 4.0	12.5 ± 1.6
<i>Chalcones</i>			
Isoliquiritigenin (1)	> 50	47.6 ± 4.0	3.33 ± 0.50
Liquiritigenin (2)	> 50	49.8 ± 3.1	12.81 ± 1.86
2',4'-Dihydroxy-4-methoxychalcone (3)	> 50	59.2 ± 3.8	6.05 ± 0.99
<i>Stilbenes</i>			
Resveratrol (4)	39.5 ± 2.8	45.3 ± 1.9	0.89 ± 0.10
Piceatannol (5)	20.9 ± 1.3	12.1 ± 1.7	0.09 ± 0.01
<i>Phenolics</i>			
Gallic acid (6)	5.1 ± 0.4	34.1 ± 2.4	6.80 ± 0.63
Methyl gallate (7)	5.3 ± 0.3	16.5 ± 1.4	7.05 ± 0.87
Ethyl gallate (8)	7.0 ± 1.0	15.8 ± 1.6	7.01 ± 0.62
<i>Flavonols</i>			
Myricetin (9)	7.3 ± 0.3	12.1 ± 1.1	0.95 ± 0.06
Afzelin (10)	33.4 ± 1.6	50.2 ± 3.4	10.25 ± 0.91
Quercitrin (11)	12.4 ± 0.6	32.1 ± 2.0	6.21 ± 0.40
Myricitrin (12)	11.6 ± 0.4	29.7 ± 1.8	5.27 ± 0.32
Myricetin-3-O-(2''-O-galloyl)-α-L-rhamnopyranoside (13)	8.6 ± 0.7	13.2 ± 2.5	4.73 ± 0.41
Syringetin-3-O-rutinoside (14)	35.1 ± 1.5	45.2 ± 2.2	11.0 ± 1.2
<i>Flavanols</i>			
(+)-Catechin (15)	15.6 ± 0.8	16.5 ± 2.0	4.71 ± 0.26
(-)-Epicatechin-3-O-gallate (16)	6.8 ± 0.5	11.9 ± 2.1	2.90 ± 0.06
(-)-Epigallocatechin-3-O-gallate (17)	6.7 ± 0.4	24.0 ± 3.5	1.00 ± 0.08
<i>Lignans</i>			
(-)-Lyoniresinol 3a-O-β-D-xylopyranoside (18)	45.7 ± 4.0	>100	37.42 ± 2.06
(+)-Lyoniresiol 3a-O-β-D-glucopyranoside (19)	42.6 ± 3.1	>100	39.10 ± 3.11
α-tocopherol	25.4 ± 0.9		6.61 ± 0.95
Caffeic acid		11.0 ± 1.8	
BHA	15.3 ± 0.6	48.8 ± 2.5	0.11 ± 0.02

Values are expressed as mean ± SD of three replicates.

**Table II.** Cytoprotective effect of the *C. Chinensis* extract and its constituents on *t*-BuOOH-induced oxidative damage

<b>Compounds</b>	<b>Cell viability (%)</b>	<b>TBARS (pmol/mg protein)</b>
Blank	100	370.2 ± 51.2**
Control	11.2 ± 1.2	8721.5 ± 809.5
EtOH extract	53.5 ± 3.8**	984.0 ± 95.2**
<i>Chalcones</i>		
Isoliquiritigenin (1)	13.9 ± 2.6	5118.1 ± 410.5*
Liquiritigenin (2)	16.2 ± 3.2*	3449.7 ± 305.8**
2',4'-Dihydroxy-4-methoxychalcone (3)	12.1 ± 1.0	8255.1 ± 576.4
<i>Stilbenes</i>		
Resveratrol (4)	18.1 ± 1.6*	2792.4 ± 259.1**
Piceatannol (5)	84.7 ± 6.9**	520.2 ± 60.7**
<i>Phenolics</i>		
Gallic acid (6)	41.5 ± 3.1**	1245.7 ± 112.4**
Methyl gallate (7)	43.0 ± 5.6**	1024.4 ± 91.9**
Ethyl gallate (8)	43.1 ± 4.1**	1115.6 ± 96.2**
<i>Flavonols</i>		
Myricetin (9)	61.0 ± 4.5**	810.7 ± 77.0**
Afzelin (10)	16.8 ± 1.3*	3295.1 ± 304.5**
Quercitrin (11)	21.4 ± 1.9*	2217.2 ± 200.3**
Myricitrin (12)	25.4 ± 1.7*	1852.6 ± 126.8**
Myricetin-3-O-(2''-O-galloyl)- $\alpha$ -L-rhamnopyranoside (13)	45.1 ± 3.6**	1019.8 ± 154.0**
Syringetin-3-O-rutinoside (14)	20.0 ± 2.6*	2928.1 ± 209.9*
<i>Flavanols</i>		
(+)-Catechin (15)	24.5 ± 1.9*	1912.4 ± 112.0**
(-)-Epicatechin-3-O-gallate (16)	49.4 ± 6.4**	1007.3 ± 95.2**
(-)-Epigallocatechin-3-O-gallate (17)	46.6 ± 5.9**	1032.6 ± 103.5**
<i>Lignans</i>		
(-)-Lyoniresinol 3a-O- $\beta$ -D-xylopyranoside (18)	13.5 ± 2.0	5981.1 ± 412.6*
(+)-Lyoniresinol 3a-O- $\beta$ -D-glucopyranoside (19)	12.9 ± 2.1	7635.7 ± 631.2

Values are expressed as mean  $\pm$  SD of three replicates.

\* P < 0.05, \*\* P < 0.001, were significantly different from the group treated *t*-BuOOH.

**Table III.** Effect of the *C. chinensis* extract and piceatannol (5) on UVB-induced lipid peroxidation in

SKH-1 mice

Sample treatment (mg/kg)		Amount of TBARS (nmol/mg protein)
Control		0.32 ± 0.05 <sup>*</sup>
UVB irradiated		0.68 ± 0.10
<i>C. chinensis</i> extract	50	0.21 ± 0.10 <sup>*</sup>
+ UVB irradiated	30	0.34 ± 0.09 <sup>*</sup>
	10	0.64 ± 0.10
Piceatannol (5)	30	0.07 ± 0.03 <sup>**</sup>
+ UVB irradiated	10	0.16 ± 0.05 <sup>*</sup>

Values are expressed as mean ± SD.

<sup>\*</sup> P < 0.05, <sup>\*\*</sup> P < 0.001, were significantly different from the UVB irradiated group.