

[PB4-2] [04/18/2003 (Fri) 09:30 - 12:30 / Hall P]

Effects of Ginsenoside Rg1 on the Expression of TNF- α from Rat microglia

Joo SeongSoo, Won TaeJoon, Lee Dolk^o

Department of Immunology, College of Pharmacy, Chung-Ang University, Seoul 156-756, Korea

Microglial cell can act for phagocytosis against abnormal particles in brain, which means that beta-amyloid produced from APP(amyloid precursor protein) can be phagocytosed by microglia when released. In contrast, when senile plaque has already been formed in brain cortex and hippocampal region, microglia can also accelerate the AD pathogenesis due to chronic inflammatory action, which lead to neuron cell cytotoxicity. In our study, we investigated the degree of activation of microglia by using Rg1. For the study, we selected TNF- α released from microglia. Experimental groups were separated in two, one for beta-amyloid and the other for non-beta-amyloid group. As a result, Rb1 showed highest release of TNF- α at 48 hour. In non-beta-amyloid group, Rg1 showed increased release of TNF- α at 0.1 μ M, 100 μ M. In addition, A β group showed that Rg1 suppresses TNF- α at 100 μ M. In conclusion, Rg1 may play a certain role in treatment and prevention of AD in a way to suppress the immune reaction in microglia adjacent around the neuron cell.

[PB4-3] [04/18/2003 (Fri) 09:30 - 12:30 / Hall P]

Effect of Ginsenoside Rg3 in Mouse Hematopoietic Cells

Joo SeongSoo, Park JeongHwan, Lee Dolk^o

Department of Immunology, College of Pharmacy, Chung-Ang University, Seoul 156-756, Korea

Rg3 is a derivative of triterpenoid dammarane, which originally extracted from Red Ginseng, which have been known to have neuroprotective, vasodilator, antioxidative, antimetastasis, and direct anticancer effects. These various backgrounds of Rg3 can provide an additional interest in respect to the "hematopoiesis" in bone marrow and spleen cells. We, therefore, have investigated what effects and correlates of Rg3 (e.g. suppression and side effects) are affected in relation with the bone marrow and spleen cells of mouse. For this study, we designed to know how Rg3 affects the process of hematopoiesis in stem cell level. From the study, we concluded that Rg3 controls the growth and differentiation of the immune cells through increase of the hematopoietic cells, whereas Rg3 reduced and rather mostly controlled the side effect of cyclophosphamide (CTX) in over a studied concentration. In conclusion, Rg3 can reduce the cytotoxicity affecting hematopoietic system inducible from anticancer agents as well as direct anticancer effects. IN addition, good potential of hepatopoiesis is expected to give an breakthrough information in clinical use when use Rg3 as an adjuvant agent both in radio- and chemotherapy.

[PB4-4] [04/18/2003 (Fri) 09:30 - 12:30 / Hall P]

Korean Propolis enhances both the presentation of DC and macrophage activation

Han ShinHa^o, Yun YunHa, Song YoungCheon, Lee SookYeon, Ha Nam-Joo, Kim KyungJae