

2-Hydroxymuconic semialdehyde dehydrogenase catalyzes the conversion of 2-hydroxymuconic semialdehyde (HMS) to an enol form of 4-oxalocrotonate which is a step in the catechol–meta cleavage pathway. A tomC gene encoding 2-HMS dehydrogenase of Burkholderia cepacia G4, a soil bacterium that can grow on toluene, cresol, phenol or trichloro ethylene, is identified in between catechol 2,3-dioxygenase gene and HMS hydrolase gene, its sequence is analysed and the enzyme is characterised. The 2-HMS dehydrogenase from B. cepacia G4 is able to oxidize many meta cleavage products (MCPs) in the presence of NAD<sup>+</sup>. The oxidative activity of the enzyme is spectrophotometrically detected with the observation of a gradual time dependent decrease in absorbance at A375 in the presence of 2-HMS and NAD<sup>+</sup>. All the tested MCPs served as substrates for 2-HMS dehydrogenase and thus it is evident that the enzyme has broader substrate specificity. But the activity of the enzyme is much lower for both 3-methyl HMS and 2-hydroxy 6-oxo 6-phenylhexa 2,4-dienoic acid (HOPDA) when compared with 2-HMS, 4-methyl HMS and 4-chloro HMS. Km and Vmax of 2-HMS dehydrogenase for 2-HMS, 4-Methyl HMS, 4-chloro HMS and for NAD<sup>+</sup> are calculated from Lineweaver–Burk plots. The kinetic parameters showed that the enzyme has high catalytic efficiency in terms of Vmax/Km towards 4-methyl HMS followed by 2-HMS but very low for 4-chloro HMS. On the other hand the catalytic efficiency of the enzyme is overall low for NAD<sup>+</sup> in the presence of different MCPs. The sequence analysis shows the open reading frame (ORF) corresponding to tomC consists of 1458 base pairs with ATG initiation codon and TGA termination codon. This gene can encode a polypeptide of molecular weight 52 KDa containing 485 amino acid residues. The deduced amino acid sequence of 2-HMS dehydrogenase encoded by tomC gene from B. cepacia G4 exhibited a highest 78% homology with that of corresponding enzyme encoded by aphC gene of Comamonas testosteroni, 64%–78% homology with those of reported HMS dehydrogenases, and 29%–70% homology with those of different kinds of dehydrogenases. From the alignment of amino acid sequence putative cofactor NAD<sup>+</sup>-binding regions and catalytic residues were identified. The 2-HMS dehydrogenase from B. cepacia showed significant phylogenetic relationship not only with the same enzyme from other bacteria, also with different dehydrogenases from evolutionarily distant organisms.

[PC1–23] [ 04/18/2003 (Fri) 09:30 – 12:30 / Hall P ]

### Differential regulation of gene expression by RNA polymerase II in response to DNA damage

Heo JeongHwa<sup>o</sup>, Han JeungWhan, Lee HyangWoo, Cho EunJung

Lab. of Biochem. & Mol. Biol., College of Pharmacy, Sungkyunkwan University

RNA polymerase II (pol II) is known to cycle between hyperphosphorylated and hypophosphorylated forms during transcription cycle. These extensive phosphorylation/dephosphorylation event occurs in the C-terminal domain (CTD) of the largest subunit of pol II which consists of a tandemly repeated heptapeptide motif with consensus of YSPTSPS. Since different phosphorylation pattern of CTD is the hallmark of transcription regulation, we asked whether changes in CTD phosphorylation is involved in global regulation of transcription in response to DNA damage.

Taking the advantage of chromatin immunoprecipitation assay, we have demonstrated that pol II or pol II with nonphosphorylated CTD uniformly associated with a transcribed gene from a promoter to a coding region. While TFIIF (Tfg2) or TFIIF (Kin28) associated predominantly at a promoter region. With several antibodies against selected transcription factors and also antibodies against different CTD phosphorylation epitopes, we describe the response of transcription complex in terms of CTD phosphorylation and its role in gene expression upon DNA damage induced by various sources.

[PC1–24] [ 04/18/2003 (Fri) 09:30 – 12:30 / Hall P ]

## PROTECTION OF PARAOXONASE BY LIPIDS

Nguyen-Duy Su<sup>o</sup>, Kim Ju Ruyng, Sok Dai Eun

College of Pharmacy, Chungnam National University, Yuseong-ku, Taejeon 305-764, KOREA

Effect of lipids on the oxidative inactivation of paraoxonase (PON1), a HDL-associated antioxidant protein, was investigated, based on the hydrolysis of phenyl acetate. First, various fatty acids were examined for the prevention against the inactivation of PON1 by ascorbate/Cu<sup>2+</sup> system, which caused a remarkable ( $\geq 90\%$ ) inactivation of PON1, accompanied by a partial loss of histidine residues. Compared to saturated fatty acid (C6-C18) exhibiting a modest protection (9-40%), monounsaturated fatty acids showed a greater protection ( $E_{max}$ , 70-82%). In addition, a remarkable protection was also expressed by conjugated linoleic acid, but not  $\alpha$ -linoleic acid. The most protective was oleic acid ( $EC_{50}$ , 2.7 mM), which fully prevented the oxidative loss of histidine residues of PON1. Noteworthy, either the introduction of hydroxyl group at C12 of oleic acid or the derivatization of carboxylic group caused the loss of protective action, emphasizing the importance of both acyl chain and anionic charge. Consistent with this, dioleoylphosphatidylglycerol, negatively-charged, was more protective than other phospholipids including dioleoylphosphatidylcholine. Similarly, such a protective action of lipids was also observed in different types of inactivation of PON1, where ascorbate/Fe<sup>2+</sup>, peroxides, or p-hydroxymercuribenzoate were employed. The same pattern of protection by lipids was also observed when PON1 activity was determined using paraoxon as substrate. These results suggest that some lipids play a beneficial role in maintaining PON1 activity from oxidative stress in vivo.

[PC1-25] [ 04/18/2003 (Fri) 09:30 - 12:30 / Hall P ]

### Proteomic analysis of proteins Secreted by Human Bronchial Epithelial Cells in Response to Pathogenic Bacterial Infections

Oh MiJung<sup>o</sup>, Park MiJa, Lee JiYeon, Park JiWoo, Lee NaGyong, Jung SungYun, Kim DaeKyong

college of pharmacy, Chung-ang univ. and Department of Bioscience, Sejong Univ.

Bacterial infection is a very complex process in which both pathogens and host cells play crucial roles, and the host cells undergo drastic changes in their physiology, releasing various proteins in response to the pathogenic infection. Human airway epithelial surface serves as a first line of defense against microorganisms and the external environment. It is well known that bronchial epithelial cells secrete various chemokines and cytokines such as IL-6 and IL-8 to cope with various respiratory pathogens. Under infectious conditions, cultured mammalian cells release a number of proteins to its culture medium. While most of these proteins localized in intracellular compartments result from cell death, some of the proteins are known to serve as immunologically responsive proteins, such as interferons, interleukins, colony stimulating factors (CSFs), transforming growth factor- $\beta$  (TGF- $\beta$ ). Although many kinds of these cytokine proteins are identified and characterized for their biological roles, such cytokine-like proteins as a functionally unknown protein could be found through high-throughput identification of the proteins in the extracellular space. In this study, we used a combined technology of column chromatography and display proteomics to identify such proteins from pathogenic bacteria-infected human bronchial epithelial (BEAS-2B) cells. Using some of ion-exchange column chromatographies, two-dimensional gel electrophoresis, we found at least twenty different proteins that specifically secreted to the culture medium in the response to the bacteria infection. While some of these proteins were revealed as known cytokines and other functional