

compounds (1,2-Dimethyl-3-phenylaziridine, amphetamine oxime, ephedrine, N-formylmethamphetamine, N-acetylmethamphetamine, acetylephedrine, 3,4-dimethyl-5-phenyl-2-oxazolidone, methamphetamine dimer) are identified impurities in illicit methamphetamine and the identity of the impurity is conformed synthesis. Identification of other impurities found on the chromatogram is under investigation. These impurities revealed that most of the seized methamphetamine in Korea was synthesized from ephedrine as a starting material.

[PD4-27] [ 04/18/2003 (Fri) 13:30 - 16:30 / Hall P ]

### MODULATION OF THE ACTIVITY OF PRO-INFLAMMATORY ENZYMES, COX-2 AND iNOS, BY CHRYSIN DERIVATIVES

Cho Heeyeong<sup>o</sup>, Yun Cheol-Won, Kong Jae-Yang, Kim Kyoung Soon, Park Youmie, Lee Sanghyun, Kim Bak-Kwang

<sup>1</sup>Pharmaceutical Screening Team, Korea Research Institute of Chemical Technology, Taejeon 305-600, South Korea and <sup>2</sup>College of Pharmacy, Seoul National University, San 56-1, Shillim-Dong, Kwanak-Gu, Seoul 151-742, South Korea.

Chrysin, a natural flavone compound contained in plants, has anti-inflammatory activity. Its anti-inflammatory effect has been previously explained in part by the suppression of promoter activities of inducible pro-inflammatory enzymes (cyclooxygenase-2 (COX-2) and inducible nitrogen synthase (iNOS)). Nitrate production triggered by the activation of lipopolysaccharides (LPS) was most highly suppressed by the treatment of chrysin, followed by 5-hydroxy-7-methoxyflavone (Ch-2), 5,7-diacetylflavone (Ch-4) in cultured Raw 264.7 cells. Here we tested the inhibitory activity of chrysin derivatives on COX-2 and iNOS enzymes. Interestingly COX-2 enzyme was strongly inhibited by Ch-2 (IC<sub>50</sub> = 2.7 mM) but not by other derivatives. Furthermore, the inhibition of COX enzyme by Ch-4 was selective on COX-2 isoenzyme over COX-1. The three-dimensional interaction models of Ch-4 complexing with COX-2 showed comparatively good fit into the topology of the binding pocket. There was a hydrogen bond between oxygen of ketone group of the 7-position of Ch-4 and the hydroxyl group of Tyr355. Docked conformation of Ch-4 into V523I mutant of COX-2 indicated that Ile523 of COX-1 might contribute to the selectivity of COX-2/COX-1. Ch-4 showed no effect on iNOS activity. Chrysin and Ch-2 weakly inhibited iNOS enzyme activity in hemoglobin assay. However, the underlying mechanisms of the inhibition of iNOS by chrysin are not clear.

[PD4-28] [ 04/18/2003 (Fri) 13:30 - 16:30 / Hall P ]

### In vitro metabolism of a new protective agent, KR-31543 in human liver microsomes

Ji HyeYoung<sup>o</sup>, Kim SookJin, Lee Hongll, Lee SeungSeok, Lee DongHa, Lim Hong, Lee HyeSuk

College of Pharmacy, Wonkwang University; AgroPharma Research Institute, DonbuHannong Chemical Co

The purpose of this paper was to identify the metabolic pathway of a new neuroprotective agent, KR-31543 for ischemia-reperfusion damage in human liver microsomes and characterize cytochrome P450 (CYP) enzymes involved in the in vitro metabolism of KR-31543. KR-31543 generates two metabolites in human liver microsomes: M1, N-(4-chlorophenyl)-N-(2-methyl-2H-tetrazol-5-ylmethyl)amine and M2, hydroxy-KR-31543. From a combination of chemical inhibition, immunoinhibition, correlation analysis in human liver microsomes and metabolism by expressed recombinant CYP enzymes, KR-31543 is metabolized predominantly by CYP3A4 and is mainly converted to M1, N-(4-chlorophenyl)-N-(2-methyl-2H-tetrazol-5-ylmethyl)amine. KR-31543 was found to be a potent inhibitor of human CYP2D6 and 3A4 in human liver

microsomes.

[PD4-29] [ 04/18/2003 (Fri) 13:30 - 16:30 / Hall P ]

**Alternation of plasma F<sub>2</sub>-isoprostanes before and after hemodialysis in patients with atherosclerosis in end-stage renal disease**

Kim KyungMee<sup>o</sup>, Jung ByungHwa, Kim SangWook, Paeng KiJung, Chung BongChul

Korea Institution of Science and Technology, Yonsei University, Korea University

F<sub>2</sub>-isoprostanes are derived *in vivo* principally from the formation of positional peroxy radicals of arachidonic acid, endocyclization to prostaglandin G<sub>2</sub>-like structures, and reduction to PGF<sub>2</sub>-like compounds. F<sub>2</sub>-isoprostanes have been proposed as biomarkers of lipid peroxidation, oxidative stress status, and oxidation of low density lipoprotein (LDL) in atherogenesis. Especially, elevated 8-iso-PGF<sub>2a</sub> is known as a useful clinical biomarker of oxidative stress in end-stage renal disease (ESRD) patients receiving hemodialysis (HD). Therefore, we studied about alternation of plasma F<sub>2</sub>-isoprostanes on before and after hemodialysis in patients with atherosclerosis in ESRD using gas chromatography-tandem mass spectrometry (GC/MS/MS). Plasma concentrations F<sub>2</sub>-isoprostanes were significantly higher in post-hemodialysis than pre-hemodialysis in atherosclerosis ( $P < 0.05$ ). This may suggest that F<sub>2</sub>-isoprostanes is affected with oxidative stress by hemolysis treatment in atherosclerosis.

[PD4-30] [ 04/18/2003 (Fri) 13:30 - 16:30 / Hall P ]

**Methyl carboxylates of naphtho[1,2-b]furan and naphtho[1,2-b]pyran from the roots of *Rubia akane***

Jung SoonJa<sup>o</sup>, Jung JiHyun, Cao LiHua, Shin YoungHee, Son JongKeun\*, Lee ChongSoon\*\*, Woo MiHee

College of Pharmacy, Catholic University of Daegu, \*College of Pharmacy, Yeungnam University, \*\*Department of Biochemistry, Yeungnam University.

*Rubia akane* (Rubiaceae) is a perennial vine plant and generally grows at the mountain, field and fence of Korea, Japan, China and Taiwan. This plant grows about two meters and has empty stem and red roots. The roots of this plant have been used as a Korean traditional medicine for the treatment of cough, the stone in the bladder and kidney, inflammation of the joints, uterine hemorrhage and uteritis.

From the screening experiment, *Rubia akane* exhibited anti-cancer activity. MeOH extract of the roots of this plant was subsequently fractionated into four parts: methylene chloride, ethyl acetate, n-butanol and water fractions. Methylene chloride fraction showed cytotoxicity against HT-29 and MCF-7 cell lines, and DNA Topoisomerase I inhibitory activity at the concentration of 5  $\mu\text{g}/\text{ml}$  and DNA Topoisomerase II inhibitory activity at the concentration of 50  $\mu\text{g}/\text{ml}$ . Ethyl acetate fraction showed DNA Topoisomerase I inhibitory activity at the concentration of 5  $\mu\text{g}/\text{ml}$  and DNA Topoisomerase II inhibitory activity at the concentration of 50  $\mu\text{g}/\text{ml}$ .

We isolated methyl carboxylates of naphtho[1,2-b]furan and naphtho[1,2-b]pyran from the methylene chloride fraction.