

P70

## Identification of Solvent Stress Gene in *Pseudomonas* sp. BCNU 106 by Differential Display Reverse Transcription-PCR

Yun Ui Bae<sup>1</sup>, Ju Soon Yoo<sup>1</sup>, Ki Wook Kim<sup>1</sup>, Su Dong Cho<sup>3</sup>,  
Ja Young Moon<sup>1,4</sup>, Dong-Wan Kim<sup>5</sup>, Young Kee Jeong<sup>5</sup>,  
and Woo Hong Joo<sup>1,2\*</sup>

<sup>1</sup>Institute of Genetic Engineering, <sup>2</sup>Department of Biology,  
<sup>3</sup>Institute for Basic Science, <sup>4</sup>Department of Biochemistry and Health Sciences,  
<sup>5</sup>Department of Microbiology, Changwon National University, Changwon, Korea  
<sup>6</sup>Department of Microbiology, Dong-Eui University, Pusan, Korea

*Pseudomonas* sp. BCNU 106 was isolated from waste water. The strain is a solvent-tolerant strain able to grow in a culture medium containing more than 50%. Solvent-tolerant strain *Pseudomonas* sp. BCNU 106 was studied to identify solvent-tolerant genes. Using the mRNA differential display technique, a total of 60 differentially expressed cDNA fragments were identified by comparing band intensities among fingerprints obtained from Luria-Bertani(LB) medium with 10% toluene. The mRNA expression levels were detected after 10%(v/v) toluene treatments for 24hrs. Transcripts induced under 10% toluene exhibited homologies to; manganese superoxide dismutase, catechol-o-transferase, catalase isozyme CatA gene, beta-hexosaminidase, putative ABC transporter, transcriptional regulator, heat-shock protein, and toluene dioxygenase. Especially, the toluene dioxygenase, one of the enzyme well known in the toluene degradation pathway of *Pseudomonas putida* was identified under toluene treatment. Therefore, toluene-responsive *Pseudomonas* sp. BCNU 106 transcripts isolated using mRNA differential display showed to encode proteins with related to antioxidation and toluene degradation pathway.