

[P-55]

Effects of Rhus Verniciflua Strokes Extracts and its Components on the Proliferation, Collagen Synthesis, and the Mrna Level of Hepatic Fibrosis Related Proteins in Rat Hepatic Stellate Cells.

Seung Hyun Jung, Jeong-Ran Kim, Chun-Soo Na*, Bum Rak Choi*, Sun Young Yoon*,
Young In Park, and Mi-Sook Dong

*School of Lifesciences and Biotechnology, Korea University, 136-701, Seoul, Korea and**
Lifetree Biotech Co., Ltd, Kyonggido, 441-350

Hepatic stellate cells (HSC) and the derived myofibroblasts are known to play a central role in liver fibrosis. Rhus verniciflua Strokes (RVS) has traditionally been used in Korea herbal medicine for a stomachic tonic. In this study, we observed the effect of RVS acetone extract (Ra) on the proliferation, the collagen synthesis, and hepatic fibrosis related proteins mRNA levels in HSC-T6 cells which is a fully activated rat hepatic stellate cell line. Ra inhibited the proliferation and decreased the content of collagen in the HSC-T6 cells. The mRNA levels of TGF β 1, Timp-1 and procollagen 1 α 1 were significantly reduced by Ra treatment. We isolated the active components of Ra by column chromatography, HPLC and determined five structures which are butein, fustin, sulfuretin, ficetin and 3,4-dihydroxybenzoic acid by ¹H-NMR and ¹³C-NMR. The anti-fibrotic activity of each component of Ra was not excellent in compared with that of total Ra judging by collagen excretion and mRNA levels of hepatic fibrosis related proteins. Collectively, Ra inhibited hepatic stellate cell proliferation and collagen synthesis that might have a protective role against liver fibrosis.

Keyword : Rhus verniciflua Strokes, antihepatofibrotic effect, rat hepatic stellate cell