## Rapid screening system of reassortant influenza A viruses using primer dependent multiplex RT-PCR

Hyun Ah. Kim<sup>1</sup>, Suk-Hoon Ha<sup>1,2</sup>, Yeon Hee Kim<sup>1</sup>, Kwang Hee Lee<sup>2</sup>, Nam J. Lee<sup>1</sup>, Jae Seoung Kim<sup>1</sup>, Wan Je Park<sup>1</sup>, and Baik Lin Seong<sup>2</sup>

<sup>1</sup>CJ Corporation, 522-1, Dokpyong-Ri, Majang-Myun, Ichon-Si, Kyonggi-Do, 467-810, Korea. TEL: +82-31-639-4732, FAX:+82-31-632-2784

<sup>2</sup>Protheon Inc., Yonsei Engineering Research Center B120E, 134 Shinchon-Dong, Seodaemun-Gu, Seoul 120-749, Korea. TEL: +82-2-2123-2885, FAX:+82-2-2123-7265

## Abstract

A simple method for rapid screening of 6:2 cold adapted reassortnats of influenza virus, primer dependent multiplex RT-PCR screening method, is described. Primers for screening purpose were selected from highly variable regions on an attenuated donor virus in comparison to all influenza A virus from constructing multiple sequence alignments. There are two kinds of primers. One kinds of primers were selected from highly variable regions on an attenuated donor virus in comparison to other wild influenza A viruses. These showed PCR products that were amplified when the template genomes were from donor strain. The others were designed on conserved region of many kinds of influenza A viruses. These were used for the presence of each segment although there were many failures of producing PCR products from wild type virus gene were observed. As a result, the difference of primer binding affinity made it possible to show the nucleotide difference between donor strains and wild viruses and to facilitate for screening of 6:2 reassortant vaccine strain. In multiplex RT-PCR format, the process of this method is highly streamlined since time consuming post-PCR analysis is omitted. We demonstrated the utility of this method by genotyping of randomly picked reassortant viruses.

**Key words:** Influenza virus; Reassortant virus; Primer-dependent multiplex RT-PCR screening system.

## References

- Cha, T. A., J. Zhao, E. Lane, M. A. Murray, and D. S. Stec, Determination of the genome composition of influenza virus reassortants using multiplex reverse transcription polymerase chain reaction followed by fluorescent single-strand conformation polymorphism analysis. 1997, Anal. Biochem. 252, 24-32.
- Copper, L. A. and K. Subbarao, A simple restriction fragment length polymorphism-based strategy that can distinguish the internal genes of human H1N1, H3N2, and H5N1 influenza viruses. 2000, J. Clin. Microbiol. 38, 2579-2583.
- 3. Cox, N. J. and C. A. Bender, The molecular epidemiology of influenza viruses. 1995, *Semin. Virol.* **6**, 359-370.
- 4. Munoz, M. F. and R. B. Couch, Current research on influenza and other respiratory viruses. 2000, *Antiviral Res.* 46, 91-124.
- Hay, A. J., A. R. Bellamy, G. Abraham, J. J. Skehel, C. M. Brand, and R. G. Webster, Procedure for characterization of genetic material of candidate vaccine strains. 1977, Dev. Biol. Stand. 39, 15-24.
- Klimov, A. J. and N. J. Cox, PCR restriction analysis of genome composition and stability of cold-adapted reassortant live influenza vaccines. 1995, J. Virol. Methods. 52, 41-49.
- Lamb, R. A. and R. M. Krug, Orthomyxoviridase: the viruses and their replication. 1996, p. 1353-1395. In B. N. Fields, D. M. Knipe, P. M. Howley, R. M. Chanock, J. L. Melnick, T. P. Monath, and B. Roizman (ed.), *Fields Virology*, 3rd ed. Lipp incott-Raven Publishers, Philadelphia, Pa.
- 8. Lee, K. H. and B. L. Seong, Current status for influenza control. 1999, *Biotechnol. bioprocess Eng.* 4, 157-164.
- Lee, K. H., J. W. Youn, H. J. Kim, and B. L. Seong, Identification and characterization of mutations in the high growth vaccine strain of influenza virus A. 2001, Arch. Virol. 146. 369-377.
- 10. Maassab, H. F. and M. L. Bryant, The development of live attenuated cold-adapted influenza virus vaccine for humans. 1999, Rev. Med. Virol. 9, 237-244.
- 11. Offringa, D. P., T. M. Victoria, Z. Ye, and R. A. Levandowski, A comprehensive systematic approach to identification of influenza A virus genotype using RT-PCR and RFLP. 2000, *J. Virol. Methods.* 88, 15-24.

- 12. Palese, P. and J. L. Schulman, Mapping of the influenza virus genome: Identification of the hemagglutination and the neuraminidase genes. 1976, *Proc. Natl. Acad. Sci. USA* 73, 2142-2146.
- Sakamoto, S., Y. Kino, T. Oka, M. L. Herlocher, and H. F. Maassab, Gene analysis of reassortant influenza virus by RT-PCR followed by restriction enzyme digestion. 1996, J. Virol. Methods 56, 61-171.
- Sambrook, J. and D. W. Russell, Mutagenesis. 2001, Molecular Cloning 3rd ed. Cold Spring Harbor Laboratory Press, Cold spring, N. Y, pp. 13.49-13.52.
- 15. Zou, S., C. Stansfield, and J. Bridge, Identification of new influenza B virus variants by multiplex reverse transcription-PCR and the heteroduplex mobility assay. 1997, J. Clin. Microbiol. 26, 1544-1548.
- Wareing, M. D., G. A. Marsh and G. A. Tannock, Preparation and characterization of attenuated cold-adapted influenza A reassortants derived from the A/Leningrad/134/17/57 donor strain. 2002, Vaccine 20, 2082-2090.
- 17. Zou, S., A practical approach to genetic screening for influenza virus variants. 1997, J. Clin. Microbiol. 35, 2623-2627.