

## The Functional Properties of *Kimchi* for the Health Benefits

Yeong-Ok Song

Department of Food Science and Nutrition and Kimchi Institute, Pusan National University,  
Busan 609-735, Korea. E-mail: yosong@pusan.ac.kr

*Kimchi* is a fermented vegetable consumed in Korea every day as a side dish. Now-a-days *kimchi* became a world wide food based upon the scientific evidences showing the health promoting activities of *kimchi*. The beneficial effects of *kimchi* on human health may come from nutrients in *kimchi* such as vitamins, minerals, fibers and photochemical, or from the biological compounds present either in *kimchi* ingredients such as garlic, ginger, red pepper powder, the fermented products of *kimchi*, or from lactic acid bacteria in *kimchi*. The functional properties of *kimchi* responsible for the health benefits have been claimed as follows. 1) Anti-oxidative activity, 2) Anti-aging activity, 3) Anti-mutagenic, anti-tumor activities, 4) Anti-microbial activity, 5) Immune-stimulatory activity, 6) Weight-controlling activity, 7) lipid-lowering activity, 8) Anti-atherogenic activity, etc.

### Anti-Oxidative Activity of *Kimchi*

The antioxidant properties of *kimchi* come from raw materials used to make *kimchi* and from other biological compounds produced during fermentation. Carotenoids, flavonoids, polyphenols, Vt. C, Vt. E, chlorophyll present in *kimchi* ingredients are known to be primary antioxidants and 3-(4'-hydroxyl-3'5'-dimethoxyphenyl)propionic acid in Chinese cabbage *kimchi* was identified as an active principle to have free radical scavenging ability (1). Antioxidant effect of *kimchi* has been confirmed in vitro, in vivo, and clinical studies. *Kimchi* retarded linoleic autooxidation (2) and LDL oxidation (3), and scavenged free radicals (4-6). Many researchers have suggested that anti-oxidative property of *kimchi* is one of the mechanisms for anti-mutagenicity/anticancer, anti-atherogenicity (7,8), and anti-aging (4-6).

### Anti-Aging Activity of *Kimchi*

Free radical theory is the one of the most acceptable hypothesis that explains the aging process. The active oxygen species or lipid radicals react with bio-chemicals in the body result in damaging. Thus animals try to reserve antioxidants in addition to the antioxidative enzymes in the body to get rid of free radicals. *Kimchi* and *kimchi* ingredients enhanced the activities of antioxidative enzymes such as superoxide dismutase, catalase, glutathione reductase/peroxidase as well as increased Vt. E and carotene levels in the plasma and liver (3,9). The concentration of total free radicals and hydroxyl radical in the plasma of elderly who consumed *kimchi* more than mean intakes (112 g) was lower than the elderly consumed *kimchi* less, while GSH and GSH/GSSG were increased. These results informed that *kimchi* might have a role either inhibit the production of free radicals or discard the free radicals more efficiently (10). This observation was confirmed in the animal study. Free radical production in the brain of SAM due to senescence (1 year feeding) was decreased by *kimchi* consumption and the activities of the antioxidative enzymes in the brain were increased (11).

### Anti-Mutagenic, Anti-Tumor Activities of *Kimchi*

The biological compounds in *kimchi* having anti-mutagenic and anti-tumor activities are vitamin C,  $\beta$ -carotene,

phenolic compounds, isothiocyanate, indole compound,  $\beta$ -sitosterol, diallylsulfide, dietary fiber, fermented products, and lactic acid bacteria (12). The chemo-preventive or chemo-therapeutic effect of kimchi may be due to enhanced detoxifying ability of the liver that metabolites the carcinogens into water-soluble compounds to be excreted, or to inhibit the transformation of pro-carcinogen to carcinogen by retarding responsible enzyme activities, or to inhibit the DNA replication and protein synthesis of tumor cells, or to stimulate the apoptosis of the tumor cell. Increased activity of hepatic glutathione S-transferase (GST) classified as phase II enzyme in the detoxification system was observed when kimchi ingredients-radish, Chinese cabbage, garlic, green onion, leek, or kimchi was given to rats. Sulforaphane or S-containing compounds in these materials are partially responsible for the increased activity of GST. Lactic acid bacteria in kimchi is claimed to have antimutagenic and anticancer activity by inactivating or inhibiting the production of carcinogens in gastro-intestinal tract, or stimulating the immune system that retards the carcinogenic process (13). The low incidence of colon cancer in Korea was explained that the activity of  $\beta$ -glucosidase and  $\beta$ -glucuronidase was inhibited by lactic acid bacteria in kimchi arrived at large intestine (23), thus less carcinogen was produced in the colon (14). Both viable and non-viable lactic acid bacteria showed anti-mutagenic ability informing that cell wall components, glycopeptides, of the lactic acid bacteria is responsible for these activities (13,15). The extracts of cell wall of *Lactobacillus plantarum* exhibited a direct anticancer effect against abdominal cancer induced by Sarcoma-180 (15). The short-chain fatty acids produced by lactic acid bacteria in the colon may induce the apoptosis (16,17). And apoptosis by kimchi extracts may occur to stimulate the DNA fragmentation (18) or to arresting the cell cycle. Kimchi extracts also showed anti-tumor activity by enhancing the phagocytic cell activity (19). Beside these effects, Chinese cabbage kimchi, radish kimchi, and yulmu kimchi showed anti-genotoxic effect against DNA damage (20). The active principles in kimchi for anti-cancer were not identified yet however the dichloromethane fraction of Chinese cabbage kimchi showed the highest anti-cancer activity (21).

### **Anti-Microbial Activity of Kimchi**

The S-containing compounds in garlic, green onion, red pepper, and ginger have an antimicrobial activity against pathogens (22) and lactic acid bacteria in kimchi is also known to have an anti-microbial activity. One hundred fifty seven strains of lactic acid bacteria were isolated from home made Chinese cabbage kimchi and 144 out of 157 showed a strong resistance to the artificial gastric juice and bile informing that most of the lactic acid bacteria in kimchi will arrive at the intestine (23). Bacteriocin produced by lactic acid bacteria is responsible for this anti-microbial activity and the fermented environment of kimchi, of pH is between 4.1~4.5, augmented this activity. In addition to this, lactic acid bacteria can absorb the harmful substances and excrete them into the feces. There has been no reports of food born disease related to the kimchi consumption since the anti-microbial action of lactic acid bacteria will destroy the harmful microorganisms exist in the intestine.

### **Immunostimulatory Activity of Kimchi**

Immunostimulatory effects of kimchi have been studied with lactic acid bacteria in kimchi. *Lactobacillus plantarum* isolated from kimchi stimulated the proliferation of splenocytes and Payer's patch, increased the production of NO by peritoneal macrophages, increased the production of intestinal secretory IgA, increased TNF- $\alpha$  and IL-2 concentration of rats, and IgG secretion against sheep red blood cell (24). These immunostimulatory effects of *Lactobacillus plantarum* was also observed in tumor induced rats. The IgA secretion, NO production by macrophage, cytokine production, and phagocytic activity of macrophage were greatly increased (15). Kimchi

extracts showed effects on stimulating the growth of spleen cell, bone marrow cell and thymus cell in the cell culture system (25), on increasing the B cell proliferation in the spleen lymphocytes of rats (26), and on increasing the phagocytic movement of macrophage. Fermented kimchi showed greater effects than fresh kimchi on immune stimulation informing that fermented products including lactic acid bacteria besides might be responsible for these effects.

### **Weight-Controlling Activity of Kimchi**

Kimchi is a natural health food that is low in carbohydrate and lipids, and abundant in vitamins, minerals, fibers, and phytochemicals. The calorie for the Chinese cabbage kimchi is approximately 32 kcal/100 g and that for the soup based radish kimchi is 9 kcal/100 g. Among the nutrients in kimchi, vitamin A, beta-carotene, chlorophyll and flavonoids are from Chinese cabbage, green anion, carrot, red pepper powder, and vitamin C mainly comes from Chinese cabbage and red pepper powder. Ca, Mg, and P are from fermented fish sauce, red pepper powder and oyster. Crude fiber in Chinese cabbage kimchi is approximately less than 10% by dry base (27). The biologically active substances in kimchi ingredients known to have lipid-lowering activities are  $\beta$ -sitosterol in Chinese cabbage, S-methylcysteine sulfoxide and S-allylcystein sulfoxide in garlic, capsaicin in red pepper. Lipolytic activity in adipocytes was observed by red pepper and the activity was increased with the degree of pungency of red pepper suggesting that capsaicin is the compounds responsible for this activity (28). Besides these, many researches have reported that kimchi has a lipid-lowering activity.

The weight reduction of rats by kimchi is positively correlated with the amount of kimchi consumed (27). Rats fed 10% kimchi for 4 weeks showed significant reduction in body weight but this phenomenon was not observed in rats fed less amount of kimchi, suggesting that large dose of Chinese cabbage kimchi intake for long term will show the diet effect (27). The size and number of adipocytes of rats fed high fat diet was reduced by kimchi, especially with fermented kimchi. This research group emphasized that the duration of fermentation was rather important than the amount of kimchi intake on reducing the epididymal fat pad size and adipose cell number (26). In clinical trial, obese girl whose BMI is over 25 consumed 3 g of freeze-dried kimchi for 6 weeks as a pill form, total body fat content was decreased (-5.11%) but changes in the body weight and abdominal fat was not significant (29). This result was in agreement with the observation from the animal study. Thus large dose and long term intake of Chinese cabbage kimchi is recommended for the weight reduction.

### **Lipid Lowering Effect of Kimchi**

The biologically active substances in kimchi ingredients known to have lipid-lowering activities are  $\beta$ -sitosterol in Chinese cabbage, S-methylcysteine sulfoxide and S-allylcystein sulfoxide in garlic, capsaicin in red pepper. And lactic acid bacteria in kimchi are also known to lipid-lowering effect, especially cholesterol lowering activity. The mechanisms for lipid lowering of these compounds have been studied extensively.  $\beta$ -sitosterol is phyto-cholesterol that competes with cholesterol in the intestine for absorption. The alliin in onion enhances the lipolytic activity through the hormonal regulation by increasing the adrenalin and glucagon secretion. Allicin in garlic inhibits cholesterol synthesis by inhibiting the acetyl-CoA synthetase or HMG-CoA reductase activity. Capsaicin stimulates the  $7\alpha$ -hydroxylase activity to convert cholesterol into the bile acids (30), and it increases the energy expenditure via thyroid hormone regulation (31). Certain strain of lactic acid bacteria such as *Lb. acidophilus* can bind the cholesterol in their cell wall besides it can decompose the cholesterol for assimilation and de-conjugates the bile acids (32,33). Based on the accumulated knowledge of individual kimchi ingredients in terms of lipid

lowering activities, kimchi should have hypolipidemic effects. Nutritional survey of kimchi intake and lipid analysis showed that daily kimchi consumption level and HDL cholesterol concentration was positively correlated while LDL cholesterol was negatively correlated in 102 of healthy Korean men aged between 40 to 64 years old (34). Researches about kimchi on lipid lowering effects have been extensively carried out with feeding various kinds of kimchi to the animals and human subjects. The results concluded that kimchi has lipid-lowering effects in the plasma, liver and other organs. Triglyceride, cholesterol, LDL cholesterol in the plasma was decreased and HDL cholesterol was increased in studies with SAM (35), rats (26,27,36) and rabbits (7,8,37,38) fed kimchi. HMG-CoA reductase activity in the liver, CETP activity in the plasma, ACAT activity in the liver of rat and rabbit were decreased by kimchi diet. A pilot-scale clinical study, six nonlipidemic persons with four Caucasians and two Chinese Americans consumed 150 g kimchi (1/2 cup) daily with regular meal for 2 weeks. Lag phase duration of LDL oxidation at 2nd week in Caucasians never been exposed to kimchi was extended two times compared to that of baseline (0 week). But plasma lipid profiles and vitamin C, E, homocystein, in the plasma was not changed. Kimchi supplementation study with 12 middle aged healthy Korean who took 3 g of freeze-dried kimchi as a form of pill every day for 6 weeks showed significant decreased in plasma TG (-16.8%) than placebo group, and atherogenic index (AI) for kimchi group was decreased by 10.8% (39). Lipid lowering activity of kimchi observed in the animal and clinical studies might be due to the inhibition of HMG-CoA reductase, CETP and ACAT activity by biological compound in kimchi identified as 3-(4'-hydroxyl-3',5'-dimethoxyphenyl)propionic acid (1,40).

### **Antiarthergenic Effects of Kimchi**

Oxidized LDL and hypercholesterolemia are one of the major symptoms to develop the atherosclerosis. The functional properties of kimchi or kimchi ingredients such as antioxidant activity, free radical scavenging activity, hypolipidemic activity will be responsible for retarding the atherosclerosis. Rabbits fed high cholesterol diet developed atherosclerosis but it was prevented by kimchi ingredients. The lipid deposition in the aorta arch of rabbit fed Chinese cabbage (7), red pepper powder (41), or garlic (42) was decreased informing that these ingredients might have anti-atherogenic property. The active principle, 3-(4'-hydroxyl-3',5'-dimethoxyphenyl)propionic acid (HDMPPA) in Chinese cabbage kimchi responsible for lipid lowering activity showed anti-atherogenic effect also. HDMPPA showed both preventive and therapeutic effect on hypercholesterolemic rabbits and these beneficial effects were comparable to that of Simvastatin which is widely used drug for treating the hypercholesterolemia in the clinic. The plasma cholesterol and LDL cholesterol was decreased while HDL cholesterol was increased due to the decreased activity of HMG-CoA reductase, CETP, and ACAT (40) and the thickness of aorta arch of rabbit fed cholesterol was significantly reduced (8). The mechanisms of 3-(4'-hydroxyl-3',5'-dimethoxyphenyl)propionic acid preventing atherosclerosis was not fully understood however HDMPPA seems to inhibit form cell formation in the smooth muscle cell of aorta by prohibiting the migration of macrophage to the aorta and to inhibit inflammatory response in aorta by increasing NO synthesis and inhibiting COX-2 expression (8). HDMPPA increased GSH level and suppressed NO production in Murine macrophage cell (43) Besides this animal study, anti-atherogenic effects of kimchi was observed in the clinical studies. Nonlipidemic Caucasians (4 peoples) consumed 150 g kimchi (1/2 cup) daily with regular meal for 2 weeks showed that lag phase duration of LDL oxidation was extended two times compared to that of baseline (0 week) suggesting that kimchi might have ability to reduce the incidence of arteriosclerosis. And kimchi pill supplementation (3 g daily) study with 12 middle aged healthy Korean adults showed that atherogenic index for kimchi group was decreased compared to that for placebo group (39).

## REFERENCES

1. Lee YM, Kwon MJ, Kim JK, Suh HS, Choi JS, Song YO. 2004. Isolation and identification of active principle in Chinese cabbage kimchi responsible for antioxidant activity. *Korean J Food Sci Technol* 36: 129-133.
2. Hwang JH, Song YO, Cheigh HS. 2000. Fermentation characteristics and antioxidative effect of red mustard leaf kimchi. *J Korean Soc Food Sci Nutr* 29: 1009-1015.
3. Kwon MJ, Song YS, Song YO. 1998. Antioxidative effect of kimchi ingredients on rabbits fed cholesterol diet. *J Korean Soc Food Sci Nutr* 27: 1189-1196.
4. Ryu BM, Ryu SH, Lee YS, Jeon YS, Moon GS. 2004. Effect of different kimchi diets on oxidation and photooxidation in liver and skin of hairless mice. *J Korean Soc Food Sci Nutr* 33: 291-298.
5. Ryu SH, Jeon YS, Kwon MJ, Moon JW, Lee YS, Moon GS. 1997. Effect of kimchi extracts to reactive oxygen species in skin cell cytotoxicity. *J Korean Soc Food Sci Nutr* 26: 814-821.
6. Ryu BM, Ryu SH, Jeon YS, Lee YS, Moon GS. 2004. Inhibitory effect of solvent fraction of various kinds of kimchi on ultraviolet B induced oxidation and erythema formation of hairless mice skin. *J Korean Soc Food Sci Nutr* 33: 785-790.
7. Kwon MJ. 1998. Antiatherogenic effect of Baechu kimchi. *PhD Dissertation*. Pusan National University.
8. Kim HJ. 2004. The preventive and therapeutic effects of 3-(4'-hydroxyl-3',5'-dimethoxyphenyl)propionic acid, an active principle in kimchi, on atherosclerosis in rabbits. *PhD Dissertation*. Pusan National University.
9. Kim HJ, Kwon MJ, Song YO. 2000. Effects of solvent fraction of Korean cabbage kimchi on antioxidative enzyme activities and fatty acid composition of phospholipids of rabbit fed 1% cholesterol diet. *J Korean Soc Food Sci Nutr* 29: 900-907.
10. Kim JH, Ryu JD, Song YO. 2002. The effect of kimchi intake on free radical production and the inhibition of oxidation in young adults and the elderly people. *Korean J Community Nutr* 7: 257-265.
11. Kim JH, Ryu JD, Lee HG, Park JH, Moon GS, Cheigh HS, Song YO. 2002. The effect of kimchi on production of free radicals and anti-oxidative enzyme activities in the brain of SAM. *J Korean Soc Food Sci Nutr* 31: 117-123.
12. Park KY. 1995. The nutritional evaluation and antimutagenic and anticancer effects of kimchi. *J Korean Soc Food Nutr* 24: 169-182.
13. Park KY, Cheigh HS. 2000. Antimutagenic and anticancer effects of lactic acid bacteria isolated from kimchi. *Bioindustry News* 13: 84-90.
14. Rhee YH, Kang MS. 1996. Physico-chemical characteristics and beta-galactosidase activity of *Lactobacillus plantarum* from kimchi. *Agric Chem Biotechnol* 39: 54-59.
15. Shin K, Chae O, Park I, Hong S, Choe T. 1998. Antitumor effects of mice fed with cell lysates of *Lactobacillus plantarum* isolated from kimchi. *Korean J Biotechnol Bioeng* 13: 357-363.
16. Heerdt BG, Houston MA, Augenlicht LH. 1994. Potentiation by specific short-chain fatty acids of differentiation and apoptosis in human colonic carcinoma cell lines. *Cancer Res* 54: 3288-3294.
17. Marchetti MC, Migliorati G, Moraca G. 1997. Possible mechanisms involved in apoptosis of colon tumor cell lines induced by deoxycholic acid, short-chain fatty acids, and their mixtures. *Nutr Cancer* 28: 74-80.
18. Cho EJ, Rhee SH, Lee SM, Park KY. 1997. In vitro antimutagenic and anticancer effects of kimchi fractions. *J Korean Assoc Cancer Prevention* 2: 113-121.
19. Choi MW, Kim KH, Park KY. 1997. Effects of kimchi extracts on the growth of Sarcoma-180 cells and phagocytic activity of mice. *J Korean Soc Food Sci Nutr* 26: 254-260.
20. Choi JW, Park JH, Ji ST, Choi OB, Shin HK. 1999. Antigenotoxic effect of dominant bacterial isolated from kimchi in vitro. *Korean J Food Sci Technol* 31: 1071-1076.
21. Cho EJ, Rhee SH, Kang KS, Park KY. 1999. In vitro anticancer effect of Chinese cabbage kimchi fractions. *J Korean Soc Food Sci Nutr* 28: 1326-1331.
22. Al-Delaimy KS, Ali SH. 1970. Antibacterial action of vegetable extracts on the growth of pathogenic bacteria. *J*

*Sci Fd Agric* 21: 110-112.

23. Sheo HJ, Seo YS. 2003. The antibacterial action of Chinese cabbage kimchi juice on *Staphylococcus aureus*, *Salmonella enteritidis* and *Enterobacter cloacae*. *J Korean Soc Food Sci Nutr* 32: 1351-1356.
24. Chae O, Shin K, Chung H, Choe T. 1998. Immunostimulation effects of mice fed with cell lysate of *Lactobacillus plantarum* isolated from kimchi. *Korean J Biotechnol Bioeng* 13: 424-430.
25. Kim MJ, Kwon MJ, Song YO, Lee EK, Youn HJ, Song YS. 1997. The effects of kimchi on hematological and immunological parameters in vivo and in vitro. *J Korean Soc Food Sci Nutr* 26: 1-7.
26. Kim JY, Lee YS. 1997. The effects of kimchi intake on lipid contents of body and mitogen response of spleen lymphocytes in rats. *J Korean Soc Food Sci Nutr* 26: 1200-1207.
27. Sheo HJ, Seo YS. 2004. The effects of dietary Chinese cabbage kimchi juice on the lipid metabolism and body weight gain in rats fed high-calories-diet. *J Korean Soc Food Sci Nutr* 33: 91-100.
28. Do MS, Hong SE, Ha JH, Choi SM, Ahn IS, Yoon JY, Park KY. 2004. Increased lipolytic activity by high-pungency red pepper extract in rat adipocytes in vitro. *J Food Sci Nutr* 9: 34-38.
29. Baek YH, Kwak JR, Kim SJ, Han SS, Song YO. 2001. Effects of kimchi supplementation and/or exercise training on body composition and plasma lipids in obese middle school girls. *J Korean Soc Food Sci Nutr* 30: 906-912.
30. Srinivasan K, Sambaiah K. 1980. Hypocholesterolemic effect of red pepper and capsaicin. *Indian J Exp Biol* 898-899.
31. Yoshioka M, St-Pierre S, Suzuki M, Tremblay A. 1998. Effects of red pepper added to high-fat and high-carbohydrate meals on energy metabolism and substrate utilization in Japanese women. *Br J Nutr* 80: 503-510.
32. Gilliland SE, Nelson CR, Maxwell C. 1985. Assimilation of cholesterol by *Lactobacillus acidophilus*. *Appl Environ Microbiol* 49: 377-381.
33. Gilliland SE, Speck ML. 1977. Deconjugation of bile acids by intestinal lactobacilli. *Appl Environ Microbiol* 33: 15-18.
34. Kwon MJ, Chun JH, Song YS, Song YO. 1999. Daily kimchi consumption and its hypolipidemic effect in middle-aged men. *J Korean Soc Food Nutr* 28: 1144-1150.
35. Kim JH. 2001. The effect of kimchi intake on antiaging characteristics in SAM and human. *PhD Dissertation*. Pusan National University.
36. Kwon MJ, Song YO, Song YS. 1997. Effects of kimchi on tissue and fecal lipid composition and apolipoprotein and thyroxine levels in rats. *J Korean Soc Food Nutr* 26: 507-513.
37. Hwang JW, Song YO. 2000. The effects of solvent fractions of kimchi on plasma lipid concentration of rabbit fed high cholesterol diet. *J Korean Soc Food Nutr* 29: 204-210.
38. Jeon HN, Kwon MJ, Song YO. 2002. Effects of kimchi solvent fractions on accumulation of lipids in heart, kidney, and lung of rabbit fed high cholesterol diet. *J Korean Soc Food Nutr* 31: 814-818.
39. Choi SH, Kim HJ, Kwon MJ, Baek YH, Song YO. 2001. The effect of kimchi pill supplementation on plasma lipid concentration in healthy people. *J Korean Soc Food Sci Nutr* 30: 913-920.
40. Kim HJ, Kwon MJ, Seo JM, Kim JK, Song SH, Suh HS, Song YO. 2004. The effect of 3-(4'-hydroxyl-3',5'-dimethoxyphenyl)propionic acid in Chinese cabbage kimchi on lowering hypercholesterolemia. *J Korean Soc Food Sci Nutr* 33: 52-58.
41. Kwon MJ, Song YS, Choi MS, Song YO. 2003. Red pepper attenuates cholesteryl ester transfer protein activity and atherosclerosis in cholesterol-fed rabbits. *Clinica Chimica Acta* 332: 37-44.
42. Kwon MJ, Song YS, Choi MS, Park SJ, Jeong KS, Song YO. 2003. Cholesteryl ester transfer protein activity and atherogenic parameters in rabbits supplemented with cholesterol and garlic powder. *Life Sciences* 72: 2953-2964.
43. Song YS, Choi CY, Suh H, Song YO. 2004. 3-(4'-hydroxyl-3',5'-dimethoxyphenyl)propionic acid suppresses NO production and elevates GSH levels in murine macrophage. *J Food Sci Nutr* 9: 270-275.