

## Poster PE-1

### Proton MR Spectroscopy of the Primary Motor Cortex and Supplementary Motor Area in Parkinson's Disease

백현만<sup>1</sup>, 최보영<sup>1</sup>, 김정석<sup>1</sup>, 윤성익<sup>1</sup>, 이정우<sup>1</sup>, 전신수<sup>2</sup>  
가톨릭의대 <sup>1</sup>의공학교실, <sup>2</sup>신경외과학교실

**목적** : To investigate whether there are significant changes in regional brain metabolism in patients with Parkinson's disease after thalamotomy using proton magnetic resonance spectroscopy (1H MRS).

**대상 및 방법** : Fifteen patients with Parkinson's disease of mean age 56.5 years (7 males and 8 females; mean age, 56.5 years) and mean disease duration 7.7 years (range 2-15 years) that have treated with levodopa were included. All patients with tremor experienced amelioration of their symptoms on the sidecontralateral to the thalamotomy. Complete abolition of tremor occurred in 87% (13/15) patients with significant improvement of tremor in additional two patients. In vivo 1H MRS study was performed on a 1.5 T MRI system (GE Signa Advantage, version 4.8) using STEAM sequence after water suppression with CHESS RF pulse and dephasing gradients. As a single-voxel technique, 1H MR spectra were obtained from the volume of interested regions in M1 and SMA of Parkinson's patients. Spectral parameters were: 20 ms TE, 2000 ms TR, 128 averages, 2500 Hz spectral width, and 2048 data points. Raw data were processed by the SAGE data analysis package. Peak areas of NAA, Cr, Cho, Ins, and sum (Glx) of  $\gamma$ -Amino butyrate (GABA) and glutamate were calculated by means of fitting the spectrum to a summation of Lorentzian curves using a Marquardt algorithm. After blindly processed, we calculated the metabolite ratios of NAA/Cho, NAA/Cr, Cho/Cr, Ins/Cr, and Glx/Cr. Ratios are given as the mean  $\pm$  SD. Statistical significance was determined using Student's paired t-tests between control subjects and patients with Parkinson's disease, where  $p < 0.05$  was considered significant.

**결과** : Compared with age-matched controls, substantial difference of NAA/Cr (or, NAA/Cho) ratios from the voxels of region of interests in Parkinson's disease patients was not observed [ $1.41 \pm 0.22$  (or,  $1.61 \pm 0.29$ ),  $1.43 \pm 0.28$  (or,  $1.63 \pm 0.33$ )]. In all patients with clinical improvement following thalamotomy, NAA/Cr (or, NAA/Cho) ratios were not statistically established ( $p > 0.05$ ) from the selected voxels in M1 and SMA. Thus, NAA/Cr (or, NAA/Cho) ratios showed generally normal levels in M1 and SMA in Parkinson's disease patients with clinical outcome following thalamotomy.

**결론** : The results of this study indicated no postoperative metabolic changes of NAA/Cr (or, NAA/Cho) ratios in M1 and SMA of the hemisphere that did undergo surgery. Thus, the metabolic changes were not established in Parkinson's disease patients with clinical improvement. These changes didn't be consistent in both thalamus and motor cortex contralateral to the most affected side. Therefore, the surgical changes induced by

thalamotomy may not directly affect the specific regional brain networks associated with clinical manifestations of parkinsonism. In conclusion, the present study suggests 1H MRS may be a useful modality for the aid in better understanding the pathophysiologic process in patients. Indeed, an understanding of the functional basis for surgical response may have great importance in the design of new treatment strategies for Parkinson's disease and related disorders. It is necessary to investigate the MR spectral alterations after a long-term follow-up examination in patients with Parkinson's disease for further detail analysis.

**Acknowledgment:** This study was supported by a grant of the Center for Functional and Metabolic Imaging Technology, Ministry of Health & Welfare, Republic of Korea (02-PJ3-PG6-EV07-0002).