Expression of functional *Carassius auratus*originated novel DNA-binding histone H1 in *Escherichia coli*

Quande Wei^{1,3}, Kyoung Ro Kim^{1,2}, Xinbing Yu³, and Hyung Joon Cha^{1,2}*

¹ Division of Molecular and Life Sciences & ² Department of Chemical Engineering,
Pohang University of Science and Technology, Pohang 790-784, Korea

³ Department of Pathogenic Biology, Sun Yat-sen University, Guangzhou 510089, China

* Corresponding author. Fax: +82-54-279-5528.

E-mail address: hjcha@postech.ac.kr (H.J. Cha).

Histones are DNA-binding proteins having function to assist in DNA packaging and protection. In the present work, we cloned novel histone H1 cDNA (638 bp) from goldfish Carassius auratus and this histone H1 amino acid sequence was shared 68.1% identity and 73.9% positives with that from rainbow trout Salmo gairdneri. We successfully expressed whole length recombinant histone H1 (~20 kDa) using maltose binding protein (MBP) fusion strategy in Escherichia coli expression system. We found that this recombinant histone H1 protein had functionality on DNA binding and protection. In addition, the C-terminal domain (~ 6 kDa) of histone H1 with 61 amino acids also maintained same abilities. Based on these results, this novel histone H1 protein, especially its derived C-terminal peptide, could be used as a potential mediator for efficacious gene delivery system.

References

- 1. Ai Leen Lam, Dorothy E. Pazin, Beth A. Sullivan. (2005). Control of gene expression and assembly of chromosomal subdomains by chromatin regulators with antagonistic functions. Chromosoma, Online First: 1-10.
- Arents, G., E.N. Moudriariakis. (1995). The histone-fold: a ubiquitous architectural motif utilized in DNA compaction and protein dimerization. Proc. Natl. Acad. Sci. USA, 92:11170-1117.
- 3. Bharath, M.M., Ramesh, S., Chandra, N.R., Rao, M.R. (2002). Identification of a 34 amino acid stretch within the C-terminus of histone H1 as the DNA-condensing domain by site-directed mutagenesis. Biochemistry, 41:76177627.