

금목서(*Osmanthus fragrans* var. *aurantiacus* Makino) 꽃으로부터 이차대사산물의
분리 및 동정

경희대학교: 이도경, 최은희, 정락훈, 이재웅, 송명중, 류하나, 백남인*

Isolation and Identification of secondary metabolites from the Flower of
Osmanthus fragrans var. *aurantiacus* Makino

Graduate School of Biotechnology & Plant Metabolism Research Center, Kyung Hee
university, Yong-In 446-701, Korea

Do-Gyeong Lee, En-Jj Cui, lakoon Jung, Jae-Woong Lee, Myoung-Chong Song,
Ha-Na Lyu, Nam-In Baek*

Objectives

Osmanthus fragrans, the genus *Osmanthus* in the family of Oleaceae, is distributed in China, Japan and southern area of Korea. It is evergreen tree, the flowers are hermaphrodite (have both male and female organs) and come out from June to August. They are not only used in cosmetics for the hair and skin but used by Chinese to give a pleasant aroma to tea and wine. Also it was reported that *O. fragrans* has the biological effect of being anti-tussive. Except them, there is few report about biological and phytochemical research on this plant.

Materials and Methods

The dried and powdered flowers were extracted with 80% aqueous MeOH, and the concentrated extract was partitioned with EtOAc, *n*-BuOH and H₂O, successively. From the EtOAc fraction, three lignans were isolated through the repeated silica gel, ODS and Sephadex LH-20 column chromatographies.

Results

According to the results of physico-chemical data including NMR, MS and IR, the chemical structures of the compounds were determined such as (-)-lariciresinol and so on. These lignan compounds have been first isolated from *O. fragrans* in this study.

Corresponding author : Nam-In Baek E-mail: nibaek@khu.ac.kr, Tel: 031-201-2661