

accretion rate, flat spectrum radio quasars and BL Lac objects, having different scaling relations with $\alpha \approx 1$ and ≈ 2 , respectively. We find that modelling the periodograms of four of our sources requires the assumption of broken powerlaw spectra. From simulating lightcurves as superpositions of exponential flares we conclude that strong overlap of flares leads to featureless simple power-law periodograms of AGNs at radio wavelengths in most cases (The paper is about to be submitted to ApJ).

[7 GC-16] Acceleration of Relativistic Jets on Sub-parsec Scales

Sang-Sung Lee^{1,2}, Andrei Lobanov³, Thomas P. Krichbaum³, J. Anton Zensus³

¹*Korea Astronomy and Space Science Institute,*

²*Korea University of Science and Technology,*

³*Max-Planck Institute for Radio Astronomy*

Jets of compact radio sources are highly relativistic and Doppler boosted, making studies of their intrinsic properties difficult. Observed brightness temperatures can be used to study the intrinsic physical properties of the relativistic jets. The intrinsic properties of relativistic jets depend on inner jet models. We aimed to observationally test the inner jet models. The very long baseline interferometry (VLBI) cores of compact radio sources are optically thick at a given frequency. The distance of the core from the central engine is inversely proportional to the frequency. Under the equipartition condition between the magnetic field energy and particle energy densities, the absolute distance of the VLBI core can be predicted. We compiled the brightness temperatures of VLBI cores at various radio frequencies of 2, 8, 15, and 86-GHz. The brightness temperatures in the rest frame were investigated in the sub-parsec regions of the compact radio sources. From the vicinity of the central engine, the brightness temperatures increased slowly and then rose with steeper slope, indicating that the Lorentz factor increases along the jet. This implies that the jets are accelerated in the (sub-)parsec regions from the central engine.

[7 GC-17] The drivers and energetics of ionized gas outflows in powerful Type 2 AGN in the local Universe

Marios Karouzos¹, Jong-Hak Woo¹, Hyun-Jin Bae^{1,2}

¹*Department of Physics and Astronomy, Seoul National University,* ²*Department of Astronomy, Yonsei University*

There exist scaling relations that link the mass of supermassive black holes with both the velocity dispersion and the mass of the central stellar cusp of their host galaxies. This implies that galaxies co-evolve with their central black holes, potentially through the feedback from actively accreting supermassive black holes (AGN). We use integral field spectroscopy data from the 8.2m Gemini-North telescope to investigate ionized gas outflows in luminous local ($z < 0.1$) Type 2 AGN. Our sample of 6 galaxies was selected based on their [OIII] dust-corrected luminosity ($> 10^{42}$ erg/s) and signatures of outflows in the [OIII] line profile of their SDSS spectra. These are arguably the best candidates to explore AGN feedback in action since they are $< 1\%$ of a large local type 2 AGN SDSS sample selected based on their [OIII] kinematics. Expanding on previously reported results concerning the kinematic decomposition and size determination of these outflows, here we report their photoionization properties and energetics. We find strong evidence that connect the extreme kinematics of the ionized gas with AGN photoionization. The kinematic component related to the AGN-driven outflow is clearly separated from other kinematic components, such as gravitation- or stellar-driven motions, on the velocity and velocity dispersion diagram. Our spatially resolved kinematic analysis reveals that up to 90% of the mass and kinetic energy of the outflow is contained within the central kiloparsec of the galaxy. The total mass and kinetic energy of the outflow correlate well with the AGN bolometric luminosity, resulting in energy conversion efficiencies between 0.01% and 1%. Intriguingly, we detect ubiquitous signs of ongoing circumnuclear star formation. Their small size, the centrally contained mass and energy, and the universally detected circumnuclear star formation cast doubts on the potency of these AGN-driven outflows as agents of negative feedback.

[7 GC-18] Discovery of a Faint Quasar at $z \sim 6$ and Implications for Cosmic Reionization

Yongjung Kim^{1,2}, Myungshin Im^{1,2}, Yiseul Jeon^{1,2}, Minjin Kim^{3,4}, Changsu Choi^{1,2}, Jueun Hong^{1,2}, Minhee Hyun^{1,2}, Hyunsung David Jun^{1,5}, Marios Karouzos², Dohyeong Kim^{1,2}, Duho Kim^{1,6}, Jae-Woo Kim^{1,2}, Ji Hoon Kim⁷, Seong-Kook Lee^{1,2}, Soojong Pak⁸, Won-Kee Park³, Yoon Chan Taak^{1,2}, and Yongmin Yoon^{1,2}

¹*Center for the Exploration of the Origin of the Universe (CEO),* ²*Astronomy Program, FPRD,*

Department of Physics & Astronomy, Seoul National University, 3Korea Astronomy and Space Science Institute, 4University of Science and Technology, 5Jet Propulsion Laboratory, California Institute of Technology, 6Arizona State University, School of Earth and Space Exploration, 7Subaru Telescope, National Astronomical Observatory of Japan, 8School of Space Research and Institute of Natural Sciences, Kyung Hee University...

Recent studies suggest that faint active galactic nuclei may be responsible for the reionization of the universe. Confirmation of this scenario requires spectroscopic identification of faint quasars ($M_{1450} > -24$ mag) at $z > 6$, but only a very small number of such quasars have been spectroscopically identified so far. Here, we report the discovery of a faint quasar IMS J220417.92+011144.8 at $z \sim 6$ in a 12.5 deg^2 region of the SA22 field of the Infrared Medium-deep Survey (IMS). The spectrum of the quasar shows a sharp break at $\sim 8443 \text{ \AA}$, with emission lines redshifted to $z = 5.944 \pm 0.002$ and rest-frame ultraviolet continuum magnitude $M_{1450} = -23.59 \pm 0.10$ AB mag. The discovery of IMS J220417.92+011144.8 is consistent with the expected number of quasars at $z \sim 6$ estimated from quasar luminosity functions based on previous observations of spectroscopically identified low-luminosity quasars. This suggests that the number of $M_{1450} \sim -23$ mag quasars at $z \sim 6$ may not be high enough to fully account for the reionization of the universe. In addition, our study demonstrates that faint quasars in the early universe can be identified effectively with a moderately wide and deep near-infrared survey such as the IMS.

[7 GC-19] GRB 140304A at $z=5.283$: Implications on the high redshift universe and the observed flaring activities

Soomin Jeong
Institute for Science and Technology in Space, SKKU, Korea and Instituto de Astrofísica de Astronomía-CSIC, Granada, Spain

Gamma ray burst, the most brightest explosion phenomena in the current universe is well suited for study of high redshift universe. We report the afterglow multi-wavelength observation and GTC spectroscopy follow up of GRB 140304A which was exploded at $z=5.283$. The spectrum was shown damped Lyman alpha features and a series of absorption lines S, Si, SiII*, Oi, CII, CII*, SiIV are

clearly detected at common redshift. Clear optical flares are detected when X-ray flare happened and a possible gamma-ray excess also. At this conference, we report on implications for the GRB host and environments using its absorption features which place the results in context to other well studied high redshift GRBs and studies about the ejecta using its observed flaring activities.

[7 GC-20] Study on mapping of dark matter clustering from real space to redshift space

Yi Zheng, Yong-Seon Song
Korea Astronomy and Space Science Institute, Daejeon 34055, Republic of Korea

The mapping of dark matter clustering from real to redshift spaces introduces the anisotropic property to the measured density power spectrum in redshift space, known as the Redshift Space Distortion (hereafter RSD) effect. The mapping formula is intrinsically non-linear, which is complicated by the higher order polynomials due to the indefinite cross correlations between the density and velocity fields, and the Finger-of-God (hereafter FoG) effect due to the randomness of the peculiar velocity field. Furthermore, the rigorous test of this mapping formula is contaminated by the unknown non-linearity of the density and velocity fields, including their auto- and cross-correlations, for calculating which our theoretical calculation breaks down beyond some scales. Whilst the full higher order polynomials remains unknown, the other systematics can be controlled consistently within the same order truncation in the expansion of the mapping formula, as shown in this paper. The systematic due to the unknown non-linear density and velocity fields is removed by separately measuring all terms in the expansion using simulations. The uncertainty caused by the velocity randomness is controlled by splitting the FoG term into two pieces, 1) the non-local FoG term being independent of the separation vector between two different points, and 2) the local FoG term appearing as an indefinite polynomials which is expanded in the same order as all other perturbative polynomials. Using 100 realizations of simulations, we find that the best fitted non-local FoG function is Gaussian, with only one scale-independent free parameter, and that our new mapping formulation accurately reproduces the observed power spectrum in redshift space at the smallest scales by far, up to $k \sim 0.3 \text{ h/Mpc}$, considering the resolution of future experiments.