

around or within Virgo. Our results strongly suggest that substructures can play important roles in galaxy evolution while galaxies are falling to the cluster.

[포 GC-19] A pilot study of dense molecular gas in a Virgo spiral using a KVN single-dish

Bumhyun Lee, Junhyun Baek, Aeree Chung
Department of Astronomy, Yonsei University

NGC 4402 is a spiral galaxy located in the Virgo cluster. It is undergoing active HI gas stripping due to the strong ICM pressure, showing evidence for recent quenching of star formation. Its CO disk is also highly disturbed as HI, yet unlike HI disk, no sign of significant molecular gas stripping is found. Aiming to better understand how atomic gas stripping and disturbed molecular gas result in star formation quenching, we probe properties of molecular gas in the densest forms. As a pilot study, we observed HCN (1-0) and HCO+ (1-0) in the center of NGC 4402 using one of the Korean VLBI Network (KVN) antennas located at Yonsei site. In this work, we present the result from the KVN single-dish observations and discuss its implications.

[포 GC-20] The Contribution of Mergers on Star Formation in Nearby UV-Bright Galaxies (별탄생 은하의 별 생성에 대한 병합 작용의 기여도 연구)

Gu Lim(임구), Myungshin Im(임명신), Changsu Choi(최창수), Yongmin Yoon(윤용민)
Center for Exploration of the Origin of the Universe (CEO), Astronomy Program, Dept. of Physics & Astronomy, Seoul National University (서울대학교)

Star formation in galaxies is one of the key factors in galaxy evolution. It is believed that star formation is triggered and enhanced by mergers among galaxies or secular evolution. However, how much these two mechanisms contribute on star formation is not well known yet. Recently, many other studies show observational evidences of faint merger features(tidal tails, stellar streams) around nearby galaxies with deep optical imaging. This study aims to investigate the fraction of star forming galaxies exhibiting faint features to total galaxies. We are analyzing samples of 76 star forming galaxies ($NUV < -18$) to find merger features from stacked B, R band frames taken at

Maidanak 1.5m, McDonald 2.1m telescope and g, r frames from Canada-France-Hawaii Telescope (CFHT) MegaCam archival data. With the fraction, we can expect to know the contribution of mergers on star formation to galaxies.

[포 GC-21] Polarization of Rayleigh Scattered Ly α in Active Galactic Nuclei

Seok-Jun Chang¹, Hee-Won Lee¹, and Yujin Yang²
¹*Department of Physics and Astronomy, Sejong University*
²*Korea Astronomy and Space Science Institute*

Active galactic nuclei (AGNs) typically show a non-thermal continuum locally represented by a power-law and many prominent emission lines in the UV and optical regions. AGNs are classified by two types, where Type 1 AGNs exhibit both broad and narrow lines and only narrow lines are observed in Type 2 AGNs. The unification models of AGNs invoke the existence of a molecular torus just outside of the broad line region. In the presence of a high column HI region associated with the molecular torus, we propose that significant fraction of broad line photons near Lyman series can be scattered by atomic hydrogen in the torus. In particular, Ly α being the strongest emission line, strong linear polarization may develop around Ly α through Rayleigh scattering. We adopt a Monte Carlo technique to investigate the polarized transfer of Ly α in a thick HI region with the shape of a torus. We consider the range of HI column density $N_{\text{HI}} = 10^{20-23}$ with fixed geometric parameters of the torus such as the inner and outer radii and the height. We present the polarized spectra and angular distribution of Rayleigh scattered radiation around Ly α . We find that the Ly α core part is polarized in the direction perpendicular to the symmetry axis whereas in the far wing part it is polarized in the parallel direction. It is concluded that the unification of AGNs implies that Ly α can be uniquely polarized through Rayleigh scattering.

[포 GC-22] SNU AGN Monitoring Project (SAMP) using reverberation mapping of luminous AGNs

Yiseul Jeon¹, Jong-Hak Woo¹, and SAMP team^{1,2,3,4,5,6}
¹*Seoul National University*, ²*Korea Astronomy and Space Science Institute*, ³*University of Michigan*, ⁴*University of California, Irvine*, ⁵*University of California, Los Angeles*, ⁶*California Polytechnic State University*

The links between super-massive black hole masses and their host galaxy properties are observed, indicating that black hole growth and host galaxy evolution are closely related. Reverberation mapping, which uses the time delay from the central black hole to broad line regions, is one of the best methods to estimate masses of black holes of active galactic nuclei (AGNs). However, only masses of about 50 black holes have been determined in reverberation mapping studies so far, and most of them are limited to optical luminosities below 10^{45} erg/s due to the challenges of long-term time domain observations in both photometry and spectroscopy. In this project, we expand reverberation mapping samples to higher luminosities of $> 10^{44.5}$ erg/s at $0.1 < z < 0.35$, that have expected time lags of 40 - 250 light days. Photometric (using LOAO 1-m and MDM 1.3-m) and spectroscopic (using MDM 2.4-m and Lick 3-m) monitoring campaigns are being conducted for a 3 year duration and 20 day cadence. Precedent photometric observations in 2015B show some targets with variability and follow-up spectroscopic observations are on-going. In this presentation, we introduce our project, present reverberation mapping simulation results, and preliminary results on photometry. These reverberation mapping masses of relatively high luminous AGNs will provide a strong constraint on black hole mass calibration, e.g., the single-epoch mass estimation.

[포 GC-23] Photometric Reverberation Mapping with SQUEAN: A Test Study using Medium Bands

Yoon Chan Taak^{1,2}, Myungshin Im^{1,2}, and Changsu Choi^{1,2}

¹Center for the Exploration of the Origin of the Universe

²Astronomy Program, Department of Physics and Astronomy, Seoul National University

Photometric reverberation mapping is an effective alternative to time consuming spectroscopy. It usually employs narrow bands to track the luminosity variations of broad emission lines, such as Balmer lines, and broadbands for the continuum variability. Here, we investigate the feasibility of substituting these for medium bands, with 50nm widths, that are currently being used on the SED Camera for QUasars in EARly uNiverse (SQUEAN) installed on the 2.1m Otto Struve Telescope at McDonald Observatory. Three targets

with recent variability and/or short expected time lags were selected, and observed for 15 minutes each in the medium band containing H α , and the two adjacent bands for continuum subtraction afterwards. Analysis shows that for one of the objects, SDSS J0350+0037, the pure H α emission line flux has a S/N ~ 12 , so that variabilities up to $\sim 8\%$ are detectable. Thus, future observations using these medium bands on SQUEAN seem to be practical.

[포 GC-24] Weak-Lensing Study of Galaxy Cluster PLCKG287.0+32.9

Kyle Finner, Myungkook James Jee
Yonsei University, Department of Astronomy

Merging galaxy clusters, such as PLCKG287.0+32.9, provide a window into the formation process of the large scale structure of the universe. PLCKG287.0+32.9 is an enormous merging galaxy cluster with mass estimated to be $\sim 10^{15}$ Msun. It hosts a pair of mega-parsec sized radio relics with projected offsets from the X-ray center of approximately 350kpc and 2.7Mpc, suggesting a NW-SE merging scenario with relics originating from two separate passes (Bonafede et al. 2014). A detected radio halo coincides with the center of x-ray emission. We present the motivation for our weak lensing study of the merging galaxy cluster PLCKG287.0+32.9 using recent Subaru optical imaging. We discuss the basics of weak-lensing and the criteria for source selection. In addition, we describe our method of PSF modeling and mass reconstruction.

[포 GC-25] Parametric modelling of the velocity dispersion profiles of 11 elliptical galaxies : dark matter versus MOND

In-Taek Gong and Kyu-Hyun Chae
Department of Physics and Astronomy, Sejong University

운동학적으로 측정된 질량과 측광으로 측정된 질량이 불일치하는 질량 불일치 문제는 현대천문학의 중요한 문제이다. 현재 이러한 질량 불일치에 대한 두 가지 해결책이 제시되었다. 하나는 현대 표준우주론인 Λ CDM 패러다임의 핵심 요소인 암흑물질, 다른 하나는 Milgrom에 의해 제시된 수정된 뉴턴역학(Modified Newtonian dynamics: MOND)이다. 두 방법에 대한 많은 연구가 진행되었는데, 최근 연구 결과에 의하면 나선형 은하의 회전 속도 윤곽은 MOND와 잘 부합하나 타원은하에 대해서는 불명확하다. 여기서 우리는 ATLAS3D project 에서 얻어진 260개의 조기형 은하 중 거의 원형인 11개의 타원은하