

외부은하 / 은하단

[구 GC-01] The Limited Impact of AGN Outflows: IFU study of 20 local AGNs

Hyun-Jin Bae^{1,2}, Jong-Hak Woo², Marios Karouzos², Elena Gallo⁴, Helene Flohic⁵, Yue Shen⁶, and Suk-Jin Yoon¹

¹*Department of Astronomy and Center for Galaxy Evolution Research, Yonsei University,* ²*Astronomy Program, Department of Physics and Astronomy, Seoul National University,* ⁴*Department of Astronomy, University of Michigan, Ann Arbor, USA,* ⁵*Department of Physics, University of the Pacific, USA,* ⁶*Department of Astronomy, University of Illinois at Urbana-Champaign, USA*

To investigate AGN outflows as a tracer of AGN feedback on the host galaxies, we perform integral-field spectroscopy of 20 type 2 AGNs at $z < 0.1$ using the Magellan/IMACS and the VLT/VIMOS. The observed objects are luminous AGNs with the [O III] luminosity $> 10^{41.5}$ erg/s, and exhibit strong outflow signatures in the [O III] kinematics. We obtain the maps of the narrow and broad components of [O III] and H α lines by decomposing the emission-line profile. The broad components in both [O III] and H α represent the non-gravitational kinematics, (i.e., gas outflows), while the narrow components represent the gravitational kinematics (i.e., rotational disks), especially in H α . By using the spatially integrated spectra within the flux-weighted size of the narrow-line region, we estimate the outflow energetics. The ionized gas mass is $(1.0-38.5) \times 10^5 M_{\odot}$, and the mean mass outflow rate is $4.6 \pm 4.3 M_{\odot}/\text{yr}$, which is a factor of ~ 260 higher than the mean mass accretion rate $0.02 \pm 0.01 M_{\odot}/\text{yr}$. The mean energy injection rate is $0.8 \pm 0.6\%$ of the AGN bolometric luminosity L_{bol} , while the mean momentum flux is $(5.4 \pm 3.6) \times L_{\text{bol}}/c$, except for two most kinematically energetic AGNs. The estimated energetics are consistent with the expectations for energy-conserving outflows from AGNs, yet we do not find any supporting evidence of instantaneous star-formation quenching due to the outflows.

[구 GC-02] Star Formation of Merging Disk Galaxies with AGN Feedback Effects

Jongwon Park, Rory Smith, and Sukyoung K. Yi.
Department of Astronomy, Yonsei University

Using numerical hydrodynamics code RAMSES,

we perform idealized galaxy merger simulations and study the star formation of merging disk galaxies. In our simulations, we consider the active galactic nucleus (AGN) feedback effect. In order to investigate the star formation influenced by AGN, we run ~ 60 simulations with various initial conditions. We confirm that star formation is more efficiently suppressed in merging galaxies than in isolated galaxies. In the mergers, AGN effect is more significant when the masses of two galaxies are similar. Furthermore, we find that bulge fraction does not affect the star formation when the AGN effect is considered. We discuss the implications on semi-analytic galaxy formation models and the limitation of the current AGN prescriptions.

[구 GC-03] Interferometric Monitoring of Gamma-ray Bright AGNs: S5 0716+714

Sang-Sung Lee^{1,2}, Jee Won Lee^{1,3}, Heffrey A. Hodgson¹, Dae-Won Kim⁴, Juan-Carlos Algaba¹, Sincheol Kang^{1,2}, Jiman Kang¹, Sungsoo S. Kim³
¹*Korea Astronomy and Space Science Institute,* ²*Korea University of Science and Technology,* ³*Kyung Hee University,* ⁴*Seoul National University*

We present the results of very long baseline interferometry (VLBI) observations of gamma-ray bright blazar S5 0716+714 using the Korean VLBI Network (KVN) at the 22, 43, 86, and 129 GHz bands, which are part of the KVN key science program known as the Interferometric Monitoring of Gamma-ray Bright AGNs (iMOGABA). Multi-frequency VLBI observations were conducted in 29 sessions from January 16, 2013 to March 1, 2016. The source was detected and imaged in all available frequency bands. For all observed epochs, the source is compact on the milliarcsecond (mas) scale, yielding a compact VLBI core dominating the synchrotron emission on the mas scale. Based on the multi-wavelength data at 15 and 230 GHz, we found that the source shows multiple prominent enhancements of the flux density at the centimeter (cm) and millimeter (mm) wavelengths, with mm enhancements leading cm enhancements with a time lag of 18 ± 5 days. Turnover frequency is found to vary over our observations between 22 to 69 GHz. Taking into account the synchrotron self-absorption model of the relativistic jet in S5 0716+714, we estimated the magnetic field strength in the mas emission region to be 0.4–66 mG during the observing period, finding that the magnetic field strength is strongly correlated with the turnover frequency and the

relatively strong magnetic field (e.g., $B > 40$ mG) is correlated with flux enhancements at mm wavelengths (e.g., 86 GHz).

[박C-04] A Study of Globular Cluster Systems in the Coma, Fornax, and Virgo Clusters of Galaxies from *HST* ACS and WFC3/IR Imaging

Hyejeon Cho
Center for Galaxy Evolution Research &
Department of Astronomy, Yonsei University

I present new near-infrared (NIR) photometry of globular cluster (GC) systems associated to a cD galaxy NGC 4874 in the core of the Coma cluster and 16 early-type galaxies in the Fornax and Virgo clusters of galaxies using the Infrared Channel of the Wide Field Camera 3 (WFC3/IR) on board the *Hubble Space Telescope (HST)*. Combining these high-resolution NIR data with new *HST* Advanced Camera for Surveys (ACS) optical photometry for NGC 4874 and existing ACS GC catalogs from the ACS Fornax and Virgo Cluster Surveys, I have examined for the first time the GC systems in a statistically significant optical/NIR sample of galaxies spanning a wide range of luminosities and colors. A primary goal of this study is to explore empirically whether the distributions of purely optical and hybrid optical-NIR color indices for extragalactic GCs have different forms and whether the relations between these color indices are nonlinear, indicating that they behave differently with underlying metallicity. I find that some GC systems of large galaxies in our sample show color bimodalities that differ between the optical and optical-NIR colors, in the sense that they have disparate ratios of “blue” and “red” peak GCs, as well as differing ratios in their color dispersions. Consistent with these results, I find empirically that the dependence of hybrid optical-NIR color on purely optical color is nonlinear, with an inflection at intermediate metallicities. These findings underscore the importance of understanding the nature of galaxy-to-galaxy variations in the GC color distributions and color-color relations, as well as the exact forms of the color-metallicity transformations, in interpreting the observational data on GC color bimodality. Our ACS data for NGC 4874 shows that its GC system exhibits a very strong blue tilt, implying a very steep mass-metallicity scaling, and the centroid of this GC system is offset by 4 ± 1 kpc from the luminosity center of NGC 4874, in the direction of NGC 4872. Finally, I discuss the asymmetrical GC distribution

around a dwarf elliptical galaxy in Coma that has a very high relative velocity with respect to the cluster mean at small clustercentric radius.

[구 GC-05] New insights on the chemical evolution in proto-globular clusters and galaxy building blocks (원시 구상성단과 은하 빌딩블록의 새로운 화학적 진화모델)

Young-Wook Lee(이영욱), Jaeyeon Kim(김재연)
Department of Astronomy, Yonsei University

초기 우주의 구상성단과 같은 소규모 은하 빌딩블록이 은하의 형성에 미치는 영향을 이해하기 위하여 우리는 완전히 새로운 개념의 화학적 진화모형을 구축하였다. 최근의 이론적 진보를 바탕으로 우리의 모델에서는 이전 모델과 달리 초신성의 폭발과 분출물이 원시 구상성단 내 잔존 가스에 아무런 영향을 끼치지 못하고 대부분 빠져나간다. 이 경우 화학적 진화는 질량이 큰 별의 윈드와 점근거성계 열성의 분출물에 의해 좌우된다. 놀랍게도 우리의 모델은 오랜 난제인 구상성단 내 Na-Oxygen anticorrelation 및 다중항성종족의 기원을 자연스럽게 설명하면서 동시에 Lee, Joo, & Chung (2015) 이 최근 주장한 것처럼 은하 별지에 헬륨 함량이 매우 높은 항성이 존재할 것으로 예측한다. 이 결과는 은하의 헤일로와 별지 형성에 매우 중요한 단서를 제공한다.

[구 GC-06] Constraints on the Evolution of the Galaxy Stellar Mass Function I: Role of Star Formation, Mergers and Stellar Stripping

Emanuele Contini
Yonsei University.

We study the connection between the observed star formation rate-stellar mass (SFR-M) relation and the evolution of the stellar mass function (SMF) by means of a Subhalo Abundance Matching technique coupled to merger trees extracted from a N-body simulation. Our approach, which considers both galaxy mergers and stellar stripping, is to force the model to match the observed SMF at redshift $z > 2$, and let it evolve down to the present time according to the observed (SFR-M) relation. In this study, we use two different sets of SMFs and two SFR-M relations: a simple power law and a relation with a mass-dependent slope.

Our analysis shows that the evolution of the SMF is more consistent with a SFR-M relation with

[구 GC-07] The mass of the high- z ($z \sim 1.132$) massive galaxy cluster, SPT-CL J2106-5844 using weak-lensing analysis with HST