

relatively strong magnetic field (e.g., $B > 40$ mG) is correlated with flux enhancements at mm wavelengths (e.g., 86 GHz).

[박C-04] A Study of Globular Cluster Systems in the Coma, Fornax, and Virgo Clusters of Galaxies from *HST* ACS and WFC3/IR Imaging

Hyejeon Cho
Center for Galaxy Evolution Research &
Department of Astronomy, Yonsei University

I present new near-infrared (NIR) photometry of globular cluster (GC) systems associated to a cD galaxy NGC 4874 in the core of the Coma cluster and 16 early-type galaxies in the Fornax and Virgo clusters of galaxies using the Infrared Channel of the Wide Field Camera 3 (WFC3/IR) on board the *Hubble Space Telescope (HST)*. Combining these high-resolution NIR data with new *HST* Advanced Camera for Surveys (ACS) optical photometry for NGC 4874 and existing ACS GC catalogs from the ACS Fornax and Virgo Cluster Surveys, I have examined for the first time the GC systems in a statistically significant optical/NIR sample of galaxies spanning a wide range of luminosities and colors. A primary goal of this study is to explore empirically whether the distributions of purely optical and hybrid optical-NIR color indices for extragalactic GCs have different forms and whether the relations between these color indices are nonlinear, indicating that they behave differently with underlying metallicity. I find that some GC systems of large galaxies in our sample show color bimodalities that differ between the optical and optical-NIR colors, in the sense that they have disparate ratios of “blue” and “red” peak GCs, as well as differing ratios in their color dispersions. Consistent with these results, I find empirically that the dependence of hybrid optical-NIR color on purely optical color is nonlinear, with an inflection at intermediate metallicities. These findings underscore the importance of understanding the nature of galaxy-to-galaxy variations in the GC color distributions and color-color relations, as well as the exact forms of the color-metallicity transformations, in interpreting the observational data on GC color bimodality. Our ACS data for NGC 4874 shows that its GC system exhibits a very strong blue tilt, implying a very steep mass-metallicity scaling, and the centroid of this GC system is offset by 4 ± 1 kpc from the luminosity center of NGC 4874, in the direction of NGC 4872. Finally, I discuss the asymmetrical GC distribution

around a dwarf elliptical galaxy in Coma that has a very high relative velocity with respect to the cluster mean at small clustercentric radius.

[구 GC-05] New insights on the chemical evolution in proto-globular clusters and galaxy building blocks (원시 구상성단과 은하 빌딩블록의 새로운 화학적 진화모델)

Young-Wook Lee(이영욱), Jaeyeon Kim(김재연)
Department of Astronomy, Yonsei University

초기 우주의 구상성단과 같은 소규모 은하 빌딩블록이 은하의 형성에 미치는 영향을 이해하기 위하여 우리는 완전히 새로운 개념의 화학적 진화모형을 구축하였다. 최근의 이론적 진보를 바탕으로 우리의 모델에서는 이전 모델과 달리 초신성의 폭발과 분출물이 원시 구상성단 내 잔존 가스에 아무런 영향을 끼치지 못하고 대부분 빠져나간다. 이 경우 화학적 진화는 질량이 큰 별의 윈드와 점근거성계 열성의 분출물에 의해 좌우된다. 놀랍게도 우리의 모델은 오랜 난제인 구상성단 내 Na-Oxygen anticorrelation 및 다중항성종족의 기원을 자연스럽게 설명하면서 동시에 Lee, Joo, & Chung (2015) 이 최근 주장한 것처럼 은하 별지에 헬륨 함량이 매우 높은 항성이 존재할 것으로 예측한다. 이 결과는 은하의 헤일로와 별지 형성에 매우 중요한 단서를 제공한다.

[구 GC-06] Constraints on the Evolution of the Galaxy Stellar Mass Function I: Role of Star Formation, Mergers and Stellar Stripping

Emanuele Contini
Yonsei University.

We study the connection between the observed star formation rate-stellar mass (SFR-M) relation and the evolution of the stellar mass function (SMF) by means of a Subhalo Abundance Matching technique coupled to merger trees extracted from a N-body simulation. Our approach, which considers both galaxy mergers and stellar stripping, is to force the model to match the observed SMF at redshift $z > 2$, and let it evolve down to the present time according to the observed (SFR-M) relation. In this study, we use two different sets of SMFs and two SFR-M relations: a simple power law and a relation with a mass-dependent slope.

Our analysis shows that the evolution of the SMF is more consistent with a SFR-M relation with

[구 GC-07] The mass of the high- z ($z \sim 1.132$) massive galaxy cluster, SPT-CL J2106-5844 using weak-lensing analysis with HST

observations

Jinhyub Kim¹, Myungkook James Jee¹, and Jongwan Ko²

¹*Department of Astronomy, Yonsei University,*

²*Korea Astronomy and Space Science Institute*

We present a weak-lensing study of the galaxy cluster SPT-CL J2106-5844 at $z=1.132$ discovered in the South Pole Telescope Sunyaev-Zel'dovich (SPT-SZ) survey. The cluster is claimed to be the most massive system at $z > 1$ in the SPT-SZ survey. The inferred mass ($M_{200c} = (1.27 \pm 0.21) \times 10^{15} M_{\text{sun}}$) is somewhat unusual at such a high redshift given the current Λ CDM prediction. The mass estimates, however, may be biased because the hydrostatic assumption may not hold when the universe was about 40% of the current age. In this work, we reconstruct the dark matter distribution and measure the mass of this interesting cluster using weak-lensing analysis based on the images from the Advanced Camera for Surveys and Wide Field Camera 3 on-board the Hubble Space Telescope. We find that the mass distribution of the cluster is unimodal with no significant substructures. The centroid of the dark matter agrees with both galaxy luminosity and number density distributions, as well as the hot gas centroid. We confirm that the cluster is indeed extremely massive ($M_{200c} = (1.81 \pm 0.47) \times 10^{15} M_{\text{sun}}$) supporting the previous non-lensing measurements. We also discuss the rarity of the cluster in the Λ CDM cosmology, comparing with the expected abundance of similarly massive clusters.

[박 GC-08] KYDISC program: The Impact of Mergers on the Evolution of Galaxies

Sree Oh

Yonsei University

In the hope to detect low-surface brightness features ($\mu_r \sim 27$ mag arcsec⁻²), we carried out KASI-Yonsei Deep Imaging Survey for Clusters (KYDISC) targeting 14 local clusters at $0.016 < z < 0.145$ using Magellan/IMACS telescope and CFHT/MegaCam. Out of 1450 cluster galaxies, 18% of galaxies show the signatures of galaxy mergers. We explore merger-driven changes from various point-of-view. We first examine color-magnitude relations, and find that galaxies related to recent mergers are populated more on blue color than their counterparts. Besides, we find the extremely low frequency of mergers on low-mass

red-sequence galaxies, suggesting a migration of red galaxies into the green-valley region through merger-driven star-formation. We also study the mass-size relation of our sample, finding a larger galaxy size in galaxies related to recent mergers. Our results suggest that mergers can simultaneously change properties of galaxies, making outliers on galactic scaling relations.

[구 GC-09] Environmental effects in the stellar populations of Compact Elliptical galaxies

Suk Kim, Hyunjin Jeong, Youngdae Lee, Seok-Joo Joo, Jaehyun Lee, Eon-Chang Sung

Korea Astronomy & Space Science institute, 776 Daedeokdae-ro, Daejeon 305-348, Korea

Compact elliptical (cE) galaxies are in a rare class of stellar systems characterized by high stellar densities, small sizes, high velocity dispersion, and high metallicity corresponding to elliptical galaxies. cE galaxies have been observed around massive galaxies, so they could be formed under strong influences of tidal stripping and truncation. However, the recent discovery of isolated cE galaxies requires the need of new formation scenarios. We aim at finding cE galaxies in various environments using SDSS DR12, and studying stellar population of cEs as function of environments. Based on the typical properties of cE galaxies, we selected cE candidates by restricting that low-luminosity $M_g > 19.5$ mag, small sizes $R_e < 700$ pc, and high velocity dispersions $\sigma > 60$ km s⁻¹. Since effect radii of cE candidates are mostly smaller than the seeing size of SDSS photometry, we calculated the effective radius by fitting a Sersic profile. In addition, we assumed that host galaxies have brightness with $M_r < -21$ mag, and an environmental parameter is computed as distances between cE galaxies and host-galaxies. We found 112 cE galaxies at $z < 0.05$, which have high sersic indices (mean value is 5.2) similar to the typical massive elliptical galaxies. Mgb values of cE galaxies increase as the distances from the host galaxies decrease. Especially, for cEs close to the host galaxies (NcE: $D_{\text{host}} < 300$ pc), the Mgb values are similar to those of massive elliptical galaxies, which is consistent with the previous studies. On the other hand, cE galaxies distant from the host galaxies (DcE: $D_{\text{host}} > 300$ pc) have lower Mgb values than the conventional cE. The Mgb values follow the σ -Mgb relation of elliptical galaxies, and are connected to its faint end. This can be explained as a result of different merger histories for differing