

Junhyun Baek<sup>1</sup>, Aeree Chung<sup>1</sup>, Kevin Schawinski<sup>2</sup>, Kyuseok Oh<sup>3</sup>, Ivy Wong<sup>4</sup>, Michael Koss<sup>5</sup>, and BASS team

<sup>1</sup>*Department of Astronomy, Yonsei University, South Korea.* <sup>2</sup>*Institute for Particle Physics and Astrophysics, ETH Zurich, Switzerland.*

<sup>3</sup>*Department of Astronomy, Kyoto University, Japan.* <sup>4</sup>*ICRAR, The University of Western Australia, Australia.* <sup>5</sup>*Eureka Scientific Inc., USA*

We have conducted a 22 GHz very long baseline interferometry (VLBI) survey of 281 local ( $z < 0.05$ ) active galactic nuclei (AGNs) selected from the *Swift* Burst Alert Telescope (BAT) 70-month ultra hard X-ray (14–195 keV) catalog. The main goal is to investigate the relation between the strengths of black hole accretion and the parsec-scale nuclear jet, which is expected to tightly correlate but has not been observationally confirmed yet. The BAT AGN Spectroscopic Survey (BASS) provides the least biased AGN sample against obscuration including both Seyfert types, hence it makes an ideal parent sample for studying the nuclear jet properties of an overall AGN population. Using the Korean VLBI Network (KVN), the KVN and VERA Array (KaVA), and the Very Long Baseline Array (VLBA), we observed 281 objects with a 22 GHz flux  $> 30$  mJy, detecting 11 targets (~4% of VLBI detection rate). This implies that the fraction of X-ray AGNs which are currently ejecting a strong nuclear jet is very small. Although our 11 sources span a wide range of pc-scale morphological types, from compact to complex, they lie on a tight linear relation between accretion luminosity and nuclear jet luminosity. Our finding may indicate that the power of nuclear jet is directly responsible for the amount of black hole accretion. We also have probed the fundamental plane of black hole activity in VLBI scale (e.g., few milli-arcsecond). The results from our high-frequency VLBI radio study support that the change of jet luminosity and size follows what is predicted by the AGN evolution scenario based on the Eddington ratio ( $\lambda_{\text{Edd}}$ ) - column density ( $N_{\text{H}}$ ) plane, proposed by a previous study.

#### [7 GC-09] An observed link between AGN Eddington ratio and [NII] $\lambda 6583/\text{H}\alpha$ at $0.6 < z < 1.7$

Kyuseok Oh<sup>1,2</sup>, Yoshihiro Ueda<sup>1</sup>, Masayuki Akiyama<sup>3</sup>, Hyewon Suh<sup>4</sup>, Michael Koss<sup>5</sup>, Richard Mushotzky<sup>6</sup>, Guenther Hasinger<sup>7</sup>, Daichi Kashino<sup>8</sup>, John Silverman<sup>9</sup>

<sup>1</sup>*Kyoto University,* <sup>2</sup>*JSPS Fellow,* <sup>3</sup>*Tohoku University,* <sup>4</sup>*Subaru Telescope,* <sup>5</sup>*Eureka Scientific,* <sup>6</sup>*University of Maryland,* <sup>7</sup>*European Space Agency,* <sup>8</sup>*ETH*

*Zurich,* <sup>9</sup>*Kavli Institute for the Physics and Mathematics of the Universe*

We present the observed relationship between Eddington ratio ( $\lambda_{\text{Edd}}$ ) and optical narrow-emission-line ratio ([NII]  $\lambda 6583/\text{H}\alpha$ ) of X-ray-selected unobscured active galactic nuclei (AGN) at  $0.6 < z < 1.7$  using 27 near-infrared spectra from the Fiber Multi-Object Spectrograph mounted on the Subaru telescope along with 26 additional sources from the literature. We show that the  $\lambda_{\text{Edd}}$  and [NII]  $\lambda 6583/\text{H}\alpha$  ratio at  $0.6 < z < 1.7$  exhibits a similar distribution of  $\lambda_{\text{Edd}}$ -[NII]  $\lambda 6583/\text{H}\alpha$  anti-correlation that has been found for local ( $\langle z \rangle = 0.036$ ), hard X-ray selected AGN. The observed anti-correlation suggests that [N II]  $\lambda 6583/\text{H}\alpha$  optical narrow-line ratio in the AGN host galaxy may carry important information about the accretion state of the central supermassive black hole, suggesting the observational hint of consistent relationship from local to  $z \sim 1.7$ . Further study is necessary to determine whether the  $\lambda_{\text{Edd}}$ -[N II]  $\lambda 6583/\text{H}\alpha$  correlation in high-redshift still holds at  $\log \lambda_{\text{Edd}} < -2$  compared to local AGN.

#### [7 GC-10] Instantaneous AGN feedback at the central part of NGC 5728

Jaejin Shin<sup>1</sup>, Jong-Hak Woo<sup>1</sup>, Aeree Chung<sup>2</sup>, Junhyun Baek<sup>2</sup>, Kyuhyun Cho<sup>1</sup>, Daeun Kang<sup>1</sup>, and Hyun-Jin Bae<sup>1</sup>

<sup>1</sup>*Seoul National University,* <sup>2</sup>*Yonsei University*

Using VLT/MUSE and ALMA data, we present a spatially-resolved analysis of the central part of NGC 5728. We find enhanced star formation ( $\sim 1.8 M_{\odot}/\text{yr}/\text{kpc}^2$ ) at a region where AGN gas outflows intersect the star formation ring. In contrast, significantly weaker CO emission ( $\sim 3.5$  times) is found at the same region compared to other regions in ring, suggesting positive AGN feedback on star formation. On the other hand, we detect gas outflows outside of the spiral arms, implying that the inflowing gas in the arms is removed (i.e., negative feedback). Even though the positive and negative feedback are expected at the central part of NGC 5728, the impact of the AGN feedback in galaxy scale may be insignificant.

#### [7 GC-11] Probing the Feedback Process in Local Type-2 AGNs with Integral-Field Spectroscopy

Rongxin Luo<sup>1</sup>, Jong-Hak Woo<sup>1</sup>, Jaejin Shin<sup>1</sup>, Daeun Kang<sup>1</sup>, Hyun-Jin Bae<sup>1</sup>, Marios Karouzos<sup>2</sup>

<sup>1</sup>*Astronomy Program, Department of Physics and*

*Astronomy, Seoul National University, Seoul 151-742, Republic of Korea*

<sup>2</sup>*Nature Astronomy, Springer Nature, 4 Crinan Street, N1 9XW London, United Kingdom*

Feedback process is one of the most important topics in the study of AGNs since it plays a key role in linking the SMBHs and their host galaxies. In order to further understand the co-evolution of SMBHs and their host galaxies, we probe the feedback process in local type-2 AGNs with a series of integral-field-spectroscopy observations.

In the first part of my talk, I will introduce our GMOS observations of luminous type-2 AGNs at  $z < 0.1$ , which are selected using the integrated [O III] kinematics. Based on the dedicated emission-line diagnostics and kinematic studies, we identify the signatures of AGN-driven outflows and quantify the outflow size in the targets with extreme [O III] kinematics. For the targets without extreme [O III] kinematics, we find the presence of weak AGN-driven outflows, which are indicated by the significant differences between the kinematics of gas and stars.

Then, I will present our recent study of 40 type-2 AGNs based on the SNIFS IFU. By comparing the radial profile of velocity dispersion of gas and stars, we measure the size of AGN-driven outflows in these targets and extend the outflow size-AGN luminosity relation in our previous GMOS studies. We also discuss the feedback effect of AGN-driven outflows by connecting the outflow velocity and host galaxy properties. These results highlight the importance of spatially-resolved observation in investigating gas kinematics and identifying the signatures of AGN-driven outflows.

### [7 GC-12] The Infrared Medium-deep Survey. VI. Discovery of Faint Quasars at $z \sim 5$ with a Medium-band-based Approach

Yongjung Kim<sup>1,2</sup>, Myungshin Im<sup>1,2</sup>, Yiseul Jeon<sup>3</sup>, Minjin Kim<sup>4</sup>, Soojong Pak<sup>1,5</sup> and IMS Team<sup>1</sup>

<sup>1</sup>*Center for the Exploration of the Origin of the Universe (CEOU),* <sup>2</sup>*Seoul National University,*

<sup>3</sup>*FEROKA Inc.,* <sup>4</sup>*Kyungpook National University,*

<sup>5</sup>*Kyung Hee University*

The faint quasars with  $M_{1450} > -24$  mag are known to hold the key to the determination of the ultraviolet emissivity for the cosmic reionization. But only a few have been identified so far because of the limitations on the survey data. Here we present the first results of the  $z \sim 5$  faint quasar survey with the Infrared Medium-deep Survey (IMS), which covers  $\sim 100 \text{ deg}^2$  areas in J band to the depths of  $J_{AB} \sim 23$  mag. To improve selection

methods, the medium-band follow-up imaging has been carried out using the SED camera for QAsars in Early uNiverse (SQUEAN) on the Otto Struve 2.1 m Telescope. The optical spectra of the candidates were obtained with 8 m class telescopes. We newly discovered 10 quasars with  $-25 < M_{1450} < -23$  at  $z \sim 5$ , among which three have been missed in a previous survey using the same optical data over the same area, implying the necessity for improvements in high-redshift faint quasar selection. We derived photometric redshifts from the medium-band data and found that they have high accuracies of  $\langle |\Delta z| / (1 + z) \rangle = 0.016$ . The medium-band-based approach allows us to rule out many of the interlopers that contaminate  $\geq 20\%$  of the broadband-selected quasar candidates. These results suggest that the medium-band-based approach is a powerful way to identify  $z \sim 5$  quasars and measure their redshifts at high accuracy (1%-2%). It is also a cost-effective way to understand the contribution of quasars to the cosmic reionization history.

### [7 GC-13] Assembling the bulge from globular clusters: Evidence from sodium bimodality

Young-Wook Lee, Jenny J. Kim, Chul Chung, Sohee Jang, Dongwook Lim  
*Center for Galaxy Evolution Research & Department of Astronomy, Yonsei University, Seoul 03722, Korea*

Recent investigations of the double red clump in the color-magnitude diagram of the Milky Way bulge cast serious doubts on the structure and formation origin of the outer bulge. Unlike previous interpretation based on an X-shaped bulge, stellar evolution models and CN-band observations have suggested that this feature is another manifestation of the multiple stellar population phenomenon observed in globular clusters (GCs). This new scenario requires a significant fraction of the outer bulge stars with chemical patterns uniquely observed in GCs. Here we show from homogeneous high-quality spectroscopic data that the red giant branch stars in the outer bulge ( $>5.5^\circ$  from the Galactic center) are clearly divided into two groups according to Na abundance in the [Na/Fe] - [Fe/H] plane. The Na-rich stars are also enhanced in Al, while the differences in O and Mg are not observed between the two Na groups. The population ratio and the Na and Al differences between the two groups are also comparable with those observed in metal-rich GCs. Since these chemical patterns and characteristics are only explained by stars originated in GCs, this is compelling evidence that the outer bulge was mostly assembled from