

wide and deep multi-wavelength data. We newly found galaxy cluster candidates at $0.2 < z < 1.4$ and a LSS spanning over 100Mpc at $z \sim 0.9$ in the ELAIS-N1 field which is one of the IMS (Infrared Medium-deep Survey; Im et al. 2019, in preparation) fields. Thanks to K-GMT science program, we performed spectroscopic follow-up observation for a $z \sim 1$ galaxy cluster candidates with GMOS of Gemini North and for $z \sim 0.9$ supercluster candidates with Hectospec of MMT in 2018A and confirmed the large scale structures. We present the newly discovered galaxy overdensities from the observation and the analysis result.

[구 GC-20] Unveiling Quenching History of Cluster Galaxies Using Phase-space Analysis

Jinsu Rhee¹, Rory Smith², Sukyoung, K. Yi¹
¹Yonsei University, ²Korea Astronomy and Space Science Institute

We utilize times since infall of cluster galaxies obtained from Yonsei Zoom-in Cluster Simulation (YZiCS), the cosmological hydrodynamic N-body simulations, and star formation rates from the SDSS data release 10 to study how quickly late-type galaxies are quenched in the cluster environments. In particular, we confirm that the distributions of both simulated and observed galaxies in phase-space diagrams are comparable and that each location of phase-space can provide the information of times since infall and star formation rates of cluster galaxies. Then, by limiting the location of phase-space of simulated and observed galaxies, we associate their star formation rates at $z \sim 0.08$ with times since infall using an abundance matching technique that employs the 10 quantiles of each probability distribution.

Using a flexible quenching model covering different quenching scenarios, we find the star formation history of satellite galaxies that best reproduces the obtained relationship between time since infall and star formation rate at $z \sim 0.08$. Based on the derived star formation history, we constrain the quenching timescale (2 - 7 Gyr) with a clear stellar mass trend and confirm that the refined model is consistent with the "delayed-then-rapid" quenching scenario: the constant delayed phase as ~ 2.3 Gyr and the quenching efficiencies (i.e., e-folding timescale) outside and inside clusters as $\sim 2 - 4$ Gyr ($\propto M_*^{-1}$) and $0.5 - 1.5$ Gyr ($\propto M_*^{-2}$). Finally, we suggest: (i) ram-pressure is the main driver of quenching of satellite galaxies for the local Universe, (ii) the quenching trend on stellar mass at $z > 0.5$

indicates other quenching mechanisms as the main driver.

[구 GC-21] Cosmological QUOKKAS: A new method for measuring distances using an extended KVN to Australia

Jeffrey Hodgson¹, Sang-Sung Lee¹, Benjamin l'Hullier¹, Yannis Lioadkis², Arman Shafieloo¹
¹KASI, ²Stanford

Measuring distances at cosmological scales is one of the most important, yet most difficult to acquire astronomical quantities, allowing astronomers to determine the expansion rate of the universe. Typically, astronomers have sought to find "standard candles" that have a known intrinsic brightness in order to determine their distance. The most well known standard candles are Type Ia supernova and Cepheid variable stars making the so-called "distance ladder". Here we present a method for determining cosmological distances via light travel-time arguments, which can be extended from nearby sources to very high redshift sources.

[박 GC-22] On the Global and Local Environmental Dependence of Type Ia Supernova Luminosity from the Analysis of SALT2 and MLCS2k2 Light-Curve Fitters

Young-Lo Kim^{1,2} and Young-Wook Lee¹
¹Center for Galaxy Evolution Research & Department of Astronomy, Yonsei University, Seoul 03722, Korea
²Université de Lyon, F-69622, Lyon, France; Université de Lyon 1, Villeurbanne; CNRS/IN2P3, Institut de Physique Nucléaire de Lyon

There is growing evidence for the dependence of Type Ia supernova (SN Ia) luminosities on the environments. The origin of this correlation, however, is under debate. In order to explore the physical origin of the trend in detail, we analyze SN Ia light-curves by combining a sample of 1231 SNe Ia over a wide redshift range ($0.01 < z < 1.37$) in various SN surveys and employing two independent light-curve fitters of SALT2 and MLCS2k2. Although SALT2 is the most widely used fitter in the SN community, MLCS2k2 has a novelty in the context of an investigation of the luminosity evolution of SNe Ia. For this reason we use both fitters and analyze them separately. We also determine a stellar mass and a star formation rate (SFR) for a sample of ~ 600 host galaxies. In addition, because recent low-redshift studies suggest that this dependence manifests itself most strongly when using the local SFR at the SN