

total 24 nights observations with large phase angle coverage (From 28 to 104 degrees). The observational results have been published in Kuroda et al. (2021) [7]. We thus found the dominance of submillimeter particles on the surface of Ryugu from the comparison with other meteorite samples from the campaign observation.

In this presentation, we report our activity to modify the TRIPOL for the 1.8-m telescope and the polarimetric performance. We also examine the rotational variability of the polarization degree using the TRIPOL data.

[1] Müller et al. (2017), *Astronomy & Astrophysics*, Volume 599, id.A103.

[2] Delbo et al. (2014), *Nature*, Volume 508, Issue 7495, pp.233-236.

[2] Jaumann et al. (2019), *Science*, Volume 365, Issue 6455, pp.817-820.

[3] Morota et al. (2020), *Science*, Volume 368, Issue 6491, pp. 654-659.

[4] Geake and Dollfus (1986), *Monthly Notices of the Royal Astronomical Society*, Volume 218, pp. 75-91.

[5] Sato et al. (2019), *Research in Astronomy and Astrophysics*, Volume 19, Issue 9, 136.

[6] Kuroda et al. (2021), *Astrophysical Journal Letters* (under review)

[구 SS-14] Study of Hydrated Asteroids via Polarimetry: Correlation between Polarimetric Properties and the Degree of Aqueous Alteration of Hydrated asteroids (편광을 통한 수화한 소행성 연구)

Jooyeon Geem¹(김주연), Masateru Ishiguro¹, Hiroyuki Naito², Daisuke Kuroda³, Koki Takahashi⁴, Tomohiko Sekiguchi⁴, Seiko Takagi⁵, Tatsuharu Ono⁵, Kiyoshi Kuramoto⁵, Tomoki Nakamura⁶
¹Seoul National University, South Korea (서울대학교), ²Nayoro Observatory, Japan, ³Kyoto University, Japan, ⁴Hokkaido University of Education, Japan, ⁵Hokkaido University, Japan ⁶Tohoku University, Japan

Hydrated asteroids get widespread attention for the evolution of water in the Solar System, especially thanks to the recent successes of the Hayabusa2 and OSIRIS-REx space missions. The target asteroids of these missions are believed to be fragments that have experienced aqueous alteration in their parent bodies [3]. Although hydrated asteroids have been studied well via spectroscopy, focusing on the 0.7 um or the 2.7 um absorption bands [2, 3, 4], polarimetric properties of these asteroids have rarely been investigated.

In this study, we conducted a polarimetric

observation of 18 C-complex main-belt asteroids with the 1.6-m Pirka telescope at the Nayoro Observatory of Hokkaido University, Japan. We used a polarimetric imaging mode of the Multi-Spectral Imager (MSI) with the standard Rc-band filter (the central wavelength at 0.64 um) [5]. As a result, we found that all of these hydrated asteroids indicate deep negative branches of their polarimetric profiles. Accordingly, the hydrated asteroids have the polarization minima (Pmin), whose values are significantly lower than any other taxonomic types of asteroids (including C-group asteroids). Because Pmin depends on albedo, particle size, and porosity of the surface materials [1], we suspect that hydrated asteroids are distinctive from other asteroids in terms of these physical properties. In this presentation, we introduce our polarimetric observation and findings. We discuss why hydrated asteroids indicate such low Pmin values, comparing Pmin with spectral features at 0.7 um and 2.7 um based on the observation results.

[1] Cellino et al., 2015, *MNRAS*, 451,4; [2] Fornasier et al., 2014, *Icarus*, 233, 163-178; [3] Rivkin et al. 2002, *Asteroids III*, 1, 235; [4] Takir et al., 2013, *Meteoritics and Planetary Science*, 48, 9; [5] Watanabe et al., 2012, *SPIE*, 8446

[구 SS-15] Isotopic Compositions of Ruthenium Predicted from Stellar Evolution Using the NuGrid Project

Seonho Kim(김선호), Kwang Hyun Sung(성광현), Kyujin Kwak(곽규진)
Physics Department, Ulsan National Institute of Science and Technology, Ulsan 44919, Republic of Korea

Presolar silicon carbide (SiC) grains form around in the envelopes of asymptotic giant branch (AGB) stars by satisfying C/O>1 which is an optimal condition for SiC grains to condense in the stellar outflows. Ruthenium (Ru) isotopes are locked into the SiC grains during the condensation of SiC grains. We investigate the isotopic compositions of Ru in the stellar winds by using the NuGrid data, which are obtained by nucleosynthesis calculations during the stellar evolution. We compare the isotopic compositions of Ru obtained from the NuGrid data with measurements and the predictions obtained from different codes. Our results present a piece of evidence that SiC grains in the presolar system came from low-mass and low-metallicity AGB stars, also confirming that they were not from massive stars. We also suggest a new scenario in which the total stellar yields are also considered because SiC grains can condense during the collapse of molecular clouds.