

the global kinematics, respectively. Using the surface densities and velocity dispersions of the kinematically decomposed HI gas maps together with the rotation curve of NGC 6822, we derive Toomre-Q parameters for individual regions of the galaxy which quantify the level of local gravitational instability of the gaseous disk. We also measure the local star formation rate (SFR) of the corresponding regions in the galaxy by combining GALEX Far-ultraviolet (FUV) and WISE 22 $\mu$ m images. We then relate the gas and SFR surface densities in order to investigate the local Kennicutt-Schmidt (K-S) law of gravitationally unstable regions which are selected from the Toomre Q analysis. Of the three groups, the bulk-narrow, bulk-broad and non-bulk gas components, we find that the lower Toomre-Q values the bulk-narrow gas components have, the more consistent with the linear extension of the K-S law derived from molecular hydrogen (H<sub>2</sub>) observations.

### [포 GC-11] High-resolution mass models of the Large Magellanic Cloud

Shinna Kim<sup>1</sup>, Se-Heon Oh<sup>2\*</sup>, Bi-Qing For<sup>3</sup> and Yun-Kyeong Sheen<sup>4</sup>

<sup>1</sup>*Department of Astronomy and Space Science, Sejong University, Seoul, Korea*

<sup>2</sup>*Department of Physics and Astronomy, Sejong University, Seoul, Korea*

<sup>3</sup>*International Centre for Radio Astronomy Research (ICRAR), University of Western Australia, Crawley, Australia*

<sup>4</sup>*Korea Astronomy and Space Science Institute, Daejeon, Korea*

\*corresponding author: seheon.oh@sejong.ac.kr

We perform disk-halo decomposition of the Large Magellanic Cloud (LMC) using a novel HI velocity field extraction method, aimed at better deriving its HI kinematics and thus mass distribution in the galaxy including both baryons and dark matter. We decompose all the line-of-sight velocity profiles of the combined HI data cube of the LMC, taken from the Australia Telescope Compact Array (ATCA) and Parkes radio telescopes with an optimal number of Gaussian components. For this, we use a novel tool, the so-called BAYGAUD which performs profile decomposition based on Bayesian MCMC techniques. From this, we disentangle turbulent non-ordered HI gas motions from the decomposed gas components, and produce an HI bulk velocity field which better follows the global circular rotation of the galaxy. From a 2D tilted-ring analysis of the HI bulk velocity field, we derive the

rotation curve of the LMC after correcting for its transverse, nutation and precession motions. The dynamical contributions of baryons like stars and gaseous components which are derived using the Spitzer 3.6 micron image and the HI data are then subtracted from the total kinematics of the LMC. Here, we present the bulk HI rotation curve, the mass models of stars and gaseous components, and the resulting dark matter density profile of the LMC.

### [포 GC-12] Searching for Spectrally Variable AGNs using Multi-epoch Spectra from SDSS

Jiyeon Seong<sup>1</sup>, Minjin Kim<sup>1</sup>, Dong-Chan Kim<sup>2</sup>, Il-Sang Yoon<sup>2</sup>, Jaejin Shin<sup>1</sup>

<sup>1</sup>*Department of Astronomy and Atmospheric Science, Kyungpook National University,* <sup>2</sup>*National Radio Astronomy Observatory*

Using multi-epoch spectra of active galactic nuclei (AGN) obtained from the Sloan Digital Sky Survey, we identify 16 spectrally variable sources, for which the spectral shapes of broad emission lines significantly vary with a time scale of yrs. Out of them, 3 AGNs are already known as changing-look (CL) AGNs by previous studies. 6 AGNs are newly identified as CL AGNs from our study. A majority of these AGNs are relatively faint and their variability in the continuum is small, which may explain their non-detection in the previous studies. 7 sources are known as binary AGN candidates based on the systematic velocity offset between broad emission lines and narrow emission lines. For those sources and 3 CL AGNs, we find that the peak of broad emission lines had been shifted up to a few thousands km/s for ~10 years, implying that those can be promising candidates for pc-scale binary AGNs or recoiling black holes. We plan to conduct multiwavelength follow-up studies to nail down the physical origin of the velocity shift.

### [포 GC-13] The strategy to catch more early light curves of supernovae

Jeeun Hwang, Myungshin Im, Gregory S.H. Paek, and IMSNG team

*SNU Astronomy Research Center*

The Intensive Monitoring Survey of Nearby Galaxies (IMSNG) is a high cadence observation program monitoring nearby galaxies at < 50 Mpc with high probabilities of hosting supernovae (SNe). The current number of main IMSNG targets is 60, but with new wide-field facilities joining IMSNG,

there is a possibility of increasing the likelihood of catching the early light curves of SNe among galaxies in the vicinity of the main targets. To test the feasibility of the expansion of the sample galaxies, we examine how much the probability of catching SNe increases by adjusting the field of view of the RASA36 telescope which is one of the IMSNG facilities with a large field of view of 6.25 deg<sup>2</sup>. We calculate supernova rates (SNRs) of galaxies within the FoV that contains main IMSNG galaxies from the stellar mass and star formation rate of the galaxies. Based on the SNRs of these galaxies, we find the best pointing of the telescope towards the highest SNR region. As a result, we present improved total SNR, with respect to the ordinary pointing on average where the IMSNG main target is placed at the center of FoV. The actual observation should be followed to test the effect of this strategy.

#### [포 GC-14] Merging histories of Galaxies in Deep and Wide Images of 7 Abell Clusters with Various Dynamical States

Duho Kim<sup>1</sup>, Yun-Kyeong Sheen<sup>1</sup>, Yara L. Jaffe<sup>2</sup>, Adarsh Ranjan<sup>1</sup>, Sukyoung K. Yi<sup>3</sup> and Rory Smith<sup>1</sup>  
<sup>1</sup>*Korea Astronomy and Space Science Institute, Daedeokdae-ro 776, Yuseoung-gu Daejeon 34055, Republic of Korea,* <sup>2</sup>*Instituto de Fisica y Astronomia, Universidad de Valparaiso, Avda. Gran Bretana 1111 Valparaiso, Chile,* <sup>3</sup>*Yonsei University, Republic of Korea*

Galaxy mergers are known to have been one of the main drivers in galaxy evolution in a wide range of environments. However, in galaxy clusters, high-speed encounters have been believed to undermine the role of mergers as a driver in galaxy evolution. Nonetheless, a high fraction (~38% in Sheen et al. 2012 and ~20% in Oh et al. 2018) of galaxies with post-merging features have been reported in deep (>~28 mag/arcsec<sup>2</sup>) optical surveys of cluster galaxies. The authors argue that these galaxies could have merged outside of the cluster and, later, fallen into the cluster, sustaining their long-lasting post-merging features. On the other hand, when galaxy clusters interact, galaxy orbits might be destabilized resulting in a higher galaxy merger rate. To test this idea, we measure the ongoing-merger fraction of galaxies in deep DECam mosaic data of seven Abell clusters (A754, A2399, A2670, A3558, A3574, A3659 and A3716) with a variety of dynamical states (0.016<z<0.091) for comparison with the ongoing-merger fraction (~4%) from virialized clusters in the literature. We also publish our photometric catalogues of DECam mosaics centered on these clusters in u, g, and

r-band.

### 우주론/암흑물질, 암흑에너지

#### [포 CD-01] Probing the Early Phase of Reionization through LiteBIRD

Kyungjin Ahn<sup>1</sup>, Hina Sakamoto<sup>2</sup>, Kiyotomo Ichiki<sup>2</sup>, Hyunjin Moon<sup>1</sup>, Kenji Hasegawa<sup>2</sup>  
<sup>1</sup>*Chosun University*  
<sup>2</sup>*University of Nagoya*

Cosmic reionization imprints its history on the sky map of the cosmic microwave background (CMB) polarization. Even though mild, the signature of the reionization history during its early phase (z>15) can also impact the CMB polarization. We forecast the observational capability of the LiteBIRD(Lite(Light) satellite for the studies of B-mode polarization and Inflation from cosmic background Radiation Detection), a truly cosmic-variance limited apparatus. We focus on the capability for such an apparatus to probe the partial optical depth of the CMB photons during z>15. We show that LiteBIRD is able to probe this quantity with a modest to high significance, enabling one to tell how efficient the cosmic reionization and star formation were at z>15.

#### [포 CD-02] Detecting the Baryon Acoustic Oscillations in the N-point Spatial Statistics of SDSS Galaxies

Se Yeon Hwang, Sumi Kim, Cristiano G. Sabiu, In Kyu Park  
*Department of Physics, University of Seoul, Seoul 02504, Republic of Korea*

Baryon Acoustic Oscillations (BAO) are caused by acoustic density waves in the early universe and act as a standard ruler in the clustering pattern of galaxies in the late Universe. Measuring the BAO feature in the 2-point correlation function of a sample of galaxies allows us to estimate cosmological distances to the galaxies mean redshift, <z>, which is important for testing and constraining the cosmology model. The BAO feature is also expected to appear in the higher order statistics. In this work we measure the generalized spatial N-point point correlation functions up to 4th order.

We made measurements of the 2, 3, and 4-point correlation functions in the SDSS-III DR12 CMASS data, comprising of 777,202 galaxies. The errors