

**PB-06**

## Rice Cultivars under Salt Stress Show Differential Expression of Genes Related to the Regulation of Na<sup>+</sup>/K<sup>+</sup> Balance

Muhammad Farooq<sup>1</sup>, Kyung-Min Kim<sup>1\*</sup>

<sup>1</sup>Division of Plant Biosciences, School of Applied Biosciences, College of Agriculture and Life Science, Kyungpook National University, Daegu 41566, Korea

### [Introduction]

Rice is one of the most important staple foods for a least half the world's population and is considered a salinity-sensitive crop. Rice cultivation systems are threatened by the effects of climate change because many rice-growing areas are located in vulnerable regions.

### [Materials and Methods]

Four different rice (*O. sativa*) genotypes Pokkali (Gyehwa-20), IR28, Cheongcheong, and Nagdong were provided by the plant molecular breeding lab of Kyungpook National University. Pokkali (Gyehwa-20) is a unique salt-resistant cultivar that is cultivated in water-logged coastal regions, whereas IR28 is salt-sensitive. The cultivars Cheongcheong and Nagdong were used for comparison.

### [Results and Discussion]

we compared two famous rice cultivars namely Nagdong and Cheongcheong, with the salt-tolerant variety Pokkali and salt-sensitive variety IR28. Pokkali and Nagdong cultivars were found to have strong salt tolerance during both seed germination and seedling growth. Under 150-mM NaCl stress, these cultivars showed slight reductions in plant growth, but the plants remained vigorous in comparison to Cheongcheong and IR28. We also found that the salt-tolerant varieties Pokkali and Nagdong enhance their salt tolerance by regulating the Na<sup>+</sup>/K<sup>+</sup> ratio in the roots and shoots. Ion transport-related genes and other micronutrients were also differentially regulated among rice cultivars under salt stress. The findings of this study could help develop new salt-tolerant or salt-sensitive cultivars via CRISPR/Cas9 knockout or overexpression methods in future studies.

### [Acknowledgement]

This work was supported by the National Research Foundation of Korea Grant funded by the Korean Government (NRF-2017R1D1A3B04028676).

\*Corresponding author: Tel. +82-53-950-5711, E-mail. kkm@knu.ac.kr