

NONLINEAR VARIATIONAL INEQUALITIES AND FIXED POINT THEOREMS

SEHIE PARK AND ILHYUNG KIM

1. Introduction

P. Hartman and G. Stampacchia [6] proved the following theorem in 1966: If $f : X \rightarrow R^n$ is a continuous map on a compact convex subset X of R^n , then there exists $x_0 \in X$ such that $\langle fx_0, x_0 - x \rangle \geq 0$ for all $x \in X$. This remarkable result has been investigated and generalized by F.E. Browder [1], [2], W. Takahashi [9], S. Park [8] and others. For example, Browder extended this theorem to a map f defined on a compact convex subset X of a topological vector space E into the dual space E^* ; see [2, Theorem 2]. And Takahashi extended Browder's theorem to closed convex sets in topological vector space; see [9, Theorem 3].

In Section 2, we obtain some variational inequalities, especially, generalizations of Browder's and Takahashi's theorems. The generalization of Browder's is an earlier result of the first author [8].

In Section 3, using Theorem 1, we improve and extend some known fixed point theorems. Theorems 4 and 8 improve Takahashi's results [9, Theorems 5 and 9], respectively. Theorem 4 extends the first author's fixed point theorem [8, Theorem 8] (Theorem 5 in this paper) which is a generalization of Browder [1, Theorem 1]. Theorem 8 extends Theorem 9 which is a generalization of Browder [2, Theorem 3].

Finally, in Section 4, we obtain variational inequalities for multi-valued maps by using Theorem 1. We improve Takahashi's results [9, Theorems 21 and 22] which are generalizations of Browder [2, Theorem 6] and the Kakutani fixed point theorem [7], respectively.

Received October 26, 1988.

Supported by a grant from the Korea Science and Engineering Foundation, 1987-88.

2. Variational inequalities

Throughout this paper, we assume that a topological space is Hausdorff and a topological vector space is real. Let us start with the following useful theorem. We deduce this from the Brouwer fixed point theorem.

THEOREM 1. *Let X be a nonempty compact convex subset of a topological vector space E and f a real valued function on $X \times X$ satisfying:*

- (i) *For each $y \in X$, the function $f(x, y)$ of x is lower semicontinuous;*
- (ii) *for each $x \in X$, the function $f(x, y)$ of y is quasi-concave; and*
- (iii) *$f(x, x) \leq c$ for all $x \in X$ with some real number c .*

Then there exists an $x_0 \in X$ such that $f(x_0, y) \leq c$ for all $y \in X$.

Proof. Suppose that for each $x \in X$, there exists $y \in X$ such that $f(x, y) > c$. Then for each $y \in X$, the set $U_y = \{x \in X : f(x, y) > c\}$ is open by (i), and $\{U_y\}_{y \in X}$ is a cover of X . Since X is compact, there exists a finite family $\{y_1, y_2, \dots, y_n\}$ such that $\{U_{y_i}\}_{i=1}^n$ covers X . Let $\{\beta_1, \beta_2, \dots, \beta_n\}$ be a partition of unity subordinated to this subcover. Then each β_i is a continuous map of X into $[0, 1]$ which vanishes outside U_{y_i} , while $\sum_{i=1}^n \beta_i(x) = 1$ for all $x \in X$. For each i satisfying $\beta_i(x) \neq 0$, x lies in U_{y_i} , so that $f(x, y_i) > c$. By (ii) we have

$$f\left(x, \sum_{i=1}^n \beta_i(x) y_i\right) > c$$

for all $x \in X$. Define a continuous map p of X into the convex hull of $\{y_1, y_2, \dots, y_n\}$ by

$$p(x) = \sum_{i=1}^n \beta_i(x) y_i.$$

Since the convex hull of $\{y_1, y_2, \dots, y_n\}$ is a compact convex subset of X which lies in a finite dimensional subspace of E , by the Brouwer fixed point theorem, we have $x_1 \in X$ such that $x_1 = p(x_1) = \sum_{i=1}^n \beta_i(x_1) y_i$. Hence we have

$$c \geq f(x_1, x_1) = f\left(x_1, \sum_{i=1}^n \beta_i(x_1) y_i\right) > c,$$

which is a contradiction.

Theorem 1 improves Takahashi [9, Lemma 1]. From Theorem 1, we obtain the following due to Fan [5] by setting $g(x, y) = f(x, x) - f(x, y)$ on $X \times X$.

COROLLARY 1. *Let X be a nonempty compact convex subset of a topological vector space E and f a real valued continuous function on $X \times X$ such that for each $x \in X$, the function $f(x, y)$ of y is quasi-convex. Then there exists $x_0 \in X$ such that $f(x_0, x_0) \leq f(x_0, y)$ for all $y \in X$.*

Let X be a convex subset of a vector space E over R . For each $x \in X$, the inward and outward sets of X at x , $I_X(x)$ and $O_X(x)$, are defined as follows:

$$I_X(x) := \{x + r(u - x) \in E : u \in X, r > 0\},$$

$$O_X(x) := \{x - r(u - x) \in E : u \in X, r > 0\}.$$

If E is a topological vector space, the closures of $I_X(x)$ and $O_X(x)$ are denoted by $\bar{I}_X(x)$ and $\bar{O}_X(x)$, respectively. In the sequel, $W(x)$ denotes either $\bar{I}_X(x)$ or $\bar{O}_X(x)$.

In [8], the first author obtained the following result by using Corollary 1.

COROLLARY 2. [8] *Let X be a nonempty compact convex subset of a topological vector space E and f a real valued continuous function on $X \times E$ such that for each $x \in X$, the function $f(x, y)$ of y is convex. Then there exists an $x_0 \in X$ such that $f(x_0, x_0) \leq f(x_0, y)$ for all $y \in \bar{I}_X(x_0)$.*

By using this, the first author proved the following:

THEOREM 2. [8] *Let X be a nonempty compact convex subset of a topological vector space E and f a continuous map of X into E^* . Then there exists an $x_0 \in X$ such that $\langle fx_0, x_0 - y \rangle \geq 0$ for all $y \in W(x_0)$.*

In particular, if E is locally convex and $W(x_0)$ is replaced by X , then Theorem 2 reduces to Browder [2, Theorem 2]. In [9, Theorem 3], Takahashi generalized Browder [2, Theorem 2] to closed convex sets in topological vector spaces. In the following theorem, we improve Takahashi's result. Let H, X be nonempty subsets of a topological vector space E . We put $B_H X = \bar{X} \cap \overline{H - X}$ and $I_H X = X \cap (B_H X)^c$ where \bar{A} is the closure of $A \subset E$ and A^c is the complement of A .

THEOREM 3. *Let H be a closed convex subset of a topological vector space E and f a continuous map of H into E^* . If there exists a compact convex subset X of H such that $I_H X \neq \phi$ and for each $z \in B_H X$, there is $u_0 \in I_H X$ with $\langle fz, z - u_0 \rangle \geq 0$, then there exists $x^* \in H$ such that $\langle fx^*, y - x^* \rangle \geq 0$ for all $y \in \bar{I}_H(x^*)$.*

Proof. By Theorem 2, there exists $x^* \in X$ such that $\langle fx^*, y - x^* \rangle \geq 0$ for all $y \in I_X(x^*)$. If $x^* \in I_H X$, for each $y \in H$, we can choose λ ($0 < \lambda < 1$) so that $x = x^* + \lambda(y - x^*)$ lies in X since the map $p(\lambda) = x^* + \lambda(y - x^*)$ is continuous. Then $y = x^* + (x - x^*)/\lambda$ lies in $I_X(x^*)$. Hence we obtain $\langle fx^*, y - x^* \rangle \geq 0$ for all $y \in H$. If $x^* \in B_H X$, by the hypothesis, there exists $u_0 \in I_H X$ with $\langle fx^*, x^* - u_0 \rangle \geq 0$. Since $\langle fx^*, x - x^* \rangle \geq 0$ for all $x \in X$, it follows that

$$\langle fx^*, x - u_0 \rangle \geq 0$$

for all $x \in X$. Since $u_0 \in I_H X$, for each $y \in H$ there exists λ ($0 < \lambda < 1$) such that $x = u_0 + \lambda(y - u_0) \in X$. Hence

$$0 \leq \langle fx^*, x - u_0 \rangle = \lambda \langle fx^*, y - u_0 \rangle$$

and consequently $\langle fx^*, y - u_0 \rangle \geq 0$ for all $y \in H$. Since $u_0 \in X$ implies $\langle fx^*, u_0 - x^* \rangle \geq 0$, we obtain $\langle fx^*, y - x^* \rangle \geq 0$ for all $y \in H$. For $y \in I_H(x^*) \setminus H$, $y = x^* + r(u - x^*)$ for some $u \in H$, $r > 1$. So $\langle fx^*, y - x^* \rangle = r \langle fx^*, u - x^* \rangle \geq 0$. Hence $\langle fx^*, y - x^* \rangle \geq 0$ for all $y \in \bar{I}_H(x^*)$.

3. Fixed point theorems

In this section, using Theorem 1, we improve and extend some known fixed point theorems.

THEOREM 4. *Let X be a nonempty compact convex subset of a topological vector space E and f a continuous map of X into E . Then, either there exists $y_0 \in X$ such that y_0 and fy_0 cannot be separated by a continuous linear functional, or there exist $x_0 \in X$ and $g \in E^*$ such that*

$$g(x_0 - fx_0) < 0 \leq \inf_{y \in W(x_0)} g(x_0 - y).$$

Proof. Suppose that for each $x \in X$, there exists $h \in E^*$ such that $h(x - fx) < 0$. Setting $U_h = \{x \in X : h(x - fx) < 0\}$ for each $h \in E^*$, we have a cover $\{U_h\}_{h \in E^*}$ of X . Since X is compact, there exists a finite family $\{h_1, h_2, \dots, h_n\}$ such that $\{U_{h_i}\}_{i=1}^n$ covers X . Let $\{\beta_1, \beta_2, \dots, \beta_n\}$ be a partition of unity subordinated to this subcover. Define a real valued function p on $X \times E$ by

$$p(x, y) = \sum_{i=1}^n \beta_i(x) h_i(x - y).$$

Then, by Corollary 2, there exists $x_0 \in X$ such that

$$p(x_0, y) = \sum_{i=1}^n \beta_i(x_0) h_i(x_0 - y) \geq p(x_0, x_0) = 0$$

for all $y \in \bar{I}_X(x_0)$. On the other hand, we have

$$p(x_0, fx_0) = \sum_{i=1}^n \beta_i(x_0) h_i(x_0 - fx_0) < 0.$$

By putting $g = \sum_{i=1}^n \beta_i(x_0)h_i$, we obtain the desired result for inward case.

For outward case, define a continuous map $f' : X \rightarrow E$ by $f'x = 2x - fx$. Then, by the preceding inward case, either there exists $y_0 \in X$ such that y_0 and $f'y_0$ cannot be separated by a continuous linear functional, or there exist $x_0 \in X$ and $g' \in E^*$ such that $g'(x_0 - f'x_0) < 0 \leq \inf_{z \in I_X(x_0)} g'(x_0 - z)$. The first alternative implies that y_0 and fy_0 cannot be separated by a continuous linear functional. Suppose that the second one holds. For any $y \in O_X(x_0)$, $z = 2x_0 - y$ lies in $I_X(x_0)$. Then we have

$$\begin{aligned} (-g')(x_0 - fx_0) &= (-g')(f'x_0 - x_0) = g'(x_0 - f'x_0) < 0 \\ &\leq g'(x_0 - z) = g'(y - x_0) = (-g')(x_0 - y) \end{aligned}$$

for any $y \in O_X(x_0)$, and hence for any $y \in \bar{O}_X(x_0)$. By putting $g = -g'$, we obtain the desired result for outward case.

Theorem 4 improves Takahashi [9, Theorem 5]. As a consequence of Theorem 4, we have the following:

THEOREM 5. [8] *Let X be a nonempty compact convex subset of a topological vector space E having sufficiently many linear functionals and f a continuous map of X into E . If for each $x \in X$, there exists $\lambda < 1$ with $\lambda x + (1 - \lambda)fx \in W(x)$, then f has a fixed point.*

Proof. Suppose f has no fixed point. By Theorem 4 there exist $x_0 \in X$ and $g \in E^*$ such that

$$g(x_0 - fx_0) < 0 \leq \inf_{y \in W(x_0)} g(x_0 - y).$$

For this x_0 , we can choose $\lambda < 1$ with $y_0 := \lambda x_0 + (1 - \lambda)fx_0 \in W(x_0)$. Hence we have

$$g(x_0 - fx_0) < 0 \leq g(x_0 - y_0) = (1 - \lambda)g(x_0 - fx_0).$$

This is a contradiction. Therefore f has a fixed point.

In particular, if E is locally convex and $W(x)$ is replaced by X , then Theorem 5 reduces to Browder [1, Theorem 1]. On the other hand, if f maps X into itself, we obtain the following:

COROLLARY 3. [3] *Let X be a nonempty compact convex subset of a topological vector space E having sufficiently many linear functionals and f a continuous map of X into itself. Then f has a fixed point.*

As another consequence of Theorem 4, we have the following:

THEOREM 6. *Let H be a closed convex subset of a topological vector space E having sufficiently many linear functionals and f a continuous map of H into H . If there exists a compact convex subset X of H such that for each $x \in B_H X$, there is $\lambda < 1$ with $\lambda x + (1 - \lambda)fx \in \bar{I}_X(x)$, then f has a fixed point in H .*

Proof. Consider the restriction of f to X . If f has no fixed point in X , by Theorem 4 there exist $x_0 \in X$ and $g \in E^*$ such that

$$g(x_0 - fx_0) < 0 \leq \inf_{y \in W(x_0)} g(x_0 - y).$$

If $x_0 \in I_H X$, since $fx_0 \in H$, we can choose λ ($0 < \lambda < 1$) so that $y_0 = \lambda x_0 + (1 - \lambda)fx_0$ lies in X . Hence we have

$$g(x_0 - fx_0) < 0 \leq g(x_0 - y_0) = (1 - \lambda)g(x_0 - fx_0).$$

This is a contradiction. If $x_0 \in B_H X$, by the hypothesis, there exists $\lambda < 1$ with $y_0 = \lambda x_0 + (1 - \lambda)fx_0 \in \bar{I}_X(x_0)$. Also we have

$$g(x_0 - fx_0) < 0 \leq g(x_0 - y_0) = (1 - \lambda)g(x_0 - fx_0),$$

which is a contradiction. Therefore f has a fixed point.

In particular, if E is locally convex and $W(x) := \bar{I}_X(x)$ is replaced by X , then Theorem 6 reduces to Takahashi [9, Theorem 7].

We now generalize Theorem 4 to multi-valued maps. The following definition is due to Fan [4]. Let X be a subset of a topological vector space E . A map T of X into 2^E is said to be upper demicontinuous if for each open half-space H in E , the set $\{x \in X : Tx \subset H\}$ is open in X . An open half-space H in E is a set of the form $\{x \in E : hx > r\}$ where h is a continuous linear functional, not identically zero, and r is a real number. It is obvious that if a map T of X into 2^E is upper semicontinuous, then T is upper demicontinuous. We say that two sets A, B in E can be strictly separated by a closed hyperplane, if there exist $h \in E^*$ and $r \in \mathbb{R}$ such that $hx < r$ for all $x \in A$ and $hy > r$ for all $y \in B$. For two sets C, D in R , $C < D$ means that $x < y$ for any $x \in C$ and $y \in D$.

THEOREM 7. *Let X be a nonempty compact convex subset of a topological vector space E . Let S, T be two upper demicontinuous maps of X into 2^E such that for each $x \in X$, Sx and Tx are nonempty. Then, either there exists $y_0 \in X$ for which Sy_0 and Ty_0 cannot be strictly separated by a closed hyperplane, or there exist $x_0 \in X$ and $g \in E^*$ such that $g(x_0 - Tx_0) < g(x_0 - Sx_0)$ and $0 \leq \inf_{y \in W(x_0)} g(x_0 - y)$.*

Proof. Suppose that for each $x \in X$, Sx and Tx can be strictly separated by a closed hyperplane. Then for each $x \in X$, we can find $g_x \in E^*$ and $r_x \in R$ such that $g_x(Sx) < r_x < g_x(Tx)$. Since S, T are upper demicontinuous on X , there exists a neighborhood U_x of x in X such that $g_x(Sy) < r_x < g_x(Ty)$ for all $y \in U_x$. Hence x is in the interior $N(g_x)$ of $\{z \in X : g_x(Sz) < g_x(Tz)\}$. Thus $X = \bigcup_{x \in X} N(g_x)$. By compactness of X , there exists a finite set $\{x_1, x_2, \dots, x_n\} \subset X$ such that $X = \bigcup_{i=1}^n N(g_{x_i})$. Let $\{\beta_i\}_{i=1}^n$ be a partition of unity subordinated to the cover $\{N(g_{x_i})\}$. Define a real valued function p on $X \in E$ by

$$p(x, y) = \sum_{i=1}^n \beta_i(x) g_{x_i}(x - y).$$

By Corollary 2, there exists $x_0 \in X$ such that

$$p(x_0, y) = \sum_{i=1}^n \beta_i(x_0) g_{x_i}(x_0 - y) \geq p(x_0, x_0) = 0$$

for all $y \in \bar{I}_X(x_0)$. We also know that

$$\sum_{i=1}^n \beta_i(x_0) g_{x_i}(Sx_0) < \sum_{i=1}^n \beta_i(x_0) g_{x_i}(Tx_0).$$

By putting $g = \sum \beta_i(x_0) g_{x_i}$, we obtain the desired result for inward case.

For outward case, define upper demicontinuous maps $S', T' : X \rightarrow 2^E$ by $S'x = 2x - Sx$, $T'x = 2x - Tx$, respectively. By the preceding inward case, either there exists $y_0 \in X$ for which $S'y_0$ and $T'y_0$ cannot be strictly separated by a closed hyperplane, or there exist $x_0 \in X$ and $g' \in E^*$ such that $g'(x_0 - T'x_0) < g'(x_0 - S'x_0)$ and $0 \leq \inf_{z \in W(x_0)} g'(x_0 - z)$.

The first alternative implies that Sy_0 and Ty_0 cannot be strictly separated by a closed hyperplane. Suppose that the second one holds.

For any $y \in O_X(x_0)$, $z = 2x_0 - y$ lies in $I_X(x_0)$. Then we have

$$\begin{aligned} (-g')(x_0 - T'x_0) &= (-g')(T'x_0 - x_0) = g'(x_0 - T'x_0) \\ &< g'(x_0 - S'x_0) = (-g')(S'x_0 - x_0) = (-g')(x_0 - Sx_0), \end{aligned}$$

and

$$0 \leq g'(x_0 - z) = g'(y - x_0) = (-g')(x_0 - y)$$

for any $y \in O_X(x_0)$, and hence for any $y \in \bar{O}_X(x_0)$. By putting $g = -g'$, we obtain the desired result for outward case.

Theorem 7 improves Takahashi [9, Theorem 8]. If S is the identity map of X , then Theorem 7 reduces to the following generalization of Theorem 4.

THEOREM 8. *Let X be a nonempty compact convex subset of a topological vector space E and T an upper demicontinuous map of X into 2^E such that for each $x \in X$, Tx is nonempty. Then, either there exists $y_0 \in X$ such that y_0 and Ty_0 cannot be strictly separated by a closed hyperplane, or there exist $x_0 \in X$ and $g \in E^*$ such that*

$$g(x_0 - Tx_0) < 0 < \inf_{y \in W(x_0)} g(x_0 - y).$$

Theorem 8 improves Takahashi [9, Theorem 9]. As a consequence of Theorem 8, we have the following theorem.

THEOREM 9. *Let X be a nonempty compact convex subset of a locally convex topological vector space E and T an upper demicontinuous map X into 2^E such that for each $x \in X$, Tx is nonempty, closed and convex. If for each $x \in X$, there exists $\lambda < 1$ such that $(\lambda x + (1-\lambda)Tx) \cap W(x) \neq \emptyset$, then T has a fixed point.*

Proof. Suppose T has no fixed point. By Theorem 8 there exist $x_0 \in X$ and $g \in E^*$ such that

$$g(x_0 - Tx_0) < 0 \leq \inf_{y \in W(x_0)} g(x_0 - y).$$

For this x_0 , we can choose $\lambda < 1$ and $z_0 \in Tx_0$ such that $y_0 := \lambda x_0 + (1-\lambda)z_0 \in W(x_0)$. Hence we have

$$g(x_0 - z_0) < 0 \leq g(x_0 - y_0) = (1-\lambda)g(x_0 - z_0).$$

This is a contradiction. Therefore T has a fixed point.

In particular, if T is upper semicontinuous and $W(x)$ is replaced by X , then Theorem 9 reduces to Browder [2, Theorem 3].

From Corollary 2 for a normed vector space, we obtain the following generalization of Ky Fan [4, Theorem 2].

THEOREM 10. *Let X be a nonempty compact convex subset of a normed vector space E and f a continuous map of X into E . Then there exists $x_0 \in X$ such that*

$$\|fx_0 - x_0\| = \min_{y \in W(x_0)} \|fx_0 - y\|.$$

Proof. Define a real valued function g on $X \times E$ by $g(x, y) = \|fx - y\|$. Then g is continuous and for each $x \in X$, the function $g(x, y)$ of y is convex. Thus the desired result is obvious by Corollary 2.

4. Variational inequalities for multi-valued maps

By using Theorem 1, we generalize Theorem 2 to multi-valued

maps for inward case.

THEOREM 11. *Let X be a nonempty compact convex subset of a topological vector space E and T an upper semicontinuous map of X into 2^{E^*} such that for each $x \in X$, Tx is nonempty and compact. If for each $x \in X$,*

$$\min_{y \in X} \max_{g \in Tx} \langle g, x-y \rangle = \max_{g \in Tx} \min_{y \in X} \langle g, x-y \rangle,$$

then there exist $x_0 \in X$ and $g_0 \in Tx_0$ such that $\langle g_0, x_0-y \rangle \geq 0$ for all $y \in \bar{I}_X(x_0)$.

Proof. Define a real valued function f on $X \times X$ by

$$f(x, y) = \max_{g \in Tx} \langle g, x-y \rangle.$$

For any $y \in X$ and $c \in R$, put $A = \{x \in X : f(x, y) \geq c\}$. We show that if $\{x_\alpha : \alpha \in I\}$ is a net in A converging to x_0 , then $x_0 \in A$. For each x_α there exists $g_\alpha \in Tx_\alpha$ such that $\langle g_\alpha, x_\alpha-y \rangle \geq c$. Since $\cup\{Tx : x \in X\}$ is compact, $\{g_\alpha\}$ has a subnet $\{g_{\alpha'}\}$ converging to g_0 . Since T is upper semicontinuous, $g_0 \in Tx_0$. Also we have $c \leq \lim_{\alpha'} \langle g_{\alpha'}, x_{\alpha'}-y \rangle = \langle g_0, x_0-y \rangle$. Hence $x_0 \in A$. That is, the function $f(x, y)$ of x is upper semicontinuous. It is obvious that the function $f(x, y)$ of y is convex and $f(x, x) = 0$ for all $x \in X$. By Theorem 1 for $-f$, there exists $x_0 \in X$ such that $\max_{g \in Tx_0} \langle g, x_0-y \rangle \geq 0$ for all $y \in X$. Since

$$\min_{y \in X} \max_{g \in Tx_0} \langle g, x_0-y \rangle = \max_{g \in Tx_0} \min_{y \in X} \langle g, x_0-y \rangle,$$

we have $g_0 \in Tx_0$ such that $\langle g_0, x_0-y \rangle \geq 0$ for all $y \in X$. For $y \in I_X(x_0) \setminus X$, $y = x_0 + r(u-x_0)$ for some $u \in X$ and $r > 1$. So $\langle g_0, x_0-y \rangle = r \langle g_0, x_0-u \rangle \geq 0$. Hence $\langle g_0, x_0-y \rangle \geq 0$ for all $y \in \bar{I}_X(x_0)$.

Theorem 11 improves Takahashi [9, Theorem 21]. In particular, if Tx is convex, the minimax equality in Theorem 11 holds. So we have the following corollary which is an extension of Browder [2, Theorem 6].

COROLLARY 4. *Let X be a nonempty compact convex subset of a topological vector space E and T an upper semicontinuous map of X into 2^{E^*} such that for each $x \in X$, Tx is nonempty, compact and convex. Then there exist $x_0 \in X$ and $g_0 \in Tx_0$ such that $\langle g_0, x_0-y \rangle \geq 0$ for all $y \in \bar{I}_X(x_0)$.*

Proof. We need only show that for each $x \in X$,

$$\min_{y \in X} \max_{g \in Tx} \langle g, x-y \rangle = \max_{g \in Tx} \min_{y \in X} \langle g, x-y \rangle.$$

Let $x \in X$ and $c = \max_{g \in Tx} \min_{y \in X} \langle g, x - y \rangle$. For each $g \in Tx$, put $A(g) = \{y \in X : \langle g, x - y \rangle \leq c\}$. Let $\{g_1, g_2, \dots, g_n\}$ be a finite subset of Tx and $\{r_1, r_2, \dots, r_n\}$ be nonnegative numbers with $\sum_{i=1}^n r_i = 1$. For $\sum_{i=1}^n r_i g_i \in Tx$, there is $y_0 \in X$ such that $\sum_{i=1}^n r_i \langle g_i, x - y_0 \rangle = \langle \sum_{i=1}^n r_i g_i, x - y_0 \rangle \leq c$. Thus there exists $z \in X$ such that $\langle g_i, x - z \rangle \leq c$ for $i = 1, 2, \dots, n$. Since the family $\{A(g) : g \in Tx\}$ has the finite intersection property and X is compact, we have $\bigcap \{A(g) : g \in Tx\} \neq \emptyset$. Let $y_0 \in \bigcap \{A(g) : g \in Tx\}$. Then $\max_g \langle g, x - y_0 \rangle \leq c$. Hence we have

$$\min_y \max_g \langle g, x - y \rangle \leq \max_g \langle g, x - y_0 \rangle \leq \max_g \min_y \langle g, x - y \rangle.$$

On the other hand it is obvious that

$$\max_g \min_y \langle g, x - y \rangle \leq \min_y \max_g \langle g, x - y \rangle.$$

THEOREM 12. *Let X be a nonempty compact convex subset of Euclidean space R^n and T an upper semicontinuous map of X into 2^{R^n} such that for each $x \in X$, Tx is nonempty and compact. If for each $x \in X$,*

$$\min_{y \in X} \max_{z \in Tx} \langle z - x, x - y \rangle = \max_{z \in Tx} \min_{y \in X} \langle z - x, x - y \rangle,$$

then there exist $x_0 \in X$ and $z_0 \in Tx_0$ such that $\langle z_0 - x_0, x_0 - y \rangle \geq 0$ for all $y \in \bar{I}_X(x_0)$.

Proof. Setting $f(x, y) = \max_{z \in Tx} \langle z - x, x - y \rangle$ for $x, y \in X$ and applying the argument in Theorem 11, we obtain the desired result.

Theorem 12 improves Takahashi [9, Theorem 2.2].

COROLLARY 5. *Let X be a nonempty compact convex subset of Euclidean space R^n and T an upper semicontinuous map of X into 2^{R^n} such that for each $x \in X$, Tx is nonempty, compact and convex. Then there exist $x_0 \in X$ and $z_0 \in Tx_0$ such that $\langle z_0 - x_0, x_0 - y \rangle \geq 0$ for all $y \in \bar{I}_X(x_0)$.*

Proof. We need only show that for each $x \in X$,

$$\min_{y \in X} \max_{z \in Tx} \langle z - x, x - y \rangle = \max_{z \in Tx} \min_{y \in X} \langle z - x, x - y \rangle.$$

This follows from the argument in Corollary 4.

In Theorem 12, if T is a map of X into 2^X , by putting $y = z_0$, we obtain $z_0 = x_0$, that is, $x_0 \in Tx_0$. In particular, the Kakutani fixed point theorem is obtained.

COROLLARY 6. [7] *Let X be a nonempty compact convex subset of Eulidean space R^n and T an upper semicontinuous map of X into 2^X such that for each $x \in X$, Tx is nonempty, compact and convex. Then T has a fixed point.*

References

1. F.E. Browder, *On a new generalization of the Schauder fixed point theorem*, Math. Ann. **174**(1967), 285-290.
2. F.E. Browder, *The fixed point theory of multi-valued mappings in topological vector spaces*, Math. Ann. **177**(1968), 283-301.
3. K. Fan, *Sur un theoreme minimax*, C.R. Acad. Sci. Paris, **259**(1962), 3925-3928.
4. K. Fan, *Extensions of two fixed point theorems of F.E. Browder*, Math. Z. **112**(1969), 234-240.
5. K. Fan, *A minimax inequality and applications*, Inequalities III (Ed. O. Shisha), Academic Press, 1972, 103-113.
6. P. Hartman and G. Stampacchia, *On some nonlinear elliptic differential functional equations*, Acta Math. **115**(1966), 271-310.
7. S. Kakutani, *A generalization of Brouwer's fixed point theorem*, Duke Math. J. **8**(1941), 457-459.
8. S. Park, *Fixed point theorems on compact convex sets in topological vector spaces*, Contemporary Math. **72**(1988), 183-191.
9. W. Takahashi, *Nonlinear variational inequalities and fixed point theorems*, J. Math. Soc. Japan **28**(1976), 168-181.

Seoul National University

and

The Mathematical Sciences Research Institute of Korea

Seoul 151-742, Korea