

je Research and Scholarship Foundation in 1989.

References

- (a) M. L. Bender and M. Komiyama, "Cyclodextrin Chemistry", Springer-Verlag, New York (1977); (b) I. Tabushi, *Acc. Chem. Res.*, **15**, 66 (1982).
- (a) R. L. Van Etten, J. F. Sebastian, G. A. Clowes, and M. L. Bender, *J. Am. Chem. Soc.*, **89**, 3242, 3253 (1967); (b) H. J. Brass and M. L. Bender, *ibid.*, **95**, 5391 (1973); (c) O. S. Tee and J. M. Bennett, *ibid.*, **110**, 269 (1988).
- (a) R. Breslow, G. L. Trainor, and A. Ueono, *J. Am. Chem. Soc.*, **105**, 2739 (1983); (b) E. Anslyn and R. Breslow, *J. Am. Chem. Soc.*, **111**, 5972 (1989); (c) A. V. Veglia and R. H. de Rossi, *J. Org. Chem.*, **53**, 5281 (1988); (d) K. R. Rao and P. B. Sattur, *J. Chem. Soc., Chem. Commun.*, 342 (1989); (e) A. B. Tabor, A. B. Holmes, and R. Baker, *ibid.*, 1025 (1989).
- (a) V. Daffe and J. Fastrez, *J. Chem. Soc., Perkin II*, 789 (1983); (b) R. Fornasier, P. Scrimin, and U. Tonellato, *Tetra. Lett.*, **24**, 5541 (1983).
- (a) F. Cramer and G. Mackensen, *Angew. Chem.*, **78**, 641 (1966); (b) Y. Kitaura and M. L. Bender, *Bioorg. Chem.*, **4**, 237 (1975); (c) R. Breslow, P. Bovy, and C. L. Hersch, *J. Am. Chem. Soc.*, **102**, 2115 (1980); (d) I. Tabushi and Y. Kuroda, *J. Am. Chem. Soc.*, **106**, 4580 (1984); (e) N. Tanaka, A. Yamaguchi, Y. Araki, *Chem. Lett.*, 715 (1987).
- (a) Y. Iwakura, K. Uno, F. Toda, S. Onojuka, K. Hattori, and M. L. Bender, *J. Am. Chem. Soc.*, **97**, 4432 (1975); (b) H. Ikeda, R. Kojin, C.-J. Yoon, T. Ikeda, and F. Toda, *Chem. Lett.*, 1495 (1987); (c) M. Komiyama, *J. Am. Chem. Soc.*, **111**, 3046 (1989); (d) E. U. Akkaya and A. W. Czarnik, *ibid.*, **110**, 8553 (1988).
- K. Ohkubo, Y. Nakano, and H. Nagamura, *J. Mol. Catal.*, **29**, 1 (1985).
- (a) R. J. Bergeron, M. A. Channing, G. J. Gibeily, and D. M. Pillor, *J. Am. Chem. Soc.*, **99**, 5146 (1977); (b) Y. Inoue, F.-H. Kuan, and R. Chujo, *Bull. Chem. Soc. Jpn.*, **60**, 2539 (1987).
- The preparation of substrates and kinetic methods have been described elsewhere. (a) I. Cho and J.-S. Shin, *Makromol. Chem.*, **184**, 147 (1983); (b) I. Cho and B.-J. Lee, *Polym. Bull.*, **17**, 383 (1987); (c) I. Cho and G.-C. Kim, *J. Org. Chem.*, **53**, 5187 (1988).

Benzophenone Sensitized Photoisomerization of 1,2-Bispyrazinylethylene

Pill-Hoon Bong and Sang Chul Shim*

Department of Chemistry, Jeon Ju University, Jeon Ju 560-759

*Department of Chemistry, Korea Advanced Institute of Science & Technology, Seoul 130-650

Received January 8, 1990


We have proposed a triplet mechanism for direct *trans* → *cis* photoisomerization of 1,2-bispyrazinylethylene (BPYE) on the basis of benzophenone sensitization and azulene quenching studies.¹ Benzophenone sensitizer, however, showed an anomalous concentration effect on the photoisomerization quantum yields^{1b} which could not be clearly explained at that time. We now propose a plausible mechanism to explain this anomaly.

A simple analysis of the data^{1b} shows a decrease of benzophenone sensitized *trans* → *cis* photoisomerization quantum yields as the concentration of benzophenone increases indicating that not all the benzophenone triplets generate the olefin triplets. The simplest mechanism which will account for this observation is shown in Scheme 1.

When the steady-state approximation is applied on the benzophenone sensitized *trans* → *cis* photoisomerization,

$$\Phi_{r-c}^{obs} = f(1-\alpha)\Phi_{isc} + (1-f)(1-\alpha) \frac{k_{et}(t)}{k_{nr} + k_{et}(t) + k_{sq}(B)} \quad (1)$$

and when azulene is added as a triplet quencher, the follow-



Scheme 1. Mechanism of the *trans* → *cis* photoisomerization of *trans*-1,2-bispyrazinylethylene.

ing Stern-Volmer relationship can be derived,

$$\Phi_{r-c}^0 / \Phi_{r-c} = \frac{(1+k_q(Az)/k_{iso})(f\Phi_{isc} + \frac{(1-f)k_{et}(t)}{k_{nr} + k_{et}(t) + k_{sq}(B)})}{(f\Phi_{isc} + \frac{(1-f)k_{et}(t)}{k_{nr} + k_{et}(t) + k_{sq}(B) + k'_q(Az)})} \quad (2)$$

where, $f = E_t[t]$ which is the fraction of photons absorbed by *trans*-BPYE at 366 nm, $(1-\alpha)$ is the fraction of decay from $^3I^*$




Figure 1. Benzophenone effect on the *trans* → *cis* photoisomerization of BPyE, where the concentration of *trans*-BPyE is 2.5×10^{-3} M.




Figure 2. Stern-Volmer plots for direct and sensitized *trans* → *cis* photoisomerization of BPyE: line 1, in benzene at 366 nm ($[B]$ is 2.0×10^{-2} M (●)); line 2, in benzene at 366 nm ($[B]$ is 1.0×10^{-2} M (▲)); line 3, in benzene at 366 nm ($[B]$ is zero (■) or 1.0×10^{-3} M (○)) and at 435.8 nm ($[B]$ is zero (⊙)); line 4, in acetonitrile-water (v/v, 7:3) at 366 nm ($[B]$ is 5.0×10^{-2} M (□)); line 5, in benzene-dichloromethane (v/v, 1:1) at 366 nm ($[B]$ is zero (x) and at 435.8 nm ($[B]$ is zero or 1.2×10^{-2} M (⊗)); line 6, in acetonitrile-water (v/v, 7:3) at 366 nm ($[B]$ is zero (△) or 1.0×10^{-3} M (⊙)), where $[B]$ is the concentration of benzophenone.

to c_0 , which was found to be 0.43,^{1b} $[f]$ and $[B]$ are the concentration of *trans*-BPyE and benzophenone, respectively, k_{nr} and k_{sq} are the rate constants of nonradiative decay and self-quenching of benzophenone triplets in benzene which were reported to be $k_{nr} = 1.14 \times 10^4 \sim 1.0 \times 10^5 \text{ sec}^{-1}$ and $k_{sq} = 1.6 \times 10^5 \text{ M}^{-1}\text{sec}^{-1}$ by Schuster *et al.*^{2a} and $k_{nr} = 2.0 \times 10^4 \sim$

$5.0 \times 10^5 \text{ sec}^{-1}$ and $k_{sq} = 4.4 \times 10^5 \text{ M}^{-1}\text{sec}^{-1}$ by Singer *et al.*,^{2b} respectively, k_{et} is the rate constant of energy transfer from benzophenone triplets to *trans*-BPyE, k_q and k_q' are the quenching rate constants of *trans*-BPyE and benzophenone triplets by azulene, respectively, $1/k_{iso}$ (τ_T) is the lifetime of *trans*-BPyE triplets, Φ_{isc} is the intersystem crossing quantum yield of *trans*-BPyE, and Φ_{t+c}^0 is the *trans* → *cis* photoisomerization quantum yield of *trans*-BPyE in the absence of azulene.

We can derive the following equation (3) from the equation (1).

$$\frac{(1-f)(1-a)}{\Phi_{t+c}^{obs} - f(1-a)\Phi_{isc}} = 1 + \frac{k_{nr}}{k_{et}(t)} + \frac{k_{sq}}{k_{et}(t)} [B] \quad (3)$$

A plot of $(1-f)(1-a) (\Phi_{t+c}^{obs} - f(1-a)\Phi_{isc})^{-1}$ vs. $[B]$ gives a good linear relationship as shown in Figure 1, showing the intercept of 2.73 and the slope of $18.4 \text{ M}^{-1}\text{sec}^{-1}$, respectively, where the concentration of *trans*-BPyE is 2.5×10^{-3} M. The rate constant of the energy transfer from benzophenone triplets to *trans*-BPyE (k_{et}) must be in the range of $3.5 \sim 9.6 \times 10^6 \text{ M}^{-1}\text{sec}^{-1}$ and k_{nr} is in the range of $1.5 \sim 4.1 \times 10^4 \text{ sec}^{-1}$, respectively if we assume the k_{sq} value to fall in the range of $1.6 \sim 4.4 \times 10^5 \text{ M}^{-1}\text{sec}^{-1}$. Rate constant of the energy transfer for other related olefins, however, is $\sim 5 \times 10^9 \text{ M}^{-1}\text{sec}^{-1}$.

From the results of equation (3) and on the assumption that k_q and k_q' are the same as the diffusion controlled rate constant, we can modify the equation (2) into the equation (4).

$$\Phi_{t+c}^0 / \Phi_{t+c} = (1 + k_q[Az]/k_{iso}) \left(1 + \frac{(1-f)k_{et}(t)}{f\Phi_{isc}(k_{nr} + k_{et}(t) + k_{sq}[B])} \right) \quad (4)$$

The value of second term in equation (4) increases as $[B]$ increases because of the increase of $(1-f)$ value. The bigger Stern-Volmer constants, therefore, were obtained as $[B]$ increases as shown in Figure 2.

These phenomena can be observed when benzophenone triplets are quenched through self-quenching and triplet-triplet annihilations (TTA) of benzophenone or when the triplets of *trans*-BPyE are quenched by benzophenone. Since the triplets of *trans*-BPyE are not quenched by benzophenone when biacetyl is used as a photosensitizer at 435.8 nm, the process may be considered to be negligible. TTA mechanism seems not acceptable either since TTA occurs at high light intensities when long triplet lifetimes are involved. Since the lifetime of benzophenone triplets decreases as $[B]$ is increased, any involvement of TTA will diminish. Wolf *et al.*^{2b} also reported that *P*-type delayed fluorescence in benzophenone is negligible at concentrations < 0.05 M in benzene.

The low energy transfer rate constant of benzophenone triplets to *trans*-BPyE may be explained by proposing the formation of an AD^* complex or exciplex³ where benzophenone triplet is the donor (D) and the ground state *trans*-BPyE is the acceptor (A) and the rate constant for energy transfer as:^{3c}

$$k_{et}^{obs} = k_{sur}k_{et}/(k_{et} + k_{-sur}) \quad (5)$$

The observed energy transfer rate constant (k_{et}^{obs}) depends on the diffusion rate constant (k_{diff}) for diffusion of D^* and A and the energy transfer during the lifetime of the exciplex or complex. If we assume the rate of dissociation of the exciplex or complex to be very fast relative to energy transfer ($k_{-diff} > k_d$)⁴ as in the case of triplet-triplet energy transfer of valerophenone to 2,5-dimethyl-2,4-hexadiene,⁵ the observed rate constant is a composite of the equilibrium constant (k_{diff}/k_{-diff}) for exciplex or complex formation and the rate constant for energy transfer. In this case, the observed energy transfer rate constant is much smaller than the diffusion controlled rate constant. It indicates that the self-quenching of the benzophenone triplets must compete with energy transfer to *trans*-BPYE and the photoisomerization quantum yields decrease as the concentration of benzophenone, $[B]$, increases.

Acknowledgements. This investigation was supported by the Korea Science and Engineering Foundation. We are grateful to Prof. Jack Saltiel of Florida State University for

many helpful discussions.

References

- (a) P.-H. Bong, K. H. Chae, S. C. Shim, N. Nakashima, and K. Yoshihara, *J. Photochem.*, **31**, 223 (1985); (b) P.-H. Bong, H. J. Kim, K. H. Chae, S. C. Shim, N. Nakashima, and K. Yoshihara, *J. Am. Chem. Soc.*, **108**, 1006 (1986).
- (a) D. I. Schuster and T. M. Weii, *J. Am. Chem. Soc.*, **95**, 4091 (1973); (b) M. W. Wolf, K. D. Legg, R. E. Brown, L. A. Singer, and J. H. Parks, *ibid.*, **97**, 4490 (1975).
- (a) M. Smoluchowski, *Z. Phys. Chem.*, **92**, 129 (1917); (b) P. Debye, *Trans. Electrochem. Soc.*, **82**, 265 (1942); (c) T. R. Evans, *J. Am. Chem. Soc.*, **93**, 2081 (1970).
- J. B. Birks, "Organic Molecular Photophysics", Vol. 1, ed. J. B. Birks, New York, Wiley (1973), p. 403.
- P. J. Wagner and I. Kochevar, *J. Am. Chem. Soc.*, **90**, 2232 (1969).

Rhodium-Olefin Interaction: Double Bond Migration of *cis*-But-2-en-1,4-diol with Rhodium(I) Complexes

Chong Shik Chin* Byeongno Lee, and Kyungjoon Hong

Department of Chemistry, Sogang University, Seoul 121-742. Received January 11, 1990

The difference in affinities of rhodium(I) and iridium(I) toward a certain ligand is prominent, e.g., reactions of $\text{IrA}(\text{CO})(\text{PPh}_3)_2$ (A = monodentate anionic ligand such as halogens and ClO_4) with H_2 give stable $\text{Ir}(\text{H})_2\text{A}(\text{CO})(\text{PPh}_3)_2$ which are quantitatively isolated at room temperature¹ while $\text{Rh}(\text{H})_2\text{A}(\text{CO})(\text{PPh}_3)_2$ has never been isolated from the reactions of $\text{RhA}(\text{CO})(\text{PPh}_3)_2$ and H_2 . The M-P bond in $\text{IrCl}(\text{PPh}_3)_3$ is stronger than that in $\text{RhCl}(\text{PPh}_3)_3$; the latter is very effective catalyst for various organic reactions due to the significant dissociation of PPh_3 from rhodium in solution to provide a vacant site around rhodium for organic substrates² while the former is hardly employed as a catalyst mainly because the dissociation of PPh_3 from iridium is negligible.

It has been known that $\text{Ir}(\text{ClO}_4)(\text{CO})(\text{PPh}_3)_2$ attacks both hydroxyl groups of *cis*- $\text{HOCH}_2\text{CH}=\text{CHCH}_2\text{OH}$ (1) to give *trans*- $\text{CH}_3\text{CH}=\text{CHCHO}$ but leaves the olefinic group intact.³ The rhodium analogue $\text{Rh}(\text{ClO}_4)(\text{CO})(\text{PPh}_3)_2$ (2), however, behaves quite differently in the reaction with 1, which is described in this report along with related experimental results.

Rhodium-olefinic group interaction is apparent in the reactions of 2 and $[\text{Rh}(\text{CO})(\text{PPh}_3)_3]\text{ClO}_4$ (3) with 1 but no evidence for interaction between rhodium (in 2 and 3) and the hydroxyl groups of 1 has been observed (see below). 2-Hydroxytetrahydrofuran (4) is catalytically produced in the reactions of 1 with 2 or 3 (Table 1). Proton NMR spectral changes during the reactions clearly showed that a saturated

Table 1. Catalytic Production of 2-Hydroxytetrahydrofuran (4) in the Reactions of *cis*-But-2-en-1,4-diol (1) (2.0 mmole) with $\text{Rh}(\text{ClO}_4)(\text{CO})(\text{PPh}_3)_2$ (2) (0.1 mmole) and $[\text{Rh}(\text{CO})(\text{PPh}_3)_3]\text{ClO}_4$ (3) (0.1 mmole) at 45 °C under Nitrogen

| Catalyst | Solvent | Time, (hr) | Yield (%) |
|----------|--|------------|-----------|
| 2 | $\text{CDCl}_3(0.5 \text{ ml})$ | 32 | 100 |
| 3 | $\text{CDCl}_3(0.5 \text{ ml})$ | 24 | 100 |
| 2 | $\text{CDCl}_3(0.4 \text{ ml}) + \text{CD}_3\text{OD}(0.1 \text{ ml})$ | 5 | 100 |
| 3 | $\text{CDCl}_3(0.4 \text{ ml}) + \text{CD}_3\text{OD}(0.1 \text{ ml})$ | 3 | 100 |
| 2 | $\text{CDCl}_3(0.4 \text{ ml}) + \text{CD}_3\text{COCD}_3(0.1 \text{ ml})$ | 15 | 100 |
| 3 | $\text{CDCl}_3(0.4 \text{ ml}) + \text{CD}_3\text{COCD}_3(0.1 \text{ ml})$ | 10 | 100 |

aldehyde, presumably 4-hydroxybutanal (5), is produced and disappeared when a mixture of CD_3COCD_3 and CDCl_3 is used as a solvent (see Experimental and Table 1). No other products such as but-2-enal (which is produced in the reaction of the iridium complex $\text{Ir}(\text{ClO}_4)(\text{CO})(\text{PPh}_3)_2$ with 1³) has been found. Accordingly, the production of 4 seems to occur through the formation of 5 (equation 1). Formation of 5 is also strongly supported by the fact that complex 2 and 3 are very effective catalysts for the double bond migration of allylic alcohols to produce the corresponding saturated aldehydes⁴ and the double bond migration product 5 would readily