

Cald. for $C_{16}H_{16}O_2$: C, 80.00%, H, 6.67%. Found: C, 79.13%, H, 6.57%.

13. King, J. L.; Posner, B. A.; Mak, K. T.; Yang, N. C. *Tetrahedron Lett.* **1987**, 28, 3919.

14. Noh, T.; Kim, D. *Tetrahedron Lett.* **1996**, 37, 9329.

15. Elemental analysis of **4**: Cald. for $C_{16}H_{16}NO$: C, 81.45%, H, 4.98%, N, 6.33%. Found: C, 81.48%, H, 5.09%, N,

6.22%.

16. Sugioka, T.; Pat, C.; Sakurai, H. *Chem. Lett.* **1972**, 791.

17. Gleiter, R.; Schafer, W. *Acc. Chem. Res.* **1990**, 23, 369.

18. Wheland, G. W. *Resonance in organic chemistry*; Wiley: New York, 1955; p 85.

19. Schriver, G. W.; Gerson, D. J. *J. Am. Chem. Soc.* **1990**, 112, 4723.

The First Example of Cation Radical Induced Decarboxylation from Carbonates

Young Sook Park and Wang Keun Lee*

Department of Chemistry Education, Chonnam National University, Kwang-Ju 500-757, Korea

Received January 13, 1997

In the course of our investigation of cation radical induced oxidation of carbonates, we have discovered that thianthrene cation radical perchlorate (**1**, $Th^+ \cdot ClO_4^-$) readily oxidizes *tert*-butyl phenyl carbonate (**2**) in acetonitrile at room temperature. This is a novel reaction of carbonates, and offers the first example of cation radical induced decarboxylation. Carbonates, diester of carbonic acid, are known as stable organic compounds and as useful intermediates for a variety of industrial and synthetic applications. Alkyl aryl carbonates undergo the loss of carbon dioxide or are converted to diaryl carbonates at reasonable rates only if heated in the region 180-200 °C.¹ In contrast, the reaction (Scheme 1) of this very stable carbonate **2** with **1** at room temperature leads to rapid oxidative decarboxylation. Quantitative data are listed in Table 1. Product balances account for 88% of the *tert*-butyl groups which appear as cations, 91% of the phenoxy groups, and 95.1% of the cation radicals.


The major products are *N*-*tert*-Butylacetamide (**3**)² from *tert*-butyl cations, 5-(4-hydroxyphenyl)thianthreniumyl perchlorate (**4**)³ from phenoxy radicals, thianthrene (Th) and carbon dioxide. Thus, the intermediates which are trapped in the course of oxidative decomposition were *tert*-butyl cations and phenoxy radicals. *tert*-Butyl cations reacted with

the solvent acetonitrile to give a Ritter-type intermediate (Me_3CN^+CMe), which reacted with water during work-up, to give **3**. The simplest representation of the reaction of **1** with **2** and the formation of **3** and **4** is given in eq 1-5.



Table 1. Products of Reaction of Thianthrene Cation Radical Perchlorate with *tert*-Butyl Phenyl Carbonate in MeCN at Room Temperature^a

Reactants, mmol x 10 ³		Products (°), ^b mmol x 10 ³			
1	2	3	4	Th ^c	PhOH
100	50	44 (88)	45 (45)	49 (49)	0.5 (1)

^aReaction was carried out by adding acetonitrile by syringe to a stirred, septum-capped round-bottomed flask containing 1.00 mmol of **1** and 0.50 mmol of **2** under argon atmosphere. The color of **1** was discharged within 10 minutes, but stirring was continued overnight. Water (5 mL) was added. The solution was neutralized with NaHCO₃ and extracted repeatedly with CH₂Cl₂ (4 x 20 mL). Products were identified and quantified by GC, using the method of "standard addition" of authentic samples,² and by ¹H NMR and GC MS. GC analyses were made with a 2 m x 1/8 in. stainless steel column packed with 10% OV-101 on Chrom W. Reaction was run four times, and the averaged yields of products are given. Isolated yield, 100 x (mmol of product) / initial mmol of $Th^+ \cdot ClO_4^-$. This represents 90% of the phenoxy groups in the carbonate **2**. ^bA small amount (1.1%) of thianthrene 5-oxide (ThO) was also obtained.



Scheme 1.



Compound **3** was also obtained in 76% and 31% yields, respectively when *tert*-butyl *S*-(4,6-dimethylpyrimidine-2-yl)thiolcarbonate and *N*-(*tert*-butoxycarbonyl)glycine reacted with **1** in acetonitrile. Further separation of products proved not to be quantitative and their complete assay was not pursued.

Analogous studies on the aromatic cation radicals induced oxidative decomposition for azoalkanes have been reported.^{4,6} 1,1'-Azoadamantane,⁴ *meso*-azobis- α -phenylethane⁵ and cyclic azoalkanes⁶ were oxidized rapidly and quantitatively by **1** at room temperature to give nitrogen plus cation derived products instead of products formed by free radical reactions. While alkyl radicals formed from the first oxidation of azoalkanes in these reactions were again oxidized into alkyl cations, phenoxy radicals formed from the oxidative decomposition of carbonate (eq 2) were not oxidized to phenoxonium ion but trapped by **1** to give **4** in excellent yield (eq 5).

A question arose as to whether or not the reaction involved electron transfer and formation of radicals, which, being trapped by **1**, gave products 5-substituted thianthreniumyl perchlorates. Evidences for these steps were found in the reaction of **1** with dialkylmercurials,⁷ phenylmagnesium chloride,⁸ organotin,⁹ cyclohexanols,^{10,11} and 2,2'-azo-2-methyl-6-heptene.¹² Thus, without doubt, the fate of phenoxy radicals is primarily to be trapped by cation radicals and leads to the formation of **4**. An earlier report¹³ showed that **1** reacted with phenol to form **4** and the mole ratio of **1** to phenol was 2 : 1.

Although the major products are rationalized by eq 1-5, the origin of the 1.0% phenol remains uncertain. We can exclude the formation of phenoxonium ion because free phenoxonium ion could not be formed from mono- and disubstituted phenols.¹⁴ Moreover, neither amide nor hydroquinone from the reaction of phenoxonium ions with solvent or with water was found. Control reaction of perchloric acid and **2** in acetonitrile was carried out to know if the decomposition is catalyzed by acid that is formed in incompletely dried solvent instead of oxidative decomposition of the carbonate. Only a trace of phenol was obtained from the control experiment. This result indicates that not only the reaction is proceeding by electron transfer and not by acid catalysis but phenol may be formed by acid that is formed in incompletely dried solvent.

In conclusion, a thermally stable carbonate has been shown to undergo facile oxidative decarboxylation under mild conditions and both carbonations and free radicals were trapped by solvent and cation radical respectively. The new oxidative decomposition of *tert*-butyl phenyl carbonate further enlarges the chemistry of the versatile carboxylic linkage.

Acknowledgment. We would like to thank Chonnam National University for its financial assistance, and Dr. Kwang-Woo Jeong, of the Korea Basic Science Institute, Kwang-Ju Branch for NMR.

References

1. Shaikh, A. G.; Sivaram, S. *Chem. Rev.* **1996**, *96*, 951.
2. *N*-*tert*-Butylacetamide (**3**) was made by refluxing triphenylphosphine, bromotrichloromethane, acetic acid, and *n*-butylamine in THF and characterized by ¹H NMR (300 MHz, CDCl₃), δ 5.38 (s, 1H), 1.92 (s, 3H), 1.35 (s, 9H). Barstow, L. E.; Hruby, V. J. *J. Org. Chem.* **1971**, *36*, 1305.
3. [In order to recover 5-(4-hydroxyphenyl)thianthreniumyl perchlorate (**4**), the solid residue was dissolved in 10 mL of CH₂Cl₂ and to that solution was added 50 mL of ether. The mixture was placed in the refrigerator overnight, after which the precipitated salt was removed by filtration, weighed, and reprecipitated from CH₂Cl₂ with ether (1 : 5, v/v) and characterized by ¹H NMR (300 MHz, CD₃CN), δ 8.23 (dd, *J* = 7.62, 1.59 Hz, 2H), 7.95 (dd, *J* = 7.78, 1.51 Hz, 2H), 7.86 (dd, *J* = 7.50, 1.55 Hz, 2H), 7.77 (dd, *J* = 7.44, 1.68 Hz, 2H), 7.12 (d, *J* = 9.02 Hz, 2H), 6.93 (d, *J* = 9.60 Hz, 2H). The OH peak was not observed. Shin, S. R. Ph. D. Dissertation, Texas Tech University, 1991.
4. Bae, D. H.; Engel, P. S.; Hoque, A. K. M. M.; Keys, D. E.; Lee, W. K.; Shaw, R. W.; Shine, H. J. *J. Am. Chem. Soc.* **1985**, *107*, 2561.
5. Lee, W. K.; Chung, C. T. *Bull. Korean Chem. Soc.* **1992**, *13*, 694.
6. Engel, P. S.; Robertson, D. M.; Scholz, J. N.; Shine, H. J. *J. Org. Chem.* **1992**, *57*, 6178.
7. Sugiyama, K.; Shine, H. J. *J. Org. Chem.* **1983**, *48*, 143.
8. Lochynski, S.; Shine, H. J.; Soroka, M.; Venkatachalam, T. K. *J. Org. Chem.* **1990**, *55*, 2702.
9. Boduszek, B.; Shine, H. J.; Venkatachalam, T. K. *J. Org. Chem.* **1989**, *54*, 1616.
10. Lochynski, S.; Boduszek, B.; Shine, H. J. *J. Org. Chem.* **1991**, *56*, 914.
11. Shine, H. J.; Yueh, W. *Tetrahedron Lett.* **1992**, *33*, 6583.
12. Zhao, W.; Shine, H. J. *Tetrahedron Lett.* **1996**, *37*, 1749.
13. Chen, T.; Shine, H. J. *J. Org. Chem.* **1996**, *61*, 4716.
14. Kim, K.; Hull, V. J.; Shine, H. J. *J. Org. Chem.* **1974**, *39*, 2534.
15. Shin, S. R.; Shine, H. J. *J. Org. Chem.* **1992**, *57*, 2706.
16. Lee, M. L.; Yang, F. J.; Bartle, K. D. In *Open Tubular Column Gas Chromatography: Theory and Practice*; Wiley: New York, 1984.
17. The formation of ThO stems from the reaction of **1** with water which was present in the solvent or was added in work up. Vicil, E.; Maurey, M. M.; Cauquis, G. *Electrochimica Acta* **1984**, *29*, 1009 and references therein. Shine, H. J.; Murata, Y. *J. Org. Chem.* **1969**, *34*, 3368.