

Three New Species of Poecilosclerid Sponge from Korea

Chung Ja Sim* and Kyung Jin Lee

Department of Biology, Hannam University, Taejeon 300-791, Korea

Key Words:

Demospongiae
Poecilosclerida
New species
Korea

Three species of Poecilosclerid sponge (class Demospongiae, subclass Ceractinomorpha) are newly described from the waters around Cheju Island, Keomun Island and Keoje Island. They are *Antho (Plocamia) bakusi*, *Clathria (Clathria) koreana* and *Oxymycale rhoi*.

Ninety percent of living sponges belong to the class Demospongiae (Hajdu, 1995). The order Poecilosclerida (subclass Ceractinomorpha) is the largest and most diverse among demosponge orders (Bergquist and Fromont, 1988). The shape varies from arborescent to massive or even encrusting, occurring in all seas from intertidal habitats to deep seas. Bakus (1966) reported 23 species of this order from the San Juan Archipelago, Washington, USA. Sim and Bakus (1986) reported 15 species from Santa Catalina Island, California. Bergquist and Fromont (1988) reported 108 species including 38 new species from New Zealand. Hooper (1996) described 459 species, including 52 new species in a phylogenetic revision of the Poecilosclerid Microcionidae. Among them, 148 species were recorded from the Australian fauna. Sixty-three species of Poecilosclerid sponges were recorded from Japan (Hoshino, 1981), and 55 species from Korea (Rho & Sim 1979, Sim 1981, Sim & Kim 1988, Sim & Byeon 1989). The materials examined in the present study were collected by SCUBA and fishing-nets from around Cheju Island, Keomun Island and Keoje Island, south of Korean mainland. Sponge identification requires examination of various characters, which may include shape, color, consistency, texture, skeletal arrangement, and spicules. For skeletal arrangement, thin free-hand sections were made with a surgical blade from specimens hardened in alcohol. Spicules were examined by dissolving a piece of sponge in sodium hypochlorite (bleach). For microsclere morphology, scanning electron microscope preparations were made by drying spicules on an AKASHI ISI- SS40 (SEM) at the Department of Biology, Hannam University. SEM study of spicules followed the procedure of Rützler (1978). Type specimens of three new species are stored at the Natural History Museum (NHM), Hannam University, Taejeon, Korea.

Systematic Accounts

Class Demospongia Sollas, 1885
Subclass Ceractinomorpha Lévi, 1953
Order Poecilosclerida Topsent, 1928
Family Microcionidae Carter, 1875

Antho (Plocamia) bakusi n. sp.
(Fig. 1. A-H)

Type specimens: Holotype (Por. 23, NHM, Hannam Univ.), two paratypes (Por. 23-1, Por. 23-2, Dept. of Biology, Hannam Univ.), Daesambudo (Keomun Island) on 27 Nov. 1995, 25 m (SCUBA).

Description: Specimens always encrusting on sponge *Jaspis wondoensis* and measure up to 9×4.5×1 cm.

Surface rough with irregularly arranged tubercles. Oscules measure up to 1 mm in diameter. Texture firm and colour in life orange-red.

Ectosome. Thick styles arranged at right angles to body surface. Acanthostrongyles and acanthostyles arranged regularly. Isochelas, toxas and thin styles found in the cortex.

Choanosome. Acanthostrongyles irregularly arranged. Thick styles, isochelas and toxas rare.

Spicules. Megascleres

Styles (thick) ----- 225-650 × 10-25 μm
Styles (thin) ----- 160-445 × 4-6 μm
Acanthostyles ----- 110-185 × 5-10 μm
Acanthostrongyles ----- 115-145 × 9-15 μm

Microscleres

Toxas (large) ----- 107-207 μm
Toxas (medium) ----- 55-95 μm
Toxas (small) ----- 17.5-48 μm
Isochelas ----- 17.5-22.5 μm

Etymology: The specific name *bakusi* is named after Dr. Gerald J. Bakus, who is a professor at Department of Biological Science, University of Southern California, a marine ecologist and a sponge taxonomist.

* To whom correspondence should be addressed.
Tel: 82-42-629-7485, Fax: 82-42-629-7444




Fig. 1. *Antho (Procamia) bakusi* n. sp. A, Top-side view. B, Back-side view. C, Megascleres: a, thick style, b, thin style, c, acanthostrongyle. D, Skeletal structure. E-F, Microscleres. E, a, large toxas, b, middle toxas, c, small toxas, F, Isochela. G, Head of thin style. H, a, spined thick style, b, acanthostyle, c, isochela. Scale bars=1.35 μ m (G), 2.59 μ m (F), 26.2 μ m (E), 150 μ m (C, D), and 1 cm (A, B).




Fig. 2. *Clathria* (*Clathria*) *koreana* n. sp. A, Side view. B, Large toxa. C, Style. D, Skeletal structure. E, a, toronte, b, acanthostyle. F, a, head of acanthostyle, b, the other end. G, Small toxa. H, Isochela. Scale bars=10 μ m (F, H), 13.3 μ m (G), 100 μ m (E), 200 μ m (C, D), 230 μ m (B), and 2 cm (A).

Three New Poecilosclerid Sponge from Korea

Table 1. Comparison of spicule size in *Anto (Plocamia) bakusi* n. sp. with *P. illgi* and *P. burtoni*

Spicules	Size of spicule (µm)		
	<i>P. illgi</i>	<i>P. burtoni</i>	<i>Anto (Plocamia) bakusi</i> n. sp.
Megascleres (µm)			
Styles (thick)	254-333 × 18-23	100-350 × 5-7	225-650 × 10-25
Styles (thin)	160-252 × 4-7	190-260 × 2-3	160-445 × 4-6
Acanthostyles	179-219 × 10-19	75-100 × 7	110-185 × 5-10
Acanthostrongyles	120-159 × 10-17	75-100 × 7	115-145 × 9-15
Microscleres (µm)			
Toxas (large)	55	85-130 × 3	107-207
Toxas (medium)	40	75 × 1-2	55-95
Toxas (small)	29	15-40 × 0.5	17.5-48
Isochelas	19-25	13-14	17.5-22.5

Remark: This new species appears to be closely related to *Procambia illgi* Bakus, 1966 and *P. burtoni* Lévi, 1952. Large styles and toxas of *P. illgi* and *P. burtoni* are shorter than those of this species (Table 1).

Clathria (Clathria) koreana n. sp.
(Fig. 2. A-H)

Type specimens: Holotype (Por. 24, NHM, Hannam Univ.), Sogwipo (Cheju Island), on 14 Jul. 1982 (fishing-net).

Description: Shape flabellate with flared margins. Sponge 200 mm high with a short, firm stalk (65 mm long, 8 mm diameter), 150 mm wide, with 3-4 mm thick lamellae. Colour dark brown in alcohol. Surface velvety, and texture firm. Oscules up to 1 mm diameter.

Ectosome. Thin membrane, single principal styles protrude through surface. Subectosomal auxiliary styles tangential or paratangential and scattered over surface.

Choanosome. A reticulate skeletal architecture present with large multispicular tracts of principal styles ascending to surface, and with transverse spicules. Spicules bound together at ends by abundant spongin.

Spicules. Megascleres	
Choanosomal principal styles	----- 400-600 × 20-25 µm
Subectosomal auxiliary torontes with blunt hastate microspined point	----- 230-390 × 2-4.5 µm
Acanthostyles (with very small spines)	----- 115-145 × 8-10 µm
Microscleres	
Wing-shaped toxas (large)	----- 300-1030 µm
(small)	----- 43-78 µm
Palmate isochelas	----- 15-20 µm

Etymology: The specific name *koreana* is named for Republic of Korea.

Remark: This new species has spicules similar to those of *Axociella bitoxifera* Koltun, 1970 except that it has large spicules and lacks acanthostyles. This genus was transferred from *Axociella* to *Clathria (Thalysias)* by Hooper's (1996) definition of auxiliary spicules. The large toxas and acanthostyles with small spines in this species are different from those of other species.

Family Mycalidae Lundbeck, 1905

Oxymycale rhoi n. sp.
(Fig. 3. A-G)

Type specimens: Holotype (Por. 22, NHM, Hannam Univ.), Memuldo (Keoje Island), on 9 Jul. 1996 (fishing-net) 40 m depth. Four Species Paratypes (Por. 22-1, Por. 22-2, Por. 22-3, Por. 22-4, Dept. of Biology, Hannam Univ.), collected with holotype

Description: Specimens irregularly cushion-shaped with a size up to 14.5 × 8 × 5 cm. Surface even to conulose with several oscula of 0.8-4 mm diameter. Texture soft, compressible, and fragile like wet bread, and sponge easily torn. Colour in life dirty yellowish grey.

Ectosome. Skeleton composed of oxeas in confusion arranged parallel to surface with anisochelas (rosettes of anisochelas do not occur, only loose groups of 3-5 anisochelas).

Choanosome. Skeleton has long ascending tracts of oxeas, 50-1,000 µm in diameter, which form subectosomal tufts. Deeper part of the choanosome has much spongin mixed with oxeas.

Spicules. Megascleres	
Oxeas	----- 500-660 × 8-18 µm
Microscleres	
Anisochela	----- 90-110 µm
Anisochela 2	----- 50-80 µm
Anisochela 3	----- 30-37 µm
Anisochela 4	----- 80-90 µm
Early stage of anisochela 4	----- 80-90 µm
Sigmas (large)	----- 55-70 µm
(small)	----- 17.5 µm
Raphides	----- 75-87 µm

Remark: This new species is similar in spicules to *Oxymycale koreana* Sim (1982), but the new species has three size categories of anisochelas whereas *O. koreana* has only two size categories. The arrangement of anisochelas is not in rosettes. The specimen of *O. koreana* is club-shaped and hard in texture.

Etymology: This specific name *rhoi* is named for Dr. Boon Jo Rho, who has contributed greatly to sponge taxonomy since 1996.




Fig. 3. *Oxymycale rhoi* n. sp. A, Top-side view. B, Back-side view. C, Oxea. D, Skeletal structure of fiber. E-G, Microscleres; E, a. anisochela 3, b. early stage of anisochela 4. F, a. large sigma, b. small sigma. G, a. anisochela 1, b. anisochela 2, c. anisochela 3, d. anisochela 4, e. raphid. Scale bars=5.08 μ m (E. a), 10.2 μ m (F), 20.4 μ m (G), 34.4 μ m (E. b), 200 μ m (C), 400 μ m (D), and 3 cm (A, B).

Acknowledgements

This work was supported in part by the Basic Science Research Institute Program, Korean Ministry of Education through Research Fund (BSRI-95-4428).

We thank Dr. G. J. Bakus, Department of Biological Science, University of Southern California for his advice and review of the manuscript.

References

- Bakus GJ (1966) Marine poeciloscleridan sponges of the San Juan Archipelago, Washington. *J Zool* 149: 415-531.
- Bergquist PR and Fromont JP (1988) The marine fauna of New Zealand: Porifera, Demospongiae. Part 4. Poecilosclerida. *N Z Oceanogr Inst Mem* 96: 1-197.
- Hajdu E (1995) Macroevolutionary patterns within the demosponge Order Poecilosclerida. Ph. D. Thesis, University of Amsterdam, pp 1-136.
- Hooper JNA (1996) Revision of Microcionidae (Porifera: Poecilosclerida: Demospongiae), with description of Australian species. *Mem Queensl Mus* 40: 1-626.
- Hoshino T (1981) Shallow-water demosponges of western Japan. I. *J Sci Hiroshima Univ Ser B Div 1 Zool* 29: 107-109.
- Koltun VM (1970) Sponge fauna of the northwestern Pacific from the shallows to the hadal depths. Communication 1. In: Bogorov VG (ed), Fauna of the Kurile-Damchatka Trench and Its Environment, Akademiya Nauk SSSR. Moskva, pp 177-223.
- Lévi C (1952) Spongiaires de la côte du Sénégal. *Bull l'Insti Fondam d'Afr Noire* 14: 34-59.
- Rho BJ and Sim CJ (1979) A taxonomic study on the Korean sponges. 1. Poecilosclerina. *J Korean Res Inst Bet Liv* 23: 61-67.
- Rützler K (1978) Sponges in coral reefs. In: Stoddart DR and Johannes RE (eds), Coral Reefs: Research Methods. *Monogr Oceanogr Neth UNESCO* 5: 299-313.
- Sim CJ (1981) A systematic study on the marine sponges in Korea. 1. Ceractinomorpha and Tetractinomorpha. *Soong Jun Univ Essays Pap* 11: 83-105.
- Sim CJ (1982) A systematic study on the marine sponges from Jeju Island. *Soong Jun Univ Essays Pap* 12: 187-210.
- Sim CJ and Bakus GJ (1986) Marine sponges of Santa Catalina Island, California. *Allan Hancock Found Univ South Calif* 5: 1-23.
- Sim CJ and Byeon HS (1989) A systematic study on the marine sponges in Korea. 9. Ceractinomorpha. *Korean J Syst Zool* 5: 33-57.
- Sim CJ and Kim MH (1988) A systematic study on the marine sponges in Korea. 7. Demospongia and Hexactinellida. *Korean J Syst Zool* 4: 21-42.

[Received December 12, 1997; accepted January 14, 1998]