

References

1. Imuta, M.; Itani, H.; Nishi, K.; Ona, H.; Kimura, Y. *Bioorg. Med. Chem. Lett.* **1993**, *3*, 2199.
2. Freshly prepared 2-hydroxymethylcarbapenem **1** should be used for MnO₂ oxidation due to its unstability at room temperature.
3. Dondoni, A.; Fantin, G.; Fogagnolo, M.; Medici, A. *Angew. Chem., Int. Ed. Engl.* **1986**, *25*, 835.
4. This compound is available from Takasago Chemical Co., Ltd, Japan.
5. Angelastro, M. R.; Peet, N. P.; Bey, P. *J. Org. Chem.* **1989**, *54*, 3913.
6. Schmitt, S. M.; Saltzmann, T. N.; Shih, D. H.; Christensen, B. G. *J. Antibiot.* **1988**, *41*, 780.
7. Physical data: **2**; ¹H NMR (200 MHz, CDCl₃) δ 0.05 (6H, s), 0.85 (9H, s), 1.20 (6H, m), 3.35 (1H, dd, *J*=5.1, 3.4 Hz), 3.46 (1H, m), 4.24 (1H, m), 4.29 (1H, dd, *J*=10.3, 3.4 Hz), 4.79 (2H, m), 5.28 (1H, dd, *J*=10.3, 1.4 Hz), 5.43 (1H, dd, *J*=17.2, 1.4 Hz), 5.88-5.97 (1H, m), 10.32 (1H, s). ¹³C NMR (50 MHz, CDCl₃) δ -5.1, -4.2, 16.2, 17.9, 22.1, 25.6, 37.9, 56.3, 60.3, 65.3, 66.7, 119.2, 130.5, 141.9, 159.1, 172.1, 188.8.
3; mp 58-61 °C IR (CDCl₃) cm⁻¹ 3403, 2957, 1779, 1465, 1386, 1275. ¹H NMR (200 MHz, CDCl₃) δ 0.09 (3H, s), 0.10 (3H, s), 0.89 (9H, s), 1.29 (3H, d, *J*=6.2 Hz), 1.30 (3H, d, *J*=7.3 Hz), 3.22 (1H, dd, *J*=6.2, 2.8 Hz), 4.07 (1H, m), 4.27 (2H, m), 4.82 (2H, m), 5.29 (1H, dd, *J*=10.5, 1.4 Hz), 5.46 (1H, dd, *J*=17.2, 1.4 Hz), 5.93-6.02 (1H, m), 7.51 (1H, d, *J*=3.0 Hz), 7.94 (1H, d, *J*=3.0 Hz). ¹³C NMR (75 MHz, CDCl₃) δ -4.9, -4.1, 17.0, 18.0, 22.5, 25.7, 43.5, 55.8, 59.9, 66.2, 118.7, 122.9, 127.3, 131.2, 142.1, 142.8, 158.4, 161.1, 172.8. HR EI-MS *m/z* calcd for C₂₂H₂N₂O₄SSi 448.1857, Found 448.1877.
5; mp 93-95 °C IR (CDCl₃) cm⁻¹ 3157, 2968, 1760, 1650, 1469, 1386, 1253. ¹H NMR (200 MHz, CDCl₃) δ 0.04 (3H, s), 0.06 (3H, s), 0.83 (9H, s), 1.14 (3H, d, *J*=6.3 Hz), 1.16 (3H, d, *J*=7.0 Hz), 2.97 (1H, dd, *J*=4.7, 2.2 Hz), 3.12 (1H, m), 3.19 (3H, s), 3.70 (3H, s), 3.84 (1H, dd, *J*=4.7, 2.2 Hz), 4.18 (1H, m), 6.06 (1H, s). ¹³C NMR (75 MHz, CDCl₃) δ -5.1, -4.4, 12.4, 17.8, 22.2, 25.6, 31.9, 37.7, 52.0, 61.3, 65.1, 168.4, 175.1. FAB-MS (*m/z*) 345 (M⁺)
6a; mp 103-106 °C IR (CDCl₃) cm⁻¹; 3098, 2971, 1761, 1673, 1477, 1387, 1258. ¹H NMR (300 MHz, CDCl₃) δ 0.04 (3H, s), 0.05 (3H, s), 0.86 (9H, s), 1.11 (3H, d, *J*=6.3 Hz), 1.32 (3H, d, *J*=7.0 Hz), 3.07 (1H, dd, *J*=4.0, 2.2 Hz), 4.01 (1H, dd, *J*=5.1, 2.2 Hz), 4.16 (2H, m), 6.04 (1H, br), 7.74 (1H, d, *J*=3.0 Hz), 8.03 (1H, d, *J*=3.0 Hz). ¹³C NMR (75 MHz, CDCl₃) δ -5.1, -4.4, 12.6, 17.8, 22.3, 25.7, 43.7, 51.6, 61.6, 64.9, 127.1, 144.8, 166.4, 168.2, 195.5. HR EI-MS *m/z* calcd for C₁₇H₂₈N₂O₃SSi 368.1589, Found 368.1570.
8. Dondoni, A.; Perrone, D.; Merino, P. *J. Org. Chem.* **1995**, *60*, 8074.

Formation of *o*-/*p*-Quinomethanes and *p*-Quinodimethanes from the Photoaddition of Diphenylacetylene to *o*-Quinones[†]

Ae Rhan Kim, Yoon Jung Mah, and Sung Sik Kim*


Department of Chemistry, Chonbuk National University,
Chonju 561-756, Korea

Quinones are an important class of compounds as quinone dye-stuffs in industry or dehydrating agents in organic synthesis in addition to a vital role in biological systems. Due to their various spectroscopic properties, the photochemistry of quinones has been a subject of interest in many areas.^{1,2} Our interest in diverse reactivity of excited quinones has promoted us to investigate the type of the photoadducts of quinones.³⁻⁵ Bryce-Smith *et al.* have reported that tetrachloro-1,2-benzoquinone **1a** reacts photochemically with diphenylacetylene (DPA) **2** to give dioxenes.⁶ Photoreaction of **1a** and **2** in acetone or acetonitrile at >400 nm gave 1:2 adduct as 1,4-dioxene. We found that, when irradiated with 300 nm UV light, *o*-quinones add to **2** to give two isomeric *o*-quinomethanes, *i.e.*, **4** and **5**, **8** and **9**, and **12** and **13**, *via* spiro-oxetene intermediates, like **3**.⁷

Recently, we found an interesting fact that irradiation (300 nm) of tetrahalo-1,2-benzoquinones **1** and DPA **2** in dichloromethane gave new type of *p*-quinomethanes **6**, as well as two isomeric *o*-quinomethanes, **4** and **5**, as shown in Scheme 1. Preparative photochemical reactions were conducted in a photoreactor composed of a water-cooled system and a Pyrex reaction vessel with 300 nm UV lamps (Rayonet Photochemical Reactor, Model RPR-208), after purging with nitrogen gas (purity; 99.9%) for 30 min. The photoproducts were isolated by flash column chromatography (silica gel, 230-400 mesh, Merck Co.) using *n*-hexane and ethyl acetate as the eluents (from 97:3 to 9:1, v/v).

Irradiation of a dichloromethane solution (100 mL) of tetrachloro-1,2-benzoquinone **1a** (246 mg, 1.0 mmol) and DPA **2** (178 mg, 1.0 mmol) with 300 nm UV light for 24 h afforded not only two isomeric *o*-quinomethanes, **4a** (26%) and **5a** (28%), *via* unstable spiro-oxetene **3a**, but also a novel *p*-quinomethane **6a** (17%).⁸ The absorption peaks for

[†]This paper is dedicated to Professor Sang Chul Shim on the occasion of his 60th birthday.





Scheme 1

the four protons of quinone moiety in **6a** were observed at δ 7.57, 7.49, 7.10, and 6.91 as doublets ($J=8.0$ Hz) each in ^1H NMR spectrum (CDCl_3).

Irradiation of tetrabromo-1,2-benzoquinone **1b** (424 mg, 1.0 mmol) and DPA **2** (178 mg, 1.0 mmol) in dichloromethane (100 mL) with 300 nm UV light for 24 h yielded same types of quinomethanes, **4b**, **5b**, and **6b**, in 29%, 30%, and 23% yields, respectively.⁹ Two dimensional ^1H - ^1H correlation spectrum (CDCl_3) of **6b** shows that the proton peaks of quinone moiety at δ 7.09 and 6.90 correspond to the peaks at δ 7.56 and 7.48, respectively.

It has been known that phenanthrenequinone **7** adds photochemically to alkynes to yield 1,4-dioxene, quinomethane, or 1,3-dioxole.^{10,11} Irradiation of phenanthrenequinone **7** (208 mg, 1.0 mmol) and DPA **2** (89 mg, 0.5 mmol) in dichloromethane (200 mL) with 300 nm UV light for 48 h afforded a *p*-quinomethane **10** (26%), as well as two isomeric *o*-quinomethanes, **8** (27%) and **9** (29%).¹² The formation of **8** (or **9**) and **10** was confirmed from the mass spectra (EI), in which the molecular ions were found at m/e 386 and 402, respectively. Two dimensional ^1H - ^1H correlation spectrum (CDCl_3) of **10** showed that the four absorption peaks of quinone moiety at δ 7.57 and 7.47 correspond to the peaks at δ 7.09 and 6.90, respectively. Two carbon peaks of **10** were observed at δ 207.0 and 205.5 in ^{13}C NMR spectrum (CDCl_3).

Irradiation of acenaphthenequinone **11** (91 mg, 0.5 mmol) and DPA **2** (89 mg, 0.5 mmol) in dichloromethane (200 mL) with 300 nm UV light for 48 h gave a *p*-quinomethane **14** (19%), as well as two isomeric *o*-quinomethanes, **12** (24%) and **13** (26%).¹³ The exact structure of *o*-quinomethanes, such as **8**, **9**, **12**, and **13**, was also confirmed by irradiating in dichloromethane in the presence of molecular oxygen, in




Scheme 2

which two benzene rings in the molecules was fused to give the corresponding oxidative photocyclization products. Purified **8** (or **12**) was irradiated with 300 nm UV light in dichloromethane to give **10** (or **14**). Irradiation of the other isomer **9** (or **13**) also gave rise to **10** (or **14**). Two dimensional ^1H - ^1H correlation spectrum (CDCl_3) of **14** showed that the peaks at δ 7.58 and 7.40 correspond to the peaks at δ 7.06 and 6.80, respectively. Mass spectrum (EI) showed the molecular ion peak at m/e 376.

In order to elucidate the formation mechanism of **6a**, a dichloromethane solution of **4a** was purged with molecular oxygen for 30 min and irradiated in dichloromethane to give **6a**. Irradiation of **4** after degassing with freeze-pump-thaw method did not give **6**.

These results imply that the formation of *p*-quinomethane **6** proceeds *via* triplet state of the starting material **4**. It may be considered that irradiation of **4** leads to the formation of radical intermediate **I**, as shown in Scheme 2. The resonance-stabilized radical **I** can be alive until a molecular oxygen is added to give hydroperoxide **II**. The hydroperoxide **II** may be considered to decompose to yield the final product **6** as a novel *p*-quinomethane.



Irradiation of **1** and **2** in high concentration also gave rise to Chichibabin hydrocarbons **15** as a dimeric *p*-quinodimethane. The FAB mass spectrum of **15** (X=Cl) showed the molecular ion peak at *m/e* 845 (M+H). ¹H NMR spectrum (CDCl₃) showed hydroxyl protons and aromatic protons at δ 11.40 (s) and 7.99-7.59. The proton signals of *p*-quinodimethane moiety were observed at δ 7.57, 7.49, 7.10, and 6.91, in which all signals appeared as doublet (*J*=8.0 Hz) each. The same type of dimeric compounds, *i.e.*, **16** and **17**, were also observed and isolated from the photoreaction of **2** and **7** (or **11**), as shown below.^{14,15}

Interestingly, one-way *E*-to-*Z* isomerization was observed between the two *o*-quinodimethanes, **4** and **5**.⁷ Irradiation of 20 mg (4.7×10^{-5} mol) of *E*-isomer **4a** in 15 mL of dichloromethane with 300 nm UV light for 24 h gave *Z*-isomer **5a** (23%), and also **6a** (16%), whereas irradiation of **5a** did not give **4a**. The similar results were obtained in the case of **8** and **12**.

In conclusion, we have shown here that two isomeric *o*-quinomethanes are produced when *o*-quinones and DPA **2** are irradiated with 300 nm UV light, instead of visible light (>400 nm). It is also interesting to note that novel *p*-quinomethanes and *p*-quinodimethanes may be formed via the initially produced *o*-quinomethanes.

Extension of the chemical properties of various *o*-quinones and their photoproducts will be investigated.

Acknowledgment. We appreciate the financial support of this work by the Organic Chemistry Research Center (OCRC) sponsored by the Korea Science and Engineering Foundation (KOSEF). This paper was also supported (in part) by the Basic Science Research Institute Program, Ministry of Education, 1998, Project No. BSRI-98-3431.

References

- Laird, T. In *Comprehensive Organic Chemistry*; Stoddart, J. F. Ed.; Pergamon Press: 1979; Vol. 1, p 1213.
- Maruyama, K.; Osuka, A. In *The Chemistry of the Quinonoid Compounds*; Patai, S.; Rappoport, Z. Eds.; John Wiley & Sons: Inc.; New York, 1988; Part 1, Chapter 13, p 759.
- Kim, S. S.; Yu, Y. H.; Shim, S. C.; Cho, I. Ho. *Tetrahedron Lett.* **1994**, 35(48), 9039-9042.
- Kim, S. S.; Kim, A. R.; Oh, K. J.; Yoo, D. J.; Shim, S. C. *Chemistry Lett.* **1995**, 9, 787-788.
- Kim, A. R.; Kim, S. S.; Yoo, D. J.; Shim, S. C. *Bull. Korean Chem. Soc.* **1997**, 18(6), 665-667.
- Bryce-Smith, D.; Gilbert, A. *Chem. Commun.* **1968**, 24, 1702.
- Kim, A. R.; Mah, Y. J.; Shim, S. C.; Kim, S. S. *J. Photochem. Photobiol.* **1997**, 4(2), 49.
- 6a**: 400 MHz ¹H NMR (CDCl₃), δ 11.4 (1H, s, hydroxy), 7.99-7.59 (5H, aromatic), 7.57 (1H, d, *J*=8.0 Hz), 7.49 (1H, d, *J*=8.0 Hz), 7.10 (1H, d, *J*=8.0 Hz), 6.91 (1H, d, *J*=8.0 Hz); IR (KBr), 3369, 3064, 1732, 1682, 1030, 763 cm⁻¹; UV (MeOH), 334, 301, 279, 253, 245 nm; Mass (EI), *m/e* 438 (M), 333 (M-105), 105.
- 6b**: 400 MHz ¹H NMR (CDCl₃), δ 11.4 (1H, s, hydroxy), 7.99-7.57 (5H, aromatic), 7.56 (1H, d, *J*=8.0 Hz), 7.48 (1H, d, *J*=8.0 Hz), 7.09 (1H, d, *J*=8.0 Hz), 6.90 (1H, d, *J*=8.0 Hz); IR (KBr), 3382, 3061, 1734, 1682, 1038, 763 cm⁻¹; UV (MeOH), 336, 328, 283, 256, 245 nm; Mass (EI), *m/e* 614 (M), 105.
- Bos, H. J. T.; Polman, H.; van Montfort, P. F. E. *J. Chem. Soc., Chem. Commun.* **1973**, 188.
- Mosterd, A.; de Noten, L. J.; Bos, H. J. T. *Recl. Trav. Chim. Pays-Bas* **1977**, 96, 16.
- 10**: 400 MHz ¹H NMR (CDCl₃), δ 11.4 (1H, s, hydroxy), 8.80-7.35 (13H, aromatic), 7.57 (1H, d, *J*=8.0 Hz), 7.47 (d, *J*=8.0 Hz), 7.09 (1H, d, *J*=8.0 Hz), 6.90 (1H, d, *J*=8.0 Hz); IR (KBr), 3440, 3065, 1742, 1671, 1093, 757 cm⁻¹; ¹³C NMR (CDCl₃), 207.0, 205.5, 133.8, 130.2, 129.8, 129.4, 129.1, 129.0, 128.6, 128.5, 128.2, 127.5, 127.0, 126.3, 123.1, 122.9; UV (MeOH), 338, 301, 252 nm; Mass (EI), *m/e* 402 (M), 105.
- 14**: 400 MHz ¹H NMR (CDCl₃), δ 11.4 (1H, s, hydroxy), 8.31-7.23 (13H, aromatic), 7.58 (1H, d, *J*=8.0 Hz), 7.40 (d, *J*=8.0 Hz), 7.06 (1H, d, *J*=8.0 Hz), 6.80 (1H, d, *J*=8.0 Hz); IR (KBr), 3440, 3059, 1734, 1687, 1014, 776 cm⁻¹; Mass (EI), *m/e* 376 (M), 271 (M-105), 105.
- 16**: 400 MHz ¹H NMR (CDCl₃), δ 11.5 (1H, s, hydroxy), 8.80-7.35 (26H, aromatic), 7.57 (2H, d, *J*=8.0 Hz), 7.47 (d, *J*=8.0 Hz), 7.09 (1H, d, *J*=8.0 Hz), 6.90 (1H, d, *J*=8.0 Hz); IR (KBr), 3440, 3065, 1742, 756 cm⁻¹; Mass (FAB), *m/e* 773 (M+1).
- 17**: 400 MHz ¹H NMR (CDCl₃), δ 11.5 (1H, s, hydroxy), 8.31-7.13 (22H, aromatic), 7.58 (2H, d, *J*=8.0 Hz), 7.40 (2H, d, *J*=8.0 Hz), 7.06 (2H, d, *J*=8.0 Hz), 6.90 (2H, d, *J*=8.0 Hz); IR (KBr), 3440, 3066, 1743, 757 cm⁻¹; Mass (FAB), *m/e* 721 (M+1).