

ON M-CONTINUITY

WON KEUN MIN AND HONG SOON CHANG

ABSTRACT. In this paper, we introduce a new class of sets, called m -sets, and the notion of m -continuity. In particular, m -sets and m -continuity are used to extend known results for α -continuity and semi-continuity and precontinuity.

1. Introduction

Let X, Y and Z be topological spaces on which no separation axioms are assumed unless explicitly stated. Let S be a subset of X . The closure (resp. interior, boundary) of S will be denoted by S^- (resp. $S^0, b(S)$). A subset S of X is called semi-open set[1] (resp. preopen set[2], α -set[3]) if $S \subset S^{0-}$ (resp. $S \subset S^{-0}, S \subset S^{0-0}$). The complement of a semi-open set (resp. preopen set, α -set) is called semi-closed set (resp. preclosed set, α -closed set). The family of all semi-open sets (resp. preopen sets, α -sets) in X will be denoted by $SO(X)$ (resp. $PO(X), \alpha(X)$). A function $f : X \rightarrow Y$ is called semi-continuous[1] (resp. precontinuous[2], α -continuous [4]) if $f^{-1}(V) \in SO(X)$ (resp. $f^{-1}(V) \in PO(X), f^{-1}(V) \in \alpha(X)$ for each open set V of Y).

A subclass $\tau^* \subset P(X)$ is called a supratopology on X if $X \in \tau^*$ and τ^* is closed under arbitrary union. (X, τ^*) is called a supratopological space. The members of τ^* are called supraopen sets[5]. Let (X, τ) be a topological space and τ^* be a supratopology on X . We call τ^* a supratopology associated with τ if $\tau \subset \tau^*$. Let (X, τ^*) be a supratopological space and (Y, μ) be a topological space. A function $f : X \rightarrow Y$ is an S -continuous function if the inverse image of each open set in Y is a supraopen set in X [5]. Let (X, τ^*) and (Y, μ^*) be supratopological spaces. A function $f : X \rightarrow Y$ is an S^* -continuous

Received June 24, 1998.

1991 Mathematics Subject Classification: 54C08, 54A10.

Key words and phrases: m -sets, τ_m , m -topology, m -continuous.

function if the inverse image of each supraopen set in Y is a supraopen set in $X[5]$.

2. m -sets induced by a supratopology

DEFINITION 2.1. Let (X, τ^*) be a supratopological space. A subset A of X is called an m -set with τ^* if $A \cap T \in \tau^*$ for all $T \in \tau^*$.

The class of all m -sets with τ^* will be denoted by τ_m .

EXAMPLE 2.2. Let $X = \{a, b, c, d\}$ and $\tau^* = \{\emptyset, X, \{a\}, \{b, c\}, \{b, d\}, \{a, b, c\}, \{b, c, d\}, \{a, b, d\}\}$. Then $\tau_m = \{\emptyset, X, \{a\}, \{b, c, d\}\}$.

REMARK. Let (X, τ) be a topological space. Since $SO(X)$ is closed with respect to arbitrary union, $SO(X)$ is a supratopology on X . For any α -set A in X , $A \cap B \in SO(X)$ for all $B \in SO(X)$. Thus A is an m -set with $SO(X)$. That is, $\alpha(X)$ is τ_m with $SO(X)$.

LEMMA 2.3. Let (X, τ^*) be a supratopological space. Then the class τ_m of all m -sets with τ^* is contained in τ^* .

Proof. Let A be an m -subset with τ^* . And X is an element of τ^* . Now we take that $X \cap A = A$ belongs to the supratopology τ^* , by the definition of m -sets. \square

THEOREM 2.4. Let (X, τ^*) be a supratopological space. Then the class τ_m of all m -sets with τ^* is a supratopology.

Proof. Let $\{A_\alpha\}$ be a class of members of τ_m . By definitions of the m -set and the supratopology, $(\cup A_\alpha) \cap T = \cup(A_\alpha \cap T) \in \tau^*$ for all $T \in \tau^*$. Thus the union $\cup A_\alpha$ also belongs to τ_m . \square

THEOREM 2.5. Let (X, τ^*) be a supratopological space with $\emptyset \in \tau^*$. If a subset A of X is a singleton set and $A \in \tau^*$, then A is an m -set.

Proof. Since $A \in \tau^*$ is a singleton set, $A \cap B = \emptyset$ or A for $B \in \tau^*$. Thus A is an m -set. \square

We obtain the following, by definition of m -set.

THEOREM 2.6. *Let (X, τ^*) be a supratopological space. If T is any supraopen set of τ^* in X and A is an m -set with τ^* , then $T \cap A$ is also a supraopen set.*

COLORALLY 2.7. *Let (X, τ) be a topological space and $\tau^* = PO(X)$. If $A \in \alpha(X)$ and $B \in PO(X)$, then $A \cap B \in PO(X)$.*

Proof. Since $\alpha(X) \subset SO(X) \cap PO(X)$, $\alpha(X)$ is a subclass of m -sets with $PO(X)$, and it obtained by Theorem 2.6. \square

THEOREM 2.8. *Let (X, τ^*) be a supratopological space with $\emptyset \in \tau^*$. Then the class τ_m of all m -subsets of X is a topology on X .*

Proof. Since $\emptyset \cap T = \emptyset \in \tau^*$ and $X \cap T = T \in \tau^*$ for all $T \in \tau^*$, \emptyset and $X \in \tau_m$.

Suppose $A, B \in \tau_m$. By definition of m -set, we obtain $B \cap T \in \tau^*$ and $A \cap (B \cap T) \in \tau^*$ for all $T \in \tau^*$. Thus $(A \cap B) \in \tau_m$.

And by Theorem 2.4., the proof is completed. \square

Now the class τ_m is called an m -topology with τ^* and the members of τ_m are called m -open sets. A subset B of X is called an m -closed set if the complement of B is an m -open set. Thus the intersection of any family of m -closed sets is a m -closed set and the union of finitely many m -closed sets is an m -closed set.

In case τ_m is an m -topology with τ^* on X , the topological space (X, τ_m) with τ^* will be denoted by (X, τ_m, τ^*) .

REMARK. In a space (X, τ) , if τ^* is an associated supratopology with τ , an m -set need not be an open set, and vice versa.

EXAMPLE 2.9.

Let $X = \{a, b, c, d\}$. Consider $\tau = \{\emptyset, X, \{a, b\}\}$ and $\tau^* = \{\emptyset, X, \{a, b\}, \{b, d\}, \{a, b, d\}\}$. Then τ^* is a supratopology associated with τ and $\{a, b, d\}$ is an m -set but it is not an open set. And $\{a, b\}$ is an open set but it is not an m -set.

DEFINITION 2.10. Let (X, τ_m, τ^*) be an m -topological space.

- (1) The m -interior of A is defined as the union of all m -open sets contained in A . The m -interior of A is denoted by $\text{mint}A$.
- (2) The m -closure of A is defined as the intersection of all m -closed sets containing A . The m -closure of A is denoted by $\text{mcl}A$.

By the above definitions, we obtain the following properties.

THEOREM 2.11. Let (X, τ_m, τ^*) be an m -topological space and A be a subset of X .

- (1) A is m -open if and only if $A = \text{mint}A$.
- (2) A is m -closed if and only if $A = \text{mcl}A$.
- (3) $\text{mcl}(\text{mcl}A) = \text{mcl}A$ and $\text{mint}(\text{mint}A) = \text{mint}A$.
- (4) $A \subset B$ implies $\text{mcl}A \subset \text{mcl}B$.
- (5) $\text{mcl}A \cup \text{mcl}B = \text{mcl}(A \cup B)$.

3. m -continuity

DEFINITION 3.1. Let (X, τ_m, τ^*) be an m -topological space and (Y, μ) be a topological space. A mapping $f : X \rightarrow Y$ is called an m -continuous if the inverse image of each open set of Y is an m -open set in X .

REMARK. In general, there is no relation between the continuity and the m -continuity.

EXAMPLE 3.2. Let $X = \{a, b, c, d\}$, $\tau = \{\emptyset, X, \{a, b\}\}$ and $\mu = \{\emptyset, X, \{a, b, d\}\}$. Now we take a supratopology $\tau^* = \{\emptyset, X, \{a, b\}, \{b, d\}, \{a, b, d\}\}$ for τ . Then $\tau_m = \{\emptyset, X, \{a, b, d\}\}$. Let $f : (X, \tau, \tau^*) \rightarrow (X, \tau)$ be the identity function. Then f is continuous but it is not m -continuous. And if $f : (X, \tau, \tau^*) \rightarrow (X, \mu)$ be the identity function. Then f is m -continuous but it is not continuous.

THEOREM 3.3. Let (X, τ_m, τ^*) be an m -topological spaces and (Y, μ) be a topological spaces. If $f : (X, \tau_m, \tau^*) \rightarrow (Y, \mu)$ is a mapping, then the following statements are equivalent:

- (1) f is an m -continuous.

- (2) The inverse image of each closed set in Y is m -closed.
- (3) For each $x \in X$, and each open set $V \subset Y$ containing $f(x)$, there exists $W \in \tau_m$ such that $x \in W$, $f(W) \subset V$.
- (4) $f(mclA) \subset clf(A)$ for every $A \subset X$.
- (5) $mcl(f^{-1}(B)) \subset f^{-1}(cl(B))$ for every $B \subset Y$.

Proof. (1) \Rightarrow (2). Let B be closed in Y . Since $Y - B$ is open in Y and $X - f^{-1}(B)$ is m -open, thus $f^{-1}(B)$ is m -closed.

(2) \Rightarrow (1). Let V be open in Y . Since $Y - V$ is closed in Y and $X - f^{-1}(V)$ is m -closed, $f^{-1}(V)$ is m -open.

(1) \Rightarrow (3). For each $x \in X$, and each open set V containing $f(x)$. Set $W = f^{-1}(V)$. Then W is m -open, $x \in W$, and $f(W) \subset V$.

(3) \Rightarrow (4). We will show that for each $b \in mclA$, $f(b) \in cl(f(A))$. Let V be an open neighborhood of $f(b)$, then there exists $W \in \tau_m$ such that $b \in W$ and $f(W) \subset V$. Since $b \in mclA$, $W \cap A \neq \emptyset$. $f(W \cap A) \neq \emptyset$ and $f(W) \cap f(A) \neq \emptyset$. Thus $V \cap f(A) \neq \emptyset$ and $f(b) \in cl(f(A))$.

(4) \Rightarrow (5). Let $A = f^{-1}(B)$ for $B \subset Y$. Then $f(mcl(A)) \subset cl(f(A)) \subset cl(B)$, and $mcl(f^{-1}(B)) \subset f^{-1}(cl(B))$.

(5) \Rightarrow (2). Let $B \subset Y$ be closed. Then $mcl(f^{-1}(B)) \subset f^{-1}(cl(B)) = f^{-1}(B)$, and $f^{-1}(B)$ is an m -closed set. \square

REMARK. If $f : (X, \tau_m, \tau^*) \rightarrow (Y, \mu)$ is an m -continuous function and $g : (Y, \mu) \rightarrow (Z, \nu)$ is a continuous function, then $g \circ f$ is m -continuous.

LEMMA 3.4. Let $f : (X, \tau) \rightarrow (Y, \mu)$ be an α -continuous function. Then

- (1) For each subset A of X , $f(cl_\alpha(A)) \subset (f(A))^-$ if and only if $f(A^{-0-}) \subset (f(A))^-$.
- (2) For each subset B of Y , $cl_\alpha(f^{-1}(B)) \subset f^{-1}(B^-)$ if and only if $(f^{-1}(B))^{-0-} \subset f^{-1}(B^-)$.

Proof. Since $cl_\alpha(A) = A \cup cl(int(cl(A)))$, the properties are proved obviously. \square

By Theorem 3.3 and Lemma 3.4, easily we get the following properties.

COROLLARY 3.5. Let $f : (X, \tau_m, SO(X)) \rightarrow (Y, \mu)$ is a function, the followings are equivalent:

- (1) f is α -continuous.
- (2) The inverse image of each closed set in Y is m -closed set.
- (3) For each $x \in X$, and each open set $V \subset Y$ containing $f(x)$, there exists $W \in \tau_m$ such that $x \in W$, $f(W) \subset V$.
- (4) $f(A^{-0-}) \subset cl(f(A))$ for every $A \subset X$.
- (5) $(f^{-1}(B))^{-0-} \subset f^{-1}(cl(B))$ for every $B \subset Y$.

DEFINITION 3.6. A function $f : (X, \tau_m, \tau^*) \rightarrow (Y, \mu_m, \mu^*)$ is an mS -continuous function if the inverse image of each m -set in Y is a supraopen set in X .

The following theorem is a straightforward result of Mashhour(Theorem 2.1.[5]).

THEOREM 3.7. Let $f : (X, \tau, \tau^*) \rightarrow (Y, \mu, \mu^*)$ be a function. Then the followings are equivalent :

- (1) f is an mS -continuous.
- (2) The inverse image of each m -closed set in Y is a supraclosed set.
- (3) $(f^{-1}(V))^{sc} \subset f^{-1}(mcl(V))$, for every $V \subset Y$.
- (4) $f(U^{sc}) \subset mcl(f(U))$, for every $U \subset X$.
- (5) For any point $x \in X$ and any m -open set V of Y containing $f(x)$, there exists $U \in \tau^*$ such that $x \in U$ and $f(U) \subset V$.

REMARK. Let $f : (X, \tau, \tau^*) \rightarrow (Y, \mu, \mu^*)$ be a function. Then we can get the following diagrams :

- (1) m -continuity $\implies S$ -continuity
- (2) S^* -continuity $\implies mS$ -continuity
- (3) In $\tau \subset \tau_m$,

$$\text{continuity} \implies m\text{-continuity} \implies S\text{-continuity}$$

- (4) In $\tau \subset \tau_m$ and $\mu \subset \mu_m$,

$$m\text{-continuity} \implies S\text{-continuity} \longleftarrow mS\text{-continuity} \longleftarrow S^*\text{-continuity}$$

(5) In $\tau^* = SO(X)$,

continuity $\implies m$ -continuity ($=\alpha$ -continuity) \implies semi-continuity

(6) In $\tau^* = PO(X)$,

continuity $\implies \alpha$ -continuity $\implies m$ -continuity \implies precontinuity

References

1. N. Levine, *Semi-open sets and semi-continuity in topological spaces*, Am. Math. Monthly **70** (1963), 36-41.
2. A.S.Mashhour, M.E.Abd El-Monsef and S.N.El.Deeb, *On precontinuous and weak precontinuous mappings*, Proc. Math. Phys. Soc. Egypt **51** (1981a), 47-53.
3. O. Njastad, *On some classes of nearly open sets*, pacific journal of mathematics **15** (1964), 961-970.
4. A.S.Mashhour, I.A.Hasanein and S.N.El-deeb, *α -cotinuous and α -open mappings*, Acta Math. Hung. **41** (1983), 213-218.
5. A.S.Mashhour, A.A.Allam, F.S.Mahmoud and F.H.Khedr, *On supratopological spaces*, Indian J. Pure Appl. Math **14** (1983), 502-510.

Department of Mathematics
Kangwon National University
Chuncheon, 200-701, Korea
E-mail: wkmin@cc.kangwon.ac.kr