

## Bioisoster of Capsaicin: Synthesis of 1-Hydroxy-2-pyridone Analogue

Hye-Young Choi and Sung-Hwa Yoon\*

Department of Applied Chemistry, Ajou University, Suwon 442-749, Korea


Received March 31, 1999

After capsaicin (**1**), a pungent principle of *capsicum*,<sup>1</sup> was known to act on a subset of peripheral sensory neuron,<sup>2</sup> molecular approach toward more potent capsaicin agonists which are anticipated to be useful as novel analgesic agents has been extensively studied.<sup>3</sup> Three parts of capsaicin structure - the aromatic ring region, the amide bond region, and the side chain region - were modified and it was found that the catechol moiety in capsaicin is important for the analgesic activity.<sup>3a</sup>

By application of bioisosterism, we previously found that the analogue which replaced the catechol moiety in dopamine with 1-hydroxy-2-pyridone system also revealed the similar dopaminergic activity.<sup>4</sup> Based on our result, we now replaced the functional catechol moiety in capsaicin with 1-hydroxy-2-pyridone system whose isosteric/isoelectric character is considered to be equivalent. And, therefore, they are interchangeable as far as its contribution to biological activity.<sup>5</sup> Here, we report the synthesis and the biological activity of 1-hydroxy-2-pyridone analogue (**2**) of capsaicin.


The synthesis of target compound **2** was started with the conversion of pyridine-N-oxide into pyridone compound (Scheme 1). Thus, 4-cyanopyridine-N-oxide **3** was refluxed with acetic anhydride followed by acid hydrolysis with acetic acid to give the pyridone **4**. After *O*-methylation of pyridone,<sup>6</sup> the free amine **6** was obtained from the reduction of the cyano group in **5** with LiAlH<sub>4</sub>. Protection of the free amine with trifluoroacetyl group followed by N-oxidation of the resulting compound **7** with *m*-chloroperbenzoic acid gave the N-oxide **8**, where the trifluoroacetyl group was then deprotected under basic condition to give 4-aminomethyl-2-methoxypyridine-1-oxide (**9**). Coupling reaction of **9** with (*E*)-8-methyl-6-nonenic acid (**11**), which was prepared by the reported method,<sup>7</sup> under DCC/DMAP condition gave **10** in 52% yield. Finally, the methyl group was removed by refluxing with acetyl chloride followed by hydrolysis with acetone-water mixture<sup>8</sup> to give **2**. However, it is interesting to note that the coupling reaction of 4-aminomethyl-1-hydroxy-2-pyridone (**12**), which was prepared from **8**, with the corresponding acid **11** under various conditions was unsuccessful to obtain **2** (Scheme 2).

The analgesic activity of **2** was determined by the reported




Capsaicin (**1**)

Figure 1



**Scheme 1.** (a) i) Ac<sub>2</sub>O, reflux, ii) AcOH, reflux. (b) Ag<sub>2</sub>CO<sub>3</sub>, CH<sub>3</sub>I, benzene. (c) LiAlH<sub>4</sub>, ether. (d) TFAA, CH<sub>2</sub>Cl<sub>2</sub>. (e) *m*CPBA, CH<sub>2</sub>Cl<sub>2</sub>. (f) K<sub>2</sub>CO<sub>3</sub>, MeOH. (g) (*E*)-8-Methyl-6-nonenic acid (**11**), DCC, DMAP, CH<sub>2</sub>Cl<sub>2</sub>. (h) i) AcCl, ii) H<sub>2</sub>O, acetone.



**Scheme 2.** (a) i) AcCl, ii) H<sub>2</sub>O. (b) MeOH : H<sub>2</sub>O : c-HCl (1 : 2 : 2).

method<sup>3a</sup> and observed to be inactive (ED<sub>50</sub> > 50 μmol/kg).<sup>9</sup> It can therefore be concluded that the catechol moiety in capsaicin makes a major contribution to the analgesic activity.

### Experimental

**Instrument.** Melting points were determined on a

Fisher-Johns melting point apparatus and are uncorrected.  $^1\text{H}$  NMR and  $^{13}\text{C}$  NMR spectra were recorded on a Varian Gemini 200 spectrometer at 200 MHz and 50 MHz. Chemical shifts were given in relative tetramethylsilane. Infrared spectra were recorded on a Nicolet FT-IR 550 spectrometer. Elemental analyses were performed by Fisons Eager 200 instrument, Italy. Column chromatography was done by using Merck silica gel 60 (230-400 mesh). All reactions were performed under a nitrogen atmosphere.

**4-Cyano-2-pyridone (4).** A solution of 4-cyanopyridine-N-oxide (50.0 g, 0.140 mol) in acetic anhydride (500 mL) was refluxed for 18 h and then concentrated. The residue was dissolved in ethyl acetate to give black precipitate which was identified with reactant. After filtering the precipitate, the filtrate was concentrated to give a brown residue, which was chromatographed on a silica gel column (ethyl acetate : hexane = 1 : 1) to give 4-cyano-2-acetoxypyridine (11.5 g, 23%) as a yellow solid, mp 60 °C. IR (KBr) 2243, 1775, 1196  $\text{cm}^{-1}$ .  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  2.31 (s, 3H,  $-\text{CH}_3$ ), 7.33 (s, 1H, 3-py-H), 7.44 (d, 1H, 5-py-H,  $J = 4.8$ ), 8.53 (d, 1H, 6-py-H,  $J = 4.8$ ).  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ )  $\delta$  20.8, 115.4, 118.7, 123.2, 123.4, 149.7, 158.0, 168.0. A solution of 4-cyano-2-acetoxypyridine (8.5 g, 50 mmol) in acetic acid (300 mL) was heated at 100-110 °C for 3 h and then concentrated. The resulting residue was recrystallized from methanol to give **4** (4.4 g, 70%) as a violet solid, mp 235 °C. IR (KBr) 2237, 1676  $\text{cm}^{-1}$ .  $^1\text{H}$  NMR ( $\text{DMSO}-d_6$ )  $\delta$  6.46 (dd, 1H, 5-py-H,  $J = 1.0, 6.6$  Hz), 6.97 (s, 1H, 3-py-H), 7.65 (dd, 1H, 6-py-H,  $J = 1.0, 6.6$  Hz), 12.25 (br, 1H,  $-\text{NH}$ ).  $^{13}\text{C}$  NMR ( $\text{DMSO}-d_6$ )  $\delta$  104.8, 116.5, 123.9, 126.0, 138.4, 160.8. Anal. Calcd for  $\text{C}_6\text{H}_4\text{N}_2\text{O}$ : C, 59.99; H, 3.36; N, 23.32. Found: C, 59.78; H, 3.37; N, 23.23.

**4-Cyano-2-methoxypyridine (5).** To a solution of **4** (6.2 g, 50 mmol) in benzene (50 mL) was added silver carbonate (15 g, 54 mmol) and iodomethane (18 mL, 290 mmol). The reaction mixture was stirred for 48 h at room temperature and then concentrated. The brown residue was chromatographed on a silica gel column (ethyl acetate : hexane = 1 : 1) to give **5** (5.0 g, 71%) as a pale yellow solid, mp 64 °C. IR (KBr) 2237, 1041  $\text{cm}^{-1}$ .  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  3.97 (s, 3H,  $-\text{OCH}_3$ ), 7.00 (s, 1H, 3-py-H), 7.08 (dd, 1H, 5-py-H,  $J = 1.1, 5.2$  Hz), 8.32 (d, 1H, 6-py-H,  $J = 5.2$  Hz).  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ )  $\delta$  53.9, 114.1, 116.4, 117.5, 122.3, 148.4, 164.3. Anal. Calcd for  $\text{C}_7\text{H}_6\text{N}_2\text{O}$ : C, 62.68; H, 4.51; N, 20.88. Found: C, 62.55; H, 4.50; N, 20.64.

**4-Aminomethyl-2-methoxypyridine (6).** To a suspension of  $\text{LiAlH}_4$  (1.6 g, 2.0 eq.) in anhydrous ether (200 mL) was slowly added **5** (2.4 g, 18 mmol) in ether (15 mL) at 0 °C. The reaction mixture was stirred for 3 h at room temperature and then quenched with 10% aqueous NaOH solution (10 mL). After the resulting gray precipitate was filtered, the organic solvent was removed. The residue was chromatographed on a silica gel column (ethyl acetate : methanol = 1 : 5) to give **6** (1.7 g, 71%) as a yellow oil. IR (neat) 3367, 3317  $\text{cm}^{-1}$ .  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  3.84 (s, 2H,  $-\text{CH}_2$ ), 3.93 (s, 3H,  $-\text{OCH}_3$ ), 6.70 (s, 1H, 3-py-H), 6.83 (d, 1H, 5-py-H,  $J = 5.2$  Hz), 8.08 (d, 1H, 6-py-H,  $J = 5.2$  Hz).  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ )

$\delta$  53.9, 114.1, 116.4, 117.5, 122.3, 148.4, 164.3.

**4-Trifluoroacetylaminomethyl-2-methoxypyridine (7).** To a solution of **6** (1.0 g, 75 mmol) in methylene chloride (100 mL) was slowly added trifluoroacetic anhydride (1.7 mL). After stirring for 1 h at room temperature, the reaction mixture was neutralized with 10% aqueous  $\text{NaHCO}_3$  solution and then extracted with methylene chloride ( $3 \times 50$  mL). The organic layer was washed with water ( $2 \times 50$  mL), dried ( $\text{Na}_2\text{SO}_4$ ), filtered, and concentrated to give **7** (1.37 g, 78%) as a yellow solid, mp 58 °C. IR (KBr) 3310, 1701, 1147  $\text{cm}^{-1}$ .  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  3.78 (s, 3H,  $-\text{OCH}_3$ ), 4.34 (d, 2H,  $-\text{CH}_2$ ,  $J = 6.0$  Hz), 6.51 (s, 1H, 3-py-H), 6.67 (d, 1H, 5-py-H,  $J = 5.4$  Hz), 7.98 (d, 1H, 6-py-H,  $J = 5.4$  Hz).  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ )  $\delta$  41.9, 53.1, 107.9, 118.5, 146.8, 148.3, 158.2, 164.4. Anal. Calcd for  $\text{C}_9\text{H}_9\text{N}_2\text{O}_2\text{F}_3$ : C, 46.16; H, 3.87; N, 11.96. Found: C, 46.00; H, 3.87; N, 11.87.

**4-Trifluoroacetylaminomethyl-2-methoxypyridine-N-oxide (8).** To a solution of **7** (0.98 g, 4.2 mmol) in  $\text{CH}_2\text{Cl}_2$  (50 mL) was added *m*-chloroperbenzoic acid (70-75%, 1.8 g). After the resulting mixture was stirred for 12 h at room temperature, the organic solvent was removed. The residue was chromatographed on a silica gel column (ethyl acetate : hexane = 3 : 1) to give **8** (0.75 g, 72%) as a white solid, mp 180 °C. IR (KBr) 3159, 1724, 1211  $\text{cm}^{-1}$ .  $^1\text{H}$  NMR ( $\text{DMSO}-d_6$ )  $\delta$  3.98 (s, 3H,  $-\text{OCH}_3$ ), 4.42 (d, 2H,  $-\text{CH}_2$ ,  $J = 3.1$  Hz), 6.93 (d, 1H, 5-py-H,  $J = 6.1$  Hz), 7.16 (s, 1H, 3-py-H), 8.23 (d, 1H, 6-py-H,  $J = 6.1$  Hz), 10.13 (br, 1H,  $-\text{NH}$ ).  $^{13}\text{C}$  NMR ( $\text{DMSO}-d_6$ )  $\delta$  41.4, 57.2, 108.1, 116.6, 137.0, 139.3, 156.4, 157.1, 158.0.

**4-Aminomethyl-2-methoxypyridine-N-oxide (9).** To a solution of **8** (0.75 g, 30 mmol) in water (10 mL) and methanol (40 mL) was added potassium carbonate (0.81 g). The reaction mixture was stirred for 5 h at room temperature and then concentrated. The residue was chromatographed on a silica gel column (methanol) to afford **9** (0.15 g, 32%) as a yellow oil. IR (KBr) 3418, 1176  $\text{cm}^{-1}$ .  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  3.86 (s, 2H,  $-\text{CH}_2$ ), 4.02 (s, 3H,  $-\text{OCH}_3$ ), 6.85 (dd, 1H, 5-py-H,  $J = 1.5, 6.7$  Hz), 6.95 (s, 1H, 3-py-H), 8.14 (d, 1H, 6-py-H,  $J = 6.7$  Hz).

**(E)-8-Methyl-6-nonenic Acid (11).** This was prepared by following the procedure reported by H. Kaga *et al.*,<sup>7</sup> yellow oil. IR (neat) 3409, 1706, 1265  $\text{cm}^{-1}$ .  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  0.96 (d, 6H,  $(\text{CH}_3)_2$ ), 1.42 (q, 2H,  $\text{C}_3\text{-H}$ ), 1.63 (q, 2H,  $\text{C}_4\text{-H}$ ), 2.02 (q, 2H,  $\text{C}_5\text{-H}$ ), 2.32 (m, 1H,  $\text{C}_8\text{-H}$ ), 2.36 (t, 2H,  $\text{C}_2\text{-H}$ ), 5.35-5.38 (m, 2H,  $\text{CH}=\text{CH}$ ).  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ )  $\delta$  23.1, 24.2, 26.4, 26.8, 29.2, 34.0, 126.5, 138.0, 180.3.

**1-Oxo-2-methoxypyridyl-8-methyl-6-trans-nonenamide (10).** The mixture of **9** (0.10 g, 64 mmol), (E)-acid **11** (0.12 g, 69 mmol), DCC (0.12 g, 58 mmol), and a catalytic amount of DMAP in methylene chloride (25 mL) was stirred for 8 h at room temperature. After filtration of the precipitate, the filtrate was concentrated. The resulting residue was chromatographed on a silica gel column (ethyl acetate : methanol = 1 : 1) to give **10** (98 mg, 52%) as a yellow oil. IR (KBr) 3267, 1654, 1265  $\text{cm}^{-1}$ .  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  0.93 (d, 6H,  $-(\text{CH}_3)_2$ ), 1.43 (q, 2H,  $\text{C}_3\text{-H}$ ), 1.64 (q, 2H,  $\text{C}_4\text{-H}$ ), 1.98 (q, 2H,  $\text{C}_5\text{-H}$ ), 2.04 (m, 1H,  $\text{C}_8\text{-H}$ ), 2.26 (t, 2H,  $\text{C}_2\text{-H}$ ), 3.98

(s, 3H, -OCH<sub>3</sub>), 4.36 (d, 2H, -CH<sub>2</sub>NH-,  $J = 6.0$  Hz), 5.35-5.38 (m, 2H, CH=CH), 6.78 (d, 2H, 3,5-py-H,  $J = 6.6$  Hz), 8.09 (d, 1H, 6-py-H,  $J = 5.5$  Hz), 8.12 (br., 1H, -NH-). <sup>13</sup>C NMR (CDCl<sub>3</sub>)  $\delta$  22.6, 25.2, 29.3, 30.9, 32.2, 36.2, 41.7, 57.2, 107.1, 116.3, 126.3, 138.0, 139.2, 141.9, 157.9, 173.7.

**4-Aminomethyl-1-hydroxy-2-pyridone (12).** A solution of **8** (150 mg, 6.4 mmol) in acetyl chloride (20 mL) was refluxed for 1 h and then concentrated. The resulting yellow residue dissolved in water (20 mL) was stirred for 14 h at room temperature. After the solvent was evaporated, the crude 1-hydroxy-2-pyridone derivative was obtained as a white solid, which was dissolved in small amount of methanol-water-c-HCl (1 : 2 : 2, 20 mL) again and then refluxed for 17 h. After the reaction mixture was concentrated, the yellow precipitate was recrystallized from ethanol-water to give the HCl salt form of **12** (54 mg, 48%) as a yellow solid, mp 230°C. IR (KBr) 3427, 1650 cm<sup>-1</sup>. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>)  $\delta$  3.90 (s, 2H, CH<sub>2</sub>), 6.35 (d, 1H, 3-py-H,  $J = 5.0$  Hz), 6.60 (s, 1H, 5-py-H), 7.94 (d, 1H, 6-py-H,  $J = 7.2$  Hz), 8.50 (br., 3H, -NH<sub>3</sub><sup>+</sup>). <sup>13</sup>C NMR (DMSO-d<sub>6</sub>)  $\delta$  104.1, 118.1, 136.0, 144.8, 157.4. Free base was obtained from the following method: after the salt was dissolved in ammonia solution, extraction with methylene chloride followed by general work-up gave the free base, which was used for the coupling reaction immediately.

**N-Hydroxy-2-oxopyridyl-8-methyl-6-trans-nonenamide (2).** A solution of **10** (70 mg, 2.0 mmol) in acetyl chloride (6 mL) was refluxed for 1 h and then neutralized with 10% aqueous NaHCO<sub>3</sub> solution. The aqueous solution was extracted with methylene chloride (3  $\times$  10 mL). The organic layer was washed with water, dried (Na<sub>2</sub>SO<sub>4</sub>), filtered, and concentrated. The resulting yellow residue dissolved in small amount of acetone and water was stirred for 8 h at room temperature. The precipitate was filtered and then dried to give **2** (50 mg, 71%) as a slightly yellow solid, mp 107°C. IR (KBr) 3410, 3279, 1639 cm<sup>-1</sup>. <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  0.96 (d, 6H, (CH<sub>3</sub>)<sub>2</sub>), 1.35 (q, 2H, C<sub>3</sub>-H), 1.64 (q, 2H, C<sub>4</sub>-H), 1.98 (q, 2H, C<sub>5</sub>-H), 2.15 (m, 1H, C<sub>8</sub>-H), 2.27 (t, 2H, C<sub>2</sub>-H), 4.24 (s, 2H, -CH<sub>2</sub>NH-), 5.34 (m, 2H, CH=CH), 6.21 (s, 1H, 5-py-H), 6.40 (s, 1H, 3-py-H), 7.00 (br., 1H, -NH-), 7.64 (d, 1H, 6-py-H,  $J = 2.1$  Hz). <sup>13</sup>C-NMR (CDCl<sub>3</sub>)  $\delta$  19.5, 20.8,

21.7, 22.6, 24.5, 30.2, 30.4, 92.9, 107.1, 115.8, 123.3, 124.5, 126.4, 141.8, 176.8.

**Acknowledgment.** The authors wish to acknowledge the financial support of Korean Research Foundation made in the program year of 1997.

## References

1. Szolcsanyi, J., In *Handbook of Experimental Pharmacology. Pyretics and Antipyretics*, Milton, A. S., Ed.; Springer: New York, 1982; pp 437-478.
2. Fitzgerald, M.; Capsaicin and Sensory Neuron- A Review, *Pain* **1983**, *15*, 109.
3. (a) Walpole, C. S. J.; Wrigglesworth, R.; Bevan, S.; Campbell, E. A.; Dray, A.; James, I. F.; Perkins, M. N.; Reid, D. J.; Winter, J. *J. Med. Chem.* **1993**, *36*, 2362. (b) Walpole, C. S. J.; Wrigglesworth, R.; Bevan, S.; Campbell, E. A.; Dray, A.; James, I. F.; Perkins, M. N.; Reid, D. J.; Winter, J. *J. Med. Chem.* **1993**, *36*, 2373. (c) Walpole, C. S. J.; Wrigglesworth, R.; Bevan, S.; Campbell, E. A.; Dray, A.; James, I. F.; Perkins, M. N.; Reid, D. J.; Winter, J. *J. Med. Chem.* **1993**, *36*, 2381. (d) Wrigglesworth, R.; Walpole, C. S. J.; Bevan, S.; Campbell, E. A.; Dray, A.; James, I. F.; Perkins, M. N.; Reid, D. J.; Winter, J. *J. Med. Chem.* **1996**, *39*, 4942. (e) Park, N-S.; Choi, J-K.; Hong, M-S.; Lee, J. C.; Choi, S. W.; Lee, B-Y. *Korean J. Med. Chem.* **1993**, *3*, 142. (f) Back, G. H.; Jung, Y. S.; Cho, S. J.; Seong, C. M.; Park, N-S. *Arch. Pharm. Res.* **1997**, *20*, 659.
4. Yoon, S-H.; Bodor, N. S.; Simpkins, J. W. *Drug Design and Discovery* **1993**, *10*, 35.
5. Friedman, H. I., Influence of isosteric replacements upon biological activity. Symposium on Chemical-Biological correlation. National Academy of Science-National Research Council, Publication no. 206, Washington D. C. 1951; p 295.
6. Chung, m. M.; Tieckmann, H. *J. Org. Chem.* **1970**, *35*, 2517.
7. Kaga, H.; Miura, M.; Orito, K. *J. Org. Chem.* **1989**, *54*, 2477.
8. Paquette, L. A. *J. Am. Chem. Soc.* **1965**, *87*, 5186.
9. Mouse tail flick test was performed at Center for Drug Discovery, University of Florida, U.S.A. according to the procedure described in reference 3a.