

□ Brief Communication □

Infestation status of head louse and treatment with lindane shampoo in children of primary school and kindergarten in Chinju-shi, Kyongsangnam-do, Korea

Yong Chan HA^{1)*}, Jung Min HEO¹⁾, Hyung Joon KIM¹⁾, Gi Moon GO¹⁾,
Sang Jeong LEE¹⁾, Song Hoan JEONG¹⁾, Seung In AHN¹⁾, Min Cheol KIM¹⁾,
Jeong Eun KIM¹⁾, Hye Young SONG¹⁾, Jeong Woon PARK¹⁾,
Byung Sung KIM²⁾ and Woon-Mok SOHN³⁾

*The Chinju Association of Public Health Doctors¹⁾, Chinju-shi Health Center²⁾,
Chinju 660-030, and Department of Parasitology³⁾, College of Medicine,
Gyeongsang National University, Chinju 660-751, Korea*

Abstract: The infestation status of head louse among children attending primary schools and kindergartens in Chinju-shi, Kyongsangnam-do, Korea, was investigated between June and July 1999. Out of 2,288 children examined, 3.9% of boys (48/1,242) and 23.5% of girls (246/1,046) were infested with nits or adult/nymphs of lice. The effectiveness of lindane shampoo (1% gamma benzene hexachloride solution) was evaluated after one or two time applications to all the children infested. The negative conversion rate of pediculosis was 93.5%. Effective control measures are needed to control and prevent such ectoparasite infestation amongst children.

Key words: head louse infestation, *Pediculus humanus capitis*, pediculosis, treatment, lindane shampoo, Chinju-shi

Since the first report on the head louse infestation in Korea (Lee et al., 1984), investigators have reported some prevalent cases from school children (Kim et al., 1984; Pai and Huh, 1987; Pai et al., 1989; Ree et al., 1992; Huh et al., 1993; Hong et al., 1995) and those of people who were admitted to mental hospitals and public welfare facilities (Pai, 1992; Huh et al., 1994; Huh and Pai, 1995). Furthermore, the drug effectiveness for delousing was evaluated (Pai, 1992; Ree et al., 1992). In this study, we investigated the infestation status of head louse and the effect

of lindane shampoo on children in primary schools and kindergartens.

A total of 2,288 primary school and kindergarten children from 11 regions (myons) in Chinju-shi was examined for the presence of nits, nymphs and/or adult head lice by naked eyes from June through July 1999. All children infested with lice were treated with lindane shampoo (1% gamma benzene hexachloride). The effectiveness of the treatment was evaluated by examining the presence of nits or lice at 7-10 days after the treatment. If the live nits or lice were detected, the children were subjected to the second shampoo treatment.

Out of 2,288 children examined, 294 (12.8%) were found to be infested with nits

• Received 13 September 1999, accepted after revision 19 January 2000.

*Corresponding author (e-mail: hychan@hitel.net)

Table 1. Prevalence of head louse infestation in primary school and kindergarten children in 11 regions (myons), Chinju-shi

Administrative area (myon)	Name of School	No. of student examined	No. of student infested (%)
Ibansong	Ibansong	125	45 (36.0)
	Ibansong (kind.) ^{a)}	104	29 (27.9)
Ilbansong	Ilbansong	314	64 (20.4)
Sabong	Sabong	109	15 (13.8)
Chisu	Songjong	66	11 (16.7)
Chinsong	Chinseong	166	12 (7.2)
Kumkok	Kumkok	92	20 (21.7)
Munsan	Munsan	603	68 (11.3)
Taepyong	Taepyong	53	0 (0)
Sukok	Sukok	91	10 (11.0)
	Sukok (kind.) ^{a)}	48	1 (2.1)
Taekok	Tuko	82	13 (15.9)
	Taekok	238	0 (0)
	Tanmok	56	0 (0)
	Chiphyun	89	0 (0)
Michon	Angan	52	6 (11.5)
Total		2,288	294 (12.8)

^{a)}kindergarten

Table 2. Infestation status of head louse by different grades and sexes

Grade	Boys		Girls		Total	
	No. examined	No.(%) positive	No. examined	No.(%) positive	No. examined	No.(%) positive
Kind. ^{a)}	207	17(8.2)	163	32(19.6)	370	49(13.2)
1	190	5(2.6)	140	33(23.6)	330	38(11.5)
2	173	6(3.5)	150	41(27.3)	323	47(14.6)
3	175	4(2.3)	148	39(26.4)	323	43(13.3)
4	165	8(4.8)	153	37(24.2)	318	45(14.2)
5	172	4(2.3)	150	27(18.0)	322	31 (9.6)
6	160	4(2.5)	142	37(26.1)	302	41(13.6)
Total	1,242	48(3.9)	1,046	246(23.5)	2,288	294(12.8)

^{a)}kindergarten

and/or adults/nymphs of head lice. The prevalent cases by each school and each kindergarten are shown in Table 1. The positive rate of head louse was 3.9% (48/1,242) for the boys and 23.5% (246/1,046) for the girls. The prevalent cases according to different academic grades are presented in Table 2. The negative conversion rate of head lice infestation after the second treatment with lindane shampoo was 93.5% (Table 3).

In this study, the infestation rate was not significantly high when compared to those of previous reports (Kim et al., 1984; Lee et al., 1984; Pai and Huh, 1987; Pai et al., 1989; Huh et al., 1993). The infestation rate for the girls, however, was remarkably higher than the boys. This finding was consistent with those of previous studies (Pai et al., 1989; Huh et al., 1993; Hong et al., 1995). However, there were no significant differences in the

Table 3. The negative conversion rate of head lice infestation after treatment with lindane shampoo

Items	Boys	Girls	Total
No. of cases treated	48	246	294
No. of negative conversion cases after first treatment	41	177	218
No. of negative conversion cases after second treatment	5	52	57
Total No. (%) of negative conversion cases	46 (95.8)	229 (93.1)	275 (93.5)

infestation rates among different academic grades.

The negative conversion rate of head louse infestation after the second application of lindane shampoo was relatively high (93.5%). Taplin et al. (1986) obtained 43.3% pediculicidal efficacy in the treatment with 1% lindane shampoo, and 96.6% with 1% permethrin cream rinse. Pai et al. (1989) reported 87.1% reduction in patients after a double treatment with 0.2% permethrin solution. Pai et al. (1991) compared the pediculicidal effects between 24% benzylbenzoate solution and 0.2% permethrin solution after two sequential treatments. The reduction rates were 87.9% for the 24% benzylbenzoate solution group and 72.3% for the 0.2% permethrin solution group. On the other hand, Ree et al. (1992) reported a 93.4% reduction rate after a single application of Sumithrin powder in the mass treatment of head louse infestation.

For a successful control of such ectoparasites among children in primary schools and kindergartens, cooperations between the school authorities and the public health centers should be established, and a team of physicians, teachers, parents, and health care officials will be needed.

REFERENCES

- Hong HK, Kim CM, Lee JS, Lee WJ, Yang YC (1995) Infestation rate of head lice in primary school children in Incheon, Korea. *Korean J Parasitol* **33**: 243-244.
- Huh S, Pai KS (1995) Head louse infestation in vagrants and children admitted to public welfare facilities, Republic of Korea. *Korean J Parasitol* **33**: 69-71.
- Huh S, Kook J, Chai JY, Pai KS (1994) Infestation rate of lice in a mental hospital, Seoul, Korea. *Korean J Parasitol* **32**: 275-276.
- Huh S, Pai KS, Lee SJ, Kim KJ, Kim NH (1993) Prevalence of head louse infestation in primary school children in Kangwon-do, Korea. *Korean J Parasitol* **31**: 67-69.
- Kim TK, Park CP, Huh S (1984) Head louse infestation among the students in Yongyang-gun, Kyongsangbuk-do. *Korean J Parasitol* **22**: 273-276 (in Korean).
- Lee SH, Oh CW, Chai JY (1984) Head louse infestation among primary school children in Seosan-gun, Chungnam Province. *Korean J Parasitol* **22**: 141-143 (in Korean).
- Pai KS (1992) Head louse infestation among girls in an orphanage and women in a mental hospital and mass-delousing with Bio-allethrin (PARA™ aerosol). *Korean J Parasitol* **30**: 49-52 (in Korean).
- Pai KS, Huh S (1987) Head louse infestation among school children in Sanbok-myon, Munkyeong-gun, Kyongsangbuk-do (1985). *Korean J Parasitol* **25**: 85-86 (in Korean).
- Pai KS, Park MS, Lee YS, et al. (1989) The prevalence of head louse infestation among urban and rural children in Korea. *Korean J Parasitol* **27**: 271-275 (in Korean).
- Pai KS, Park MS, Lee YS, Lee JS, Kim PK, Huh S (1991) Therapeutic effects of 24% benzylbenzoate and 0.2% permethrin on head louse infestation. *J Korean Med Assoc* **34**: 1208-1218 (in Korean).
- Ree HI, Yong TS, Shin HJ, et al. (1992) Mass treatment of head louse infestation with Sumithrin powder in primary schools in Korea. *Korean J Parasitol* **30**: 349-354.
- Taplin D, Meinking TL, Castillero PM, Sanchez R (1986) Permethrin 1% creme rinse for the treatment of *Pediculus humanus* var. *capitis* infestation. *Pediatric Dermatol* **3**: 344-348.

IN MEMORIAM

CHONG-HWAN KIM
(1931-1999)



Professor Chong-Hwan Kim passed away on December 30, 1999, just 1 month after his 78th birthday. He was born on November 28, 1931 in Seochon-gun, Kyungsangbuk-do, and spent his childhood in Haman-gun. Professor Kim graduated from the Department of Biology, College of Teaching, at Seoul National University in 1957, and later earned his doctorate in Biology from Kon-kuk University in 1969.

Professor Kim's career in the field of Parasitology began in 1959 when he worked as a Teaching Assistant. He later became a full-time lecturer, Assistant Professor and Associate Professor in the Department of Parasitology, College of Medicine, Yonsei University where he worked until 1973. In 1973, Professor Kim established the Department of Parasitology, College of Medicine, Chungnam National University where he worked until 1979. He continued his research and teaching for the Department of Biology, College of Natural Science at Chungnam National University until his retirement in 1997.

Professor Kim's research interests dealt with various trematodes, snails and their relationships. In recognition of his efforts and studies, Professor Kim received the Academic Award of the Korean Society for Parasitology in 1981. Professor Kim left an inspiring legacy for those pursue research and teaching excellence in the field of Parasitology. His memory leaves those who knew him with a sense of affection and great admiration. He will not be forgotten for his mentorship and achievements by those, both young and old, who follow him in the challenges of Parasitology academia. May he rest in peace and never be forgotten.