

(1994 1999)

가

= Abstract =

Salmonellosis in Children in Daejeon, Korea, 1994 1999

Jung-Soo Chun, M.D., Kyung-Yil Lee, M.D., Hyung-Shin Lee, M.D.
Sang-Yong Kim, M.D., Ji-Whan Han, M.D., Jae-Kyun Her, M.D.
and Tyung-Tai Whang, M.D.

Department of Pediatrics, College of Medicine, The Catholic University of Korea, Seoul, Korea

Purpose : Salmonellosis including typhoid fever is still prevalent in Korea. Recently it has been reported that the incidence of salmonella gastroenteritis is increasing with a reduction of typhoid fever. We studied the clinical and laboratory features of salmonellosis.

Methods : We evaluated the clinical records of 83 patients with salmonellosis who had been confirmed by stool culture from 1994 to 1999.

Results : Annual incidence of salmonellosis showed an increasing tendency during recent 4 years. Seasonally, summer(45.8%) was the most prevalent, followed by fall(32.5%). Male to female ratio was 1.4 : 1. In age distribution, 64 cases(77.1%) were under 5 year-old, and 18 cases(21.7%) were younger than 1 year-old. Clinical features included diarrhea(96.4%), fever(91.6%), vomiting(49.4%), bloody stool(42.1%), abdominal pain(40.1%) and tenesmus(12.0%). In serogroups, there was no group A and group B, group C, group D(including 2 cases of *S.typhi*) and group E were in 41.0%(34 cases), 3.6%(3 cases), 51.8%(43cases) and 3.6% (3 cases), respectively. In Widal test, 5 cases(13.5%) and 1 case(2.7%) of group B, C and E(total 37 cases) were observed the O titer above 1 : 80 and 1 : 320, respectively. However, in 36 cases of group D, 19 cases(52.7%) and 9 cases(25.0%) were above the O titer 1 : 80 and 1 : 320, respectively. Antibiotics resistant rates to ampicillin, trimethoprim/sulfamethoxazole and chloramphenicol were 23.2%, 10.1% and 51.4%, respectively.

Conclusion : Salmonellosis has become a common cause of gastroenteritis in children, especially under the pre-school age. Isolation of salmonella is necessary for accurate diagnosis of gastroenteritis and typhoid fever. A careful attention of the use of antibiotics is needed to reduce the multi-drug resistant strains.

Key Words : Salmonellosis, Serogroup, Widal test

: , 가

Tel : 042)220-9541 Fax : 042)221-2925 E-mail : leekyungyil@yahoo.com

S.typhi 가 1994 1 1999 12 6 가 83 , Widal 1994 1 1999 12 6 가 83 McConkey , Selenite-F broth Salmonella-shigella 1 , 4 (TSI , MIO , Citrate , Urease) (serotype) 7 (Poly, Vi, A, B, C, D, E) Kirby-Bauer ampicilline, trimethoprim-sulfamethoxazole(bactrim), chloramphenicol, gentamicin, tobramycin, amikacin, ciprofloxacin, aztreonam, imipenem, cephalothin, cefuroxime, ceftriaxone, sulperazone 13

1. 1994 3 , 1995 7 1996 4 가 1999 14 23 가 1995 2 (Fig. 1). 38 (45.8%) 가 가 27 (32.5%), 12 (14.5%), 6 (7.2%) (Fig. 2). 2. 0 14 가 48 (57.85%), 가 35 14 : 1 . 5 가 64 (77.1%) 18 (21.7%) (Fig. 3). 3. 2 81 가 80 (96.4%) 가 (91.6%), (49.4%), (42.1%), (40.1%), tenesmus(12.0%), (7.2%), (3.6%), (2.4%) (Table 1). 가 38 , 가 45 2




Fig. 1. Annual cases of salmonellosis in children, 1994 1999.




Fig. 2. Monthly distribution of salmonellosis.




Fig. 3. Age and sex distribution of salmonellosis in children.

4 .

(serogroup) 83 A , B
 34 (41.0%), C 3 (3.6%), D 43 (51.8%),
 E 3 (3.6%) (Table 2). Widal
 가 B, C E 37 O titer 1:80
 5 (13.5%), 1:320 1 (2.7%)가
 , H titer 1:80 1 (2.7%)
 D 43 Widal 36

Table 1. Clinical Features on Admission

Symptoms & Signs	No. of Patients	%
Diarrhea	80	96.4
Fever	76	91.6
Vomiting	41	49.4
Bloody stool	35	42.1
Abdominal pain	33	40.1
Tenesmus	10	12.0
Headache	6	7.2
Convulsion	3	3.6
Lethargy	2	2.4

Table 2. Serogroups of Salmonella

Serotype	No.(% of Total)
A	0(0.0%)
B	34(41.0%)
C	3(3.6%)
D	43(51.8%)*
E	3(3.6%)
Total	83(100.0%)

* : contained 2 cases of *S.typhi*

Table 3. Results of Widal Test

Serotype	No. tested	Widal test(O titer)	
		1 : 80 No.(%)	1 : 320 No.(%)
A	0		
B	32	5(15.6)	1(3.1)
C	3	0(0.0)	0(0.0)
D	36	19(52.7)	9(25.0)
E	2	0(0.0)	0(0.0)

* : contained 2 cases of *S.typhi*

O titer 1 : 80 19 (52.7%), 1 : 320 9 (25.0%), H titer 1 : 80 5 (11.6%)가 (Table 3). D *S.typhi* 2 O titer 320 : 1, H titer 1 : 160 5,000/mm³ 5 (6.0%), 5,000 10,000/mm³ 31 (37.3%), 10,000/mm³ 47 (56.6%) . 가 AST, ALT 7 . CRP 69 가 (83.1%), 1 5mg/dL 49 , 6 10mg/dL 8 , 11 15mg/dL 6 , 15 20mg/dL 4 , 21mg/dL 2 . 5 .

13

(Table 4). Ampicil-

line 69 53 (76.8%)가 chloramphenicol 48.6% 가 ,

Table 4. Antibiotics Susceptibilities of Isolated Salmonella

Antibiotics	No. tested	No. of positive	% of total
Ampicillin	69	53	76.8
Bactrim	69	62	89.9
Chloramphenicol	37	18	48.6
Gentamicin	47	45	95.7
Tobramycin	14	14	100.0
Amikacin	54	53	98.1
Ciprofloxacin	74	68	91.9
Aztreonam	35	34	97.1
Imipenem	44	44	100.0
Cephalothin	48	34	70.8
Cefuroxime	49	47	95.9
Ceftriaxone	12	12	100.0
Sulperazone	25	25	100.0

aminoglycoside 3 cephalosporin 100% .

noncapsulated . Enterobacteriaceae family . *S.typhi*, *S.choleraesuis* *S.enteritidis* 3 species . *S.typhi* *S.choleraesuis* *S.enteritidis* 2,000

so-matic O Kauffmann-White A, B, C, D E 5가

(outbreak)

6)

가 8, 9)

1 가

(enteric fever),
 가 , ,
 serogroup
 1 5) . 70 가 , ,
 90 S.typhi가 , ,
 (Table 1).
 가 . 1970 , ,
 S.typhi가 90% 1980 가 ,
 60% , 90 25% 가 ,
 90 B (rapid gastric emptying)
 (S.typhimurium) , 1996 D 13) .
 (S.enteritidis) 가 . 가
 가 83 14) .
 D 41 (49.4%), B 34 (41.0%),
 C 3 (3.6%), E 3 (3.6%) S.typhi 2 1
 (2.4%)가 (Table 2). 가 15) .
 1994 가 ,
 4) 52 : 48 19 가
 24.2% , 4 가
 11.1% .
 . 1984 34
 10) 2 3 가 15 (44.1%)
 가 , B 17 가
 (50.0%) 가 C (23.5%), D (23.5%), E
 (8.8%), A 2.9% 5
 가 64 (77.1%)
 1 18 (21.7%) (Fig. 3).
 5 9
 4, 5, 11) , 가
 38 (45.8%) 27 (32.5%)가
 , , 가
 , , ,
 가 가 , , ,
 가 가 , , ,
 8, 12) . 가

Widal	가	36	D	2	S.ty-
phi	O titer가	1 : 320	,		
	D (S.enteritidis)		34	O titer	
	1 : 80	19	, 1 : 320	9	

(Table 3).

14, 18),
S.typhi
 가
 3 5, 19, 20)
 가
 ampicil-
 lin, trimethoprim-sulfamethoxazole(Bactrim) chlor-
 amphenicol 23.2%, 10.1% 51.4%
 aminoglycoside 3 cephalo-
 sporin
 (Table 4).

: 1996 1999
 14 23 4 가
 45.8%, 가 32.5%가
 가 48 (57.8%) 14 : 1 , 5
 가 64 (77.1%) 1
 18 (21.7%)
 가 80 (96.4%) 가
 (91.6%), (49.4%), (42.1%),
 (40.1%), tenesmus(12.0%), (7.2%), (3.6
 %), (2.45%)
 (serogroup) 83 A
 0%, B 41.0%, C 3.6%, D 51.8% E
 3.6% . Widal B, C E 37
 O titer 1 : 80 5 (13.5%), 1 : 320
 1 (2.7%)가 , D 36 O titer 1 : 80
 19 (52.7%), 1 : 320 9 (25.0%)
 ampicillin, tri-
 methoprim-sulfamethoxazole chloramphenicol
 23.2%, 10.1% 51.4% ,
 aminoglycoside 3 cephalosporin

가
 가
 : 1994 1 1999 12 6
 가
 83

- 1) , , . 1972
 1973;8:1-5.
- 2) KS Chung, KD Park, YH Ryu. A report on
 the *salmonella* cultures isolated in Korea(1977).
 1997;14:57-63.
- 3) , , , , ,
 . *Salmonella Shigella*
 (1987).
 1987;24:4 13-28.
- 4) , , , , .

- 1994;31:14-23.
- 5)
1996;33:25-36.
- 6) Saphra I, Winter JW. Clinical manifestations of Salmonellosis in man. N Eng J Med 1957;256: 1128-34.
- 7) Aserkoff B, Schroeder SA, Brachman PS. Salmonellosis in the United States- a five-year review. Am J Epidemiol 1970;92:13-24.
- 8) Rodriguez DC. International increase in Salmonella enteritidis :A new pandemic? Epidemiol Infect 1990;105:21-7.
- 9) Choo KE, Razif AR, Oppenheimer SJ, Ariffin WA, Lau J, Abraham T. Usefulness of the widal test in diagnosing childhood typhoid fever in endemic areas. J Pediatr. Child Health 1993; 29:36-9.
- 10)
Salmonella, Shigella .
1984;19:55-64.
- 11) salmonellosis 1994;37:199-204.
- 12)
2000;11:41-7.
- 13) Agunod M, Yamaguchi N, Lopez R, et al. Correlative study of hydrochloric acid, pepsin, and intrinsic factor secretion in newborns and infants. Am J Dig Dis 1969;14:400-14.
- 14) Aserkoff B, Bennett JV. Effect of antibiotic therapy in acute salmonellosis on the fecal excretion of salmonellae. N Eng J Med 1969;281: 636-40.
- 15) Centers for Disease Control and Prevention. Summary of notifiable disease. United States, 1994. MMWR 1994;43:1-80.
- 16) Kulkarni ML. Value of single Widal test in the diagnosis of typhoid fever. Indian Pediatr 1994; 31:1373-7.
- 17) Schroeder SA. Interpretation of serologic tests for typhoid fever. JAMA 1968;206:839-40.
- 18) Buchbald DS, Blazer MJ. A review of human salmonellosis.II. Duration of excretion following infection with nontyphi salmonella. Rev Infect Dis 1984;6:345-56.
- 19) *Chloramphenicol, Ampicillin, Carbenicillin* *Salmonella typhimurium* *Salmonella* (1980 1986). 1988;20: 145-55.
- 20) *Salmonella* 1989;32:1230-8.