

## Microbial Production of Riboflavin Using Riboflavin Overproducers, *Ashbya gossypii*, *Bacillus subtilis*, and *Candida famate*: An Overview

Seong Han Lim<sup>1</sup>, Jong Soo Choi<sup>2</sup>, and Enoch Y. Park<sup>1\*</sup>

<sup>1</sup>Laboratory of Biotechnology, Department of Applied Biological Chemistry, Faculty of Agriculture, Shizuoka University, 836 Ohya, Shizuoka 422-8529, Japan,

<sup>2</sup>Laboratory of Fermentation, R&D Center, BASF Co. Ltd., JeonBuk 573-400, Korea

**Abstract** In this paper, the microbial production of riboflavin is reviewed and includes descriptions of riboflavin overproducers, and the biosynthesis and details of the key-enzyme genes related to riboflavin. Three kinds of riboflavin overproducers are known; *Bacillus subtilis* and *Candida famate* utilize glucose as a carbon source, but the fungus *Ashbya gossypii* requires plant oil as its sole carbon source. The starting material in riboflavin biosynthesis is guanosine triphosphate (GTP), which is converted to riboflavin through six enzymatic reactions. Though *Bacillus subtilis*, *Candida famate*, and *Ashbya gossypii* operate via different pathways until GTP, they follow the same pathway from GTP to riboflavin. From the metabolic viewpoint, with respect to improved riboflavin production, the supplementation of GTP, a process-limiting precursor must be considered. The GTP fluxes originate from three sources, serine, threonine and glyoxylate cycles. The development of pathways to strengthen GTP supplementation using biotechnological techniques remains an issue for future research.

**Keywords:** riboflavin, *Ashbya gossypii*, vitamin B<sub>2</sub>, metabolic engineering

### INTRODUCTION

Microorganisms produce valuable polymers, such as proteins, nucleic acids, and polysaccharides as well as many smaller molecules useful in human and animal health [1]. The vitamins are an example, and are classified as chemical substances that control and affect the physiological processes, and which are essential for the metabolism and growth of animals, especially mammals. However, these compounds cannot be biosynthesized by mammalian cells, which have lost the capacity to synthesize vitamins, whereas lower organisms retain this ability. Therefore, vitamins are synthesized by a variety of plants and microorganisms, which are nutritional requirements for human and animals. Although many studies have been undertaken in the field of vitamin biosynthesis, progress has been slow because organisms require such small quantities of these compounds.

Riboflavin, the so called vitamin B<sub>2</sub>, is widely distributed in plants and animals, and plays an important role in organisms because it is the precursor of flavin mononucleotide (riboflavin 5'-monophosphate, FMN) and flavin adenine dinucleotide (FAD), which function as coenzymes for a wide variety of enzymes in the intermediate metabolism. Therefore, a daily dose of 2 mg of

riboflavin is essential for humans. Furthermore the vitamin is useful as an animal feed supplement. Riboflavin is primarily used in the pharmaceutical industry, though some are used in the food industry as an additive, and in the animal feed industry. Thus, vitamins have become one of the important fermentation products in biotechnology industry.

The world consumption of riboflavin was reported to be  $1.25 \times 10^6$  kg per year, about 30 years ago, again largely for human and animal uses [2]. However, at present, the amount of riboflavin produced is believed to exceed 3,000 tons per year. Riboflavin is commercially produced by chemical or by biochemical synthesis, the latter being the preferred route because it is cheaper [3]. As a result, current riboflavin production by fermentation is estimated to be about 2,500 tons per year. This process has the merits of saving half the cost, reducing waste and energy requirements, and using renewable resources like sugar or plant oil [4].

Recently, with the progress of metabolic engineering, many researchers turn their attentions to biological metabolisms to improve the production efficiency of process that produce bioproducts from microorganisms. Riboflavin as an essential compound for humans and animals, and is one of target biotechnical processes, though a vast amount of knowledge has already been accumulated. In this review, we describe and compare riboflavin overproducers, its biochemical and metabolic biosynthesis, and the genes related to key enzymes in riboflavin biosynthesis.

\* Corresponding author

Tel: +81-54-238-4887 Fax: +81-54-238-4887

e-mail: yspark@agr.shizuoka.ac.jp

## DEVELOPMENT OF MICROORGANISMS

Riboflavin is a unique vitamin that can be synthesized in large enough amounts by some fungi and bacteria to be exploited successfully on an industrial scale. Many microorganisms have been screened and studied for industrial production. *Clostridium acetobutylicum* was one of the earliest organisms used to produce riboflavin, and its productivity was only 100 mg/L [5]. Some organisms of the genus *Candida* [6-12] were also found to produce riboflavin, and among these, *Candida flarerii* achieved about 600 mg/L of riboflavin [13]. But in order to produce higher levels of riboflavin production using these yeasts, inhibition due to iron should be overcome [6]. The biochemical key to riboflavin overproduction appears to involve resistance to the repressive effects of iron. In the biochemical synthesis of riboflavin of normal microorganisms, it appears that iron represses almost all of the riboflavin biosynthetic enzymes, whereas riboflavin or a derivative inhibits the first enzyme of the pathway, GTP cyclohydrolase II. New processes to overcome iron inhibitions and novel mutation techniques [4] that inhibit purine biosynthesis using antimetabolite, tuberculin, in *Candida famate* have recently been developed and give riboflavin yields of 20 to 30 g/L [1,14]. On the other hand, unlike the yeast. Bacteria, iron inhibition is not a factor for the two yeast-like molds, *Eremothecium ashbyii* and *Ashbya gossypii*, which are natural overproducers of riboflavin. The productivity of these ascomycetes presently is believed to exceed 20 g/L [1]. Both are pathogenic on cotton and form needle-shaped ascospores.

### *Eremothecium ashbyii*

The overproduction of riboflavin by this strain was discovered by Guilliermond and co-workers [15] in 1935. Hence, because it is an overproducer of riboflavin, much work has been undertaken up on *E. ashbyii* over decades. In the early 1950s, media that supported riboflavin overproduction at 200 mg/L were developed [16-18]. Further addition of Tween 80 and certain proteins, casein or glycine to this allowed yields to reach 1.4 g/L [18]. Currently, yields of 1.5 to 2.5 g/L [5, 19] are possible when very crude sources are used as media.

To increase productivity, 8-azaguanine, an analog of guanine, which is a precursor of the riboflavin biosynthetic pathway, was used. Mutants were obtained to get this compound. The level of vitamin production obtained from the majority of mutants in a synthetic medium containing peptone was 2-4 times higher than the level of the original strain [20].

Recent studies with wild type of *E. ashbyii* have yielded 3.3 g/L of riboflavin using molasses and peanut seed cake as carbon and nitrogen source, respectively [21]. Some studies with yeast-like strains have been undertaken from the viewpoint of morphological observation. The ideal colony for riboflavin production was characterized by its medium growth intensity (colony size, 8-10 mm in diameter over a culturing period of 96

h), intense yellow color, absence of sector formation, and a smooth surface covered by air mycelia [22]. Production of riboflavin using a colony obtained on the basis of these observations reached 1.1 g/L from a starting position of 0.2 g/L, and these results were confirmed by subsequent experiments [23]. The capacity for riboflavin biosynthesis by *E. ashbyii* was characterized by the considerable heterogeneity of the population. This heterogeneity was expressed in a high frequency of splitting off of lemon colony, which had a vitamin production level three times lower than that of the orange colonies. A population of a culture of *E. ashbyii* showed high morphological instability, as well as considerable heterogeneity with respect to its ability to synthesize riboflavin [23]. The relationship between the pH value of the medium and morphological variations was also observed. Highest riboflavin productivity was reported at initial pH of 6.5, using *E. ashbyii* [24,25]. At this constant pH value, morphological variation was shown to involve the gradual increase of swollen hyphal cells and the formation of ascospores [26]. However, obtained at various pH values showed that the highest riboflavin productivity was obtained at pH value of 4.5 to 5.5 and morphological examination revealed that asci and ascospores outnumbered the other forms [26].

### *Ashbya gossypii*

Another riboflavin overproducer, *A. gossypii* was known to yield more riboflavin than *E. ashbyii* [27]. However, with similar constructs and equivalent yield, *E. ashbyii* showed neither nuclear fusion nor meiosis [28]. On the contrary, *E. ashbyii* showed instable in storage, for example during lyophilization or storage in slant at room temperature [29].

The ability of *A. gossypii* to overproduce riboflavin was discovered in 1946 [30]. The characteristics and fermentation of this strain are very similar to those of *E. ashbyii*. After the discovery of this yeast-like, filamentous fungus, the industrial producer of riboflavin has moved to *A. gossypii*, which has replaced *E. ashbyii* [31]. The first application of *A. gossypii* to industrial scale production was made in 1968 [2]. Since the first quantitative report [30] numerous studies have been carried out to overproduce riboflavin. In early studies, some defined media were established using Tween 80 and purine, and 145 mg/L and later 370 mg/L of riboflavin were produced in use of these media [32,33]. Many media were then developed [34] which contained glucose, vitamins, and amino acids, and mineral salts. In these markedly improved media, the production of riboflavin reached 1 g/L. It was found that glycine was an important component and Tween 80 served to prevent mycelium lysis and stimulator for riboflavin overproduction. Apart from these media, more complex media for industrial application were studied, and using soybean oil as a carbon source and collagenous protein and CSL as nitrogen sources, yields of 5 g/L were obtained [35,36]. The same amount of riboflavin was produced using other newly developed media, which included 3% bone

fat and 2% soybean oil as carbon sources, supplemented with additional soybean oil after 72 h in a 120 h cultivation [37]. Peptone and gelatin were investigated as complex nitrogen sources [35,38], and some waste proteins of animal origin, such as skin and bone glues were also tested [39]. Skin and bone glues contain several important amino acids for riboflavin production, such as glycine (about 20%) and threonine. These amino acids were found strongly stimulate riboflavin synthesis in *A. gossypii* [34,38,40], and in medium containing 3% skin glue as a nitrogen source, and 4% gelatin lard as a carbon source, to which 2.5% gelatin lard was added after 72 and 120 h of cultivation, the amount of riboflavin produced was higher than 5.5 g/L [39].

Microscopical characteristics, such as hyphae, asci, spore and autolysis was found to be varied in *A. gossypii* with various additional components and time courses [41]. Granula and hyaline droplets were described in the other study [42]. In 1990, *A. gossypii* was selected as a subject for fractal geometry, which at that time was being introduced to microbiology [43]. It was the first example of fractal aggregates in biological systems, with a cell as the smallest aggregating unit and the colony as an aggregate. Fractal dimensions increased during growth and demonstrated that the technique might serve as an important way of characterizing the morphology. The experimental results showed that fractal geometry provides a suitable method of describing the biological growth pattern of the *A. gossypii*. On the other hand, with morphological observations made using Nile red staining, found microdroplets in the hyphae of *A. gossypii* [44]. These lipid bodies accumulated neutral lipid as an energy reserve, and the fact was confirmed by separate experiments as follows; when soybean oil was used, the neutral lipid reached about 22% of the mycelium dry weight, while when the mycelium was grown on glucose it produced only 12% triacylglycerol, and these amounts decreased to 3–4% when the glucose declined in the medium. Although many fungi have been found to produce neutral lipid and utilize it, especially during spore germination [45], *A. gossypii* accumulated extraordinarily high amounts of neutral fatty acids. These results showed that the fat metabolism of *A. gossypii* was more efficient than that of using other fungi [44]. The yield enhancing effect of plant oil as a substrate in riboflavin production using *A. gossypii* had been shown earlier [2,25,36].

In terms of using plant oil as the sole carbon source in vitamin formation, the glyoxylate cycle is indispensable because the fatty acids can be degraded by isocitrate lyase, the key enzyme of this anaplerotic pathway, and isocitrate lyase from *A. gossypii* was firstly isolated and characterized [46]. During the production of riboflavin using *A. gossypii*, it was interesting that this enzyme appeared of the start of the riboflavin accumulation [47]. This implies that riboflavin synthesis might begin from storage lipids or this was observed with Nile red, to depended upon isocitrate lyase, which catalyzes the cleavage of isocitrate to glyoxylate and succinate. Itaconate as antimetabolite inhibits this pathway and

eventually riboflavin production. For this reason itaconate-resistant mutant grown on soybean oil had a significantly higher isocitrate lyase specific activity, which was accompanied by a large overproduction of riboflavin compared to the wild type strain [47]. Itaconate-resistant mutants were achieved easily using the characteristics of riboflavin. Because riboflavin is yellow and this can be used on agar plates to search for improved riboflavin overproducers, and this method led to *A. gossypii* to riboflavin overproducers being used for industrial applications [4]. When plant oils were used as a carbon source in riboflavin biosynthesis, lipase also becomes important, because the plant oils are degraded to fatty acids and glycerol by an extracellular lipase before their uptake into intracellular organelles. The fungal lipase of *A. gossypii* was hardly detected, because the activity was deactivated by interfacial characteristics [48]. Meanwhile, researches on osmoadaptation mechanisms [49] and vacuole of *A. gossypii* have been much studied [50].

#### Comparison of the Riboflavin Productivities of Several Microorganisms

In the field of microbial fermentation, there are several riboflavin-producing microorganisms, as shown in Table 1. Among these, the Gram-positive bacteria *Bacillus subtilis*, the yeast *Candida famate*, and the filamentous fungus *Ashbya gossypii* are mainly used for the biosynthesis of riboflavin via large-scale production [4]. *Candida* sp. is a natural overproducer of riboflavin, but these yeasts have to overcome iron inhibition in order to overbiosynthesize riboflavin [31]. *Candida famate* with tubercidine-, and iron-resistance has been developed to the level of plant operation, and it has been recently reported that production of riboflavin reaches to 20–30 g/L using this yeast [1,14]. Like *Candida* sp., bacteria are also inhibited by iron [31]. Among these *Bacillus subtilis* is mainly used to study riboflavin production, especially in terms of biosynthesis [51]. And a recombinant, riboflavin-producing strain of *Bacillus subtilis* was reported to allow productivities of 80 mg/L at 0.3 h<sup>-1</sup> using a glucose-limited chemostat [52]. To obtain riboflavin production using this Gram-positive bacterium requires at least the deregulation of purine synthesis and a mutation in a flavokinase/FAD-synthase. Currently recombinant *B. subtilis* with these features is in use in large-scale fermentations and producing concentrations exceeding 15 g/L [53].

*A. gossypii* as a strong overproducer and has been reported to give product yields in the range of 15–20 g/L [1,27]. About 30% of the world's industrial riboflavin output is produced by direct fermentation with *A. gossypii* [27], which has substituted for *E. ashbyii*, although its use in industry has been limited to date [21].

In addition, some experiments have been performed using a minor producer of riboflavin. *Candida guilliermondii*, which produced small amounts (220 mg/L) of riboflavin from liquid brewery waste [54] and *Mycobacterium pheli* also produced small quantities from

Table 1. Summary of the riboflavin productivities of several strains

	Strains	Carbon source	Nitrogen source	Supplements	Max. riboflavin concentration (g/L)	Culture time	References
Bacteria	<i>Clostridium acetobutylicum</i>	-	-	-	0.1	-	5
	<i>Bacillus subtilis</i>	Glucose	-	-	0.08	0.3 h	52
Yeast	<i>Candida flareri</i>	Glucose, fructose	Urea, glycin, serin, threonine	Guanine, xanthine	0.6	-	13
	<i>Candida guilliermondii</i>	Liquid brewery waste	-	biotin	0.2	3 day	54
	<i>Mycobacterium pheli</i>	Beet molasses	Peptone	-	0.1	6 day	55
	<i>Aspergillus terreus</i>	Beet molasses	Peptone	Asparagine	1.0	16 day	58
	<i>Eremothecium ashbyii</i>	Glucose, fructose, sucrose	Asparagine, serine, threonine	Biotin, thiamine, inositol	0.2	-	16, 17
Glucose		Caseine, glycine	Tween 80, lecithine	1.4	-	18	
Glucose, sucrose		Soybean meal	-	2.0	-	20	
Glucose, molasses		CSL, dry yeast, meat extract	-	1.0-2.0	4 day	22, 23	
Molasses		Peasnut seed cake	-	3.3	7 day	21	
Fungi		Glucose	Asparagine, glycine, tyrosine	Tween 80, hypoxanthine, biotin, thiamine, nositol	0.4-1.0	-	33, 34
	<i>Ashbya gossypii</i>	Corn oil, soybean oil	CSL, yeast extract, collagenous Proteins	-	5.0	-	35, 36
		Soybean oil, bone fat	CSL, gelatin	Glycine	5.0	8 day	37
		Soybean oil	CSL, gelatin, skin glue	Glycine	5.5	8 day	39

beet molasses [55]. A filamentous fungus, *Aspergillus terreus* was isolated from crude oil [56,57] and produced 1 g/L of riboflavin from beet molasses [58].

## RIBOFLAVIN BIOSYNTHESIS

There are three kinds of riboflavin overproducers, yeast, gram-positive bacteria, and fungus, the metabolic fluxes of which are shown in Fig. 1. *Candida famata* has a simple pathway from glucose to guanosine triphosphate (GTP), which is the entrance of riboflavin biosynthesis. The *Candida* uses glucose as its carbon source and converts this to ribulose-5-phosphate (Ribu-5P), which is a precursor of L-3, 4-dihydroxy-2-butanone-4-phosphate (DBP). 3-Phosphoglycerate, an intermediate of glycolysis, is also used for the synthesis of glycine that is the precursor of GTP.

*Bacilli* cells follow a different pathway from yeast, and convert glucose to guanosine monophosphate (GMP), which is also a precursor of GTP.

The most popular riboflavin overproducers are fungi, *A. gossypii* or *E. ashbyii*. Of these two strains, *Ashbya-*

*gossypii* specifically uses plant oil, which is obtained by decomposing fatty acids using extracellular lipase, and the resulting fatty acids enter to the peroxisomes. Biosynthesis begins from acetyl-CoA and forms malate, the precursor of GTP through several enzymatic reactions. On the other hand, citrate an intermediate of the TCA cycle in mitochondria is converted to isocitrate in peroxisomes. Isocitrate is partly converted to glyoxylate, and finally converted to malate.

The resulting metabolites from carbon sources in several riboflavin overproducers enter the riboflavin biosynthetic pathway. Biosynthesis starts from GTP and the final product is riboflavin; the precise fluxes are described as follows.

### Guanosine-triphosphate (GTP), as a Precursor of Riboflavin

Fig. 1 shows a proposed model of metabolic flux of riboflavin production in *Ashbya gossypii*. There are many steps from substrate to riboflavin, and the metabolic flux to riboflavin can be separated into specific




Fig. 1. Proposed model for the metabolic flux for riboflavin production by *Ashbya gossypii*, *Bacillus subtilis*, and *Candida famate*. Abbreviations: G-6P, Glucose-6-phosphate; 3PG, 3-phosphoglycerate; PEP, Phosphoenolpyruvate; Rib-5P, Ribulose-5-phosphate; OAA, Oxaloacetate; Asp, Aspartate; Thr, Thre-onine; Gly, Glycine; Ser, Serine; GTP, Guanosine triphosphate; GMP, Guanosine monophosphate; XMP, Xanthine monophosphate; IMP, Inosine monophosphate; DRTP, 2, 5-diamino-6-ribosylamino-4 (3H)-pyrimidinedione 5-phosphate; ARP, 5-amino-6-ribitylamino-2, 4 (1H, 3H)-pyrimidine; DBP, L-3, 4-dihydroxy-2-butanone-4-phosphate; DMRL, 6, 7-dimethyl-8-ribityllumazine.

and non-specific reactions. As shown in Fig. 1, most of the metabolites concentrate upon GTP, which is the first precursor of riboflavin. From this compound, six specific reaction steps are required to produce riboflavin. GTP as a purine, which was built de novo from amino acids, tetrahydrofolate derivatives, and CO<sub>2</sub> are well known to stimulate riboflavin overproduction in many organisms.

The sugar moiety is from ribose-5-phosphate from the pentose phosphate pathway. It has been shown that the 4 carbon, 5 carbon, and 7 nitrogen atoms of the purine ring originate from glycine [59]. Therefore, glycine stimulates riboflavin overproduction as a precursor of purine. Then results were obtained with *Candida* sp. [7,9,12] and *A. gossypii* [36]. Glycine in the riboflavin biosynthesis was not associated to cell growth, but only product formation [34], which is thought to be a limiting precursor in riboflavin overproduction.

In early work with *S. cerevisiae*, two pathways of glycine biosynthesis were described as follows [60]; one is a glycolytic pathway starting from the glycolytic intermediate, 3-phosphoglycerate on a glucose medium. The other is a gluconeogenic pathway starting from glyoxylate, a product of the anaplerotic glyoxylate cycle, on ethanol or acetate medium. Recent work [61] identi-




Fig. 2. Glycine biosynthesis in *S. cerevisiae*.

fied an additional glycine biosynthetic pathway, which starts from threonine and act as a major source of glycine in *S. cerevisiae* (Fig. 2).

Unlike *Candida* and *Ashbya*, *E. ashbyii* was not stimulated by glycine. Instead, serine and threonine served the same function [62]. It was elucidated that the two compounds are converted to riboflavin through glycine intracellularly [63, 64].

Purine, which is incorporated by glycine, stimulates riboflavin overproduction. Among this xanthine had been postulated for some times to be an intermediate precursor because of its structural similarities. But in the experiments with guanine auxotroph (lacking XMP aminase) *Aerobacter aerogenes* [65,66], *Candida guilliermondii* [67], and *Corynebacterium* sp. [68], it was

proven that the main precursor was a guanine or guanine nucleotide and the conversion of adenine, hypoxanthine and xanthine to riboflavin passed through a guanine or guanine nucleotide.

As research progressed, it was proposed that the guanine precursor is a nucleotide or nucleoside and the first definitive evidence of this was obtained with a mutant of *Salmonella typhimurium* [69]. Such results were also obtained with *E. ashbyii* [70-72] and *B. subtilis* [73]. Much work showed that GTP (I) is a real immediate precursor of riboflavin. The most conclusive evidence was found in *E. coli* [74], *B. subtilis* [75], the yeasts *Pichia guilliermondii* (*Candida guilliermondii*) [76], and *Torulopsis candida* (*Candida flareri*) [77].

### Which Compound Follows GTP?

During biosynthesis, the 8-carbon atom is eliminated as a single carbon unit. This result was obtained with *E. ashbyii* [78]. In the early studies, the compound following GTP was proposed to be 2,5-diamino-6-hydroxy-4-ribitylamino-pyrimidine, because this compound was accumulated in a riboflavin auxotroph of *S. cerevisiae* [79]. Through intensive researches it has recently been shown that this group is derived from the ribose moiety of GTP, which is incorporated without degradation into the ribityl side chain. These results were obtained with *Salmonella typhimurium* [69], *E. ashbyii* [72,80,81], *Pichia guilliermondii* [82, 83], and *B. subtilis* [73]. As mentioned above, the compound following GTP was suggested to be 2,5-diamino-6-hydroxy-4-ribitylamino-pyrimidine [79]. This compound was established to be 2,5-diamino-6-ribosylamino-4 (3H)-pyrimidinedione 5'-phosphate (DRTP) (II) in Fig. 3 [84] through 2,5-diamino-6-hydroxy-4-ribosylamino-pyrimidine 5'-phosphate [85].

### Two Different Steps to the Next Compound, 5-Amino-6-ribitylamino-2, 4 (1H, 3H)-pyrimidine (ARP)

In the early studies, the next compound was identified using auxotrophs of *Aspergillus nidulans* [86] and *Saccharomyces cerevisiae* [87] as 5-amino-2,6-dihydroxy-4-ribitylamino-pyrimidine. Another research group suggested that the compound was 5-amino-2,6-dioxy-4-ribitylamino-pyrimidine as the product was accumulated in resting cells of the *E. ashbyii* [88] and a riboflavin-deficient mutant of *B. subtilis* [89]. These compounds are at present known to as ARP (III) [84]. Between DRTP (II) to ARP (III), there are two steps including a deamination and a reduction as shown in Fig. 4. Reduction precedes deamination in one pathway and deamination occurs before reduction in the other pathway. First, reduction before deamination was found in a mutant *S. cerevisiae* [90]. Thereafter the same result was shown in *C. guilliermondii* [82, 84] and *A. gossypii* [91]. The second mechanism involving reduction after deamination shown to operate in *E. coli* [92] and *B. subtilis* [93].




Fig. 3. Biosynthesis from GTP (I) to DRTP (II). A denotes GTP cyclohydrolase II.




Fig. 4. Two pathways from DRTP (II) to ARP (III). B and C denote reductase and deaminase, respectively.




Fig. 5. The origin of the four-carbon moiety as a precursor of DMRL (IV). III and IV denote ARP and DMRL, respectively. DBP (VII) is synthesized by DBP synthase (F) from D-Ribu-5P (VI), one of glycolytic intermediates. D and E denote DMRL synthase and riboflavin synthase, respectively.

### Additional Four Carbon Moiety as a Precursor of Riboflavin

In riboflavin biosynthesis, the pathway from ARP (III) to 6,7-dimethyl-8-ribitylamino-pyrimidine (DMRL) (IV) in Fig. 5 was the most recently elucidated because of the origin of the four carbon moiety. Significant work was performed on the origin of four carbons. The likely origin of the four carbons was ARP (III). Even though much investigating work was performed, the origin of the 4-carbon moiety remained a mystery for a long time. At one time, it was seriously considered that acetoin might be the source of the four carbons from work upon *E. ashbyii* [94,95], and *C. flareri* [96]. But neither acetoin nor acetate is found to be precursors of the dimethylbenzene unit of riboflavin in *E. ashbyii* by labeling experiments [97]. Subsequently its oxidized form, diacetyl was considered because it was known to

condense nonenzymatically with ARP (III) to form DMRL (IV) [95]. This hypothesis was supported by other experiments with ethanol [98]. Feeding ethanol during the fermentation of *A. gossypii* showed effective stimulation of riboflavin formation. Ethanol was not considered to be directly utilized to form riboflavin, but to be metabolized to form diacetyl through acetaldehyde, pyruvate, and acetoin. Therefore, it was believed that ethanol might play a significant role in the origin of the four carbon unit, but ethanol is currently believed to be precursor of acetyl-CoA that proceeds to isocitrate through the gluconeogenic pathway. In addition, the incorporation of labeled diacetyl into riboflavin was not sufficient proof of its role as a natural precursor [99]. The stimulation of riboflavin production by adding diacetyl can be explained by another reason, which is that it acts like ethanol rather than as a precursor [100].

On the other hand, glucose and ribose were suggested to be the true donors of the 4-carbon moiety [101], because they incorporated more efficiently than either acetate or pyruvate. Some of the data obtained from measurements of the incorporation of various radioactive compounds were consistent with the view that a pentose compound was the source of the four-carbon unit [99]. Tetraose, formed via the pentose phosphate pathway, was also reported to serve as the precursor of the four-carbon biosynthetic unit [102]. Through various experiments, considerable progress on the origin of the four-carbon moiety was made in studies using *A. gossypii* [103]. It was shown in *in vivo* studies with a variety of  $^{13}\text{C}$ -labeled precursors that the C-6 $\alpha$ , C-6, C-7, and C-7 $\alpha$  of DMRL are biosynthetically equivalent to C-1, C-2, C-3, and C-5 of the pentose phosphate pool. C-4 of the pentose phosphates has no equivalent in lumazine. It was shown that this carbon atom is eliminated by an intramolecular skeletal rearrangement [104]. Further studies revealed the same results with *C. guilliermondii* [105] and *B. subtilis* [106]. Several pentose and pentulose phosphates could serve as the second substrate in the reaction to form lumazine. That is, the formation of the four-carbon moiety was determined by the loss of the internal carbon atom 4 from a pentose precursor.

On the other hand, the enzyme that catalyzed the formation of DBP (VII) from D-Ribu-5P (VI) was reported [107] from *C. guilliermondii*. DBP could yield the lumazine with ARP (III). Therefore, this compound was thought to be an obligatory intermediate of riboflavin biosynthesis (Fig. 5).

### Conversion of DMRL to Riboflavin

As the nearest precursor of riboflavin (V), DMRL (IV) in Fig. 5 was identified during the early research period [96,108]. Our understanding of the transformation step between DMRL and riboflavin was elucidated by experiments with *A. gossypii* and *E. ashbyii* [109]. It became apparent that two moles of DMRL react to give one mole of riboflavin and one mole of ARP (III). The




Fig. 6. Bioconversion of DMRL (IV) to riboflavin (V), which is catalyzed by riboflavin synthase (E).

regiospecificity of the reaction have been elucidated by experiments with [7- $^2\text{H}_3$ ] DMRL as substrate [110]. Riboflavin formed from labeled substrate contained deuterium in positions 9 and 7 $\alpha$ . Thus, the two four carbon moieties ultimately form the xylene ring (Fig. 6). This regiospecificity was later confirmed by experiments with  $^{13}\text{C}$ -labeled substrates [111] and deuterium-based experimentation [112]. The other product of the reaction, ARP (III), was suggested to be reused to form another molecule of DMRL (IV) [113], but no firm evidence exists for enzymatic reutilization except for the results obtained with *E. ashbyii* [114], which showed that the ARP (III) was reused, at least in part. At present, it has been established that the by-product, ARP, can enter the biosynthesis of riboflavin (V) as a precursor because the by-product is also the same biosynthetically and structurally to produced compound from DRTP (II).

As mentioned above, the starting materials of riboflavin biosynthesis are GTP (I) and Ribu-5P (VI), and there are many intermediates on the pathway. On the other hand, several enzymes are concerned in these reactions, and these are described in the following section.

### Enzymes Related to Riboflavin Biosynthesis and Regulations of GTP Cyclohydrolase II

On the riboflavin biosynthetic pathway, the starting material, GTP (I) is catalyzed by GTP cyclohydrolase II and forms formate and DRTP (II) as shown in Fig. 3. The enzyme was found and reported initially a few experiment with *E. coli* [74] and was suggested to be GTP cyclohydrolase II because it differed from a previously studied *E. coli* enzyme known to be involved in folic acid biosynthesis. GTP cyclohydrolase II catalyzes the initial and rate-limiting step of riboflavin biosynthesis and is subject to multiple metabolic controls. Much work has been performed upon the properties of GTP

cyclohydrolase II of *B. subtilis* [115,116] and *E. coli* [117]. In particular, enzyme of *B. subtilis* was a bifunctional enzyme that could also catalyze the formation of 3,4-dihydroxy-2-butanone 4-phosphate from ribulose 5-phosphate [118], and the GTP cyclohydrolase II of *E. coli* and *B. subtilis* was specified by the gene *ribA*. Recently, the gene *RIB1* in *S. cerevisiae*, which encodes GTP cyclohydrolase, was cloned by functional complementation of the corresponding mutation [119,120]. Contemporaneously the structure gene *RIB1* encoding GTP cyclohydrolase II of *Pichia guilliermondii* has been cloned by functional complementation of a *ribA* mutation in *E. coli* [121]. The deduced protein sequence obtained showed significant homologies to GTP cyclohydrolase from *S. cerevisiae* and its C-terminal half homologies to the enzyme of *E. coli* and *B. subtilis* [122].

### Reductase and Deaminase

The existence of two enzymes involved in the biosynthesis of riboflavin was reported in *E. coli* [92]. These two enzymes catalyzed the conversion reaction of the product by GTP cyclohydrolase II and DRTP (II), to ARP (III). There are two enzymatical steps between the two compounds. First, deamination at carbon 2 of the ring by deaminase was by-passed and instead the ribosylamino group was reduced to a ribitylamino group by reductase. These results are in general agreement with those of *S. cerevisiae* [123]. The major difference is that in yeast, reduction precedes deamination, whereas in *E. coli* deamination occurs first, as shown in Fig. 4. Reductase activity was also shown in the filamentous fungus, *A. gossypii* [91] and in the yeast, *C. guilliermondii* [84].

In the meanwhile, reductase and deaminase in yeast, *S. cerevisiae* [124], is specified by the genes *RIB7* and *RIB2*, respectively, and in other yeast, *Pichia guilliermondii* (*Candida guilliermondii*) [122], by *RIB2* and *RIB3*, respectively, but the yeast enzymes have not been studied in detail. On the contrary, the complete sequence of the *B. subtilis* [53,115] and *E. coli* [125] riboflavin operon has been determined. In *B. subtilis*, it is known that the gene *ribG* encodes deaminase and that the gene *ribT* encodes reductase [126], but further study concluded that the *ribG* gene could code for a bifunctional deaminase and reductase [127], which was arrived at by comparing the homology between the 3' end of the *ribG* gene of *B. subtilis* and the *RIB7* gene of *S. cerevisiae* coding for reductase. An *E. coli* gene with sequence homology to the *B. subtilis* *ribG* gene has also been identified as *ribD* [127,128]. In common with *ribG* of *B. subtilis*, *ribD* of *E. coli* codes for a bifunctional deaminase and reductase in regular order.

### DMRL Synthase or the $\beta$ -Subunit of Riboflavin Synthase

Lumazine, DMRL (IV) is biosynthesized from ARP (III) and DBP (VII) by the enzyme lumazine synthase,

as shown in Fig. 5. The gene coding for lumazine synthase in *S. cerevisiae* has been cloned as *RIB4* [129]. Gene *RIB4* is a single copy gene located on the left arm of chromosome XV as indicated by hybridization analysis and its deduced amino acid sequence showed extensive homology to the sequence of the  $\beta$ -subunits of riboflavin synthase from *B. subtilis*. Lumazine synthase occurs as a constituent of the enzyme complex in *B. subtilis*, which consists of 60  $\beta$ -subunits and 3  $\alpha$ -subunits [130]. The  $\beta$ -subunits indicating lumazine synthase form a capsid with icosahedral 532 symmetry [131,132] and is specified by the gene *ribH* of the riboflavin operon on the *B. subtilis* chromosome [126]. The  $\alpha$ -subunits, another component of the riboflavin synthase enzyme complex, formed a trimer that is enclosed in the central core of the icosahedral  $\beta$ -subunit capsid. Lumazine synthase of *E. coli* is specified by a gene designated *ribE*. The protein is an empty icosahedral capsid consisting of 60 subunits. In contrast to the homologous *B. subtilis* protein, it does not form a complex with riboflavin synthase [133].

### Riboflavin Synthase or the $\alpha$ -Subunits of Riboflavin Synthase

The terminal step of riboflavin biosynthesis is catalyzed by riboflavin synthase. This enzyme catalyzes the dismutation of two molecules of the substrate DMRL (IV) to yield one molecule each of riboflavin (V) and ARP (III) as shown in Fig. 6. As mentioned-above, *B. subtilis* has two different riboflavin synthases characterized by  $\alpha$ -subunits and  $\beta$ -subunits. The enzyme catalyzing the last step of riboflavin biosynthesis is specified by the gene *ribB* of the riboflavin operon on the *B. subtilis* [126]. The derived amino acids sequence showed internal homology existed between the N-terminal and C-terminal halves of the protein, which suggests that the promoter forms two structurally similar substrate-binding domains [134]. Because, the enzyme binds two molecules of the substrate DMRL, during enzymatic riboflavin production, one mole of the substrate DMRL acts as the donor and the other, as the acceptor of the four-carbon moiety. These two sites are distinguished as the acceptor site binds the product riboflavin, whereas the donor site binds the 2nd product 5-amino-2, 6-dihydroxy-4-ribitylamino-pyrimidine [135,136]. A gene *ribC* encoding mono functional riboflavin synthase in *E. coli* was specified [137] and cloned onto a 6kb DNA segment by marker rescue using a *ribC* mutant of *E. coli*. In common with *E. coli*, *S. cerevisiae* has also a monofunctional riboflavin synthase. The gene *RIB5* encoding this enzyme was isolated from a yeast genomic library by functional complementation of a mutant, *rib5-10*, lacking riboflavin synthase activity [138]. The deduced amino acid sequence showed extensive homology to the sequence of the  $\alpha$ -subunits of riboflavin synthase of *B. subtilis*. These internal homologous characteristics suggested that the Rip5p subunit contained two structurally related but catalytically different domains.



## DBP Synthase

The xylene ring of riboflavin originates by dismutation of the precursor, DMRL (IV). The formation of the latter compound requires a four-carbon unit. DBP (VII) as the four-carbon unit incorporated into riboflavin (V) [107,139]. Ribulose 5-phosphate (VI) served as substrate for the formation of DBP (VII) as catalyzed by the enzyme DBP synthase (Fig. 5). The enzyme catalyzes the release of carbon 4 of ribulose 5-phosphate as formate, which is accompanied by a complex rearrangement reaction conducive to the formation of DBP from carbon atoms 1,2,3 and 5 of the substrate [139].

As mentioned above in GTP cyclohydrolase II, the *ribA* encoding GTP cyclohydrolase II in *B. subtilis* could be also specify DBP synthase bifunctionally [118]. In contrast to *B. subtilis*, *E. coli* has two separate genes, *ribB* and *ribA*, that encode the function of DBP synthase [118] and GTP cyclohydrolase II [117], respectively. The *RIB3* gene of *S. cerevisiae* specifying synthase was cloned by the same method that was used to clone the *ribB* in *E. coli* [140].

## GENETICS FOR *A. GOSSYPII* AS AN OVERPRODUCER OF RIBOFLAVIN

In riboflavin biosynthesis, there are six specific enzymatic pathways (Fig. 1). However, the metabolic flux to riboflavin is also determined by numerous nonspecific reactions. This means that riboflavin production may be enhanced by supplementation. Ribitol [141], purine [34], or glycine [31,34,36] was studied as enhancers in the early stages of riboflavin production. Among these, glycine is thought to be a limiting precursor in riboflavin overproduction. Glycine supplementation during fermentation with *A. gossypii* doubled riboflavin production at least without growth variations. As mentioned in the riboflavin biosynthesis, there are three routes of glycine enzymatic biosynthesis, among these, serine hydroxymethyltransferase is the most widespread (Fig. 2). Another enzyme, glutamate glyoxylate aminotransferase (Fig. 2) was found in the mitochondria of a fungus. Thus this enzyme is not co-located with isocitrate lyase in the peroxisomes, which is a main location site of riboflavin metabolic flux. Glycine biosynthesis was discovered in *S. cerevisiae* through the threonine aldolase and later it was understood to be the main reaction [61,142]. Therefore, the gene *GLY1* encoding the threonine aldolase of *A. gossypii* was isolated by heterologous complementation of the glycine-auxotrophic *S. cerevisiae* [143]. In contrast to *S. cerevisiae* the *GLY1* knockout mutant of *A. gossypii* was not auxotrophic for glycine. This means that threonine aldolase plays only a minor role during glycine biosynthesis in *A. gossypii*. Nevertheless, overexpression of *AgGLY1* under the control of the constitutive *TEF* promoter and terminator led to a 10-fold increase of threonine aldolase specific activity and a 9-fold increase in riboflavin production when the medium was sup-

plemented with threonine. This strong enhancement could not be achieved by supplementation of glycine alone.

In terms of the metabolism of riboflavin, the glyoxylate cycle as well as glycine biosynthesis is important, because the glyoxylate cycle is essential for the degradation of plant oil in riboflavin production. There are degraded to fatty acids and glycerol by an extracellular lipase [48] and are the preferred substrates for riboflavin biosynthesis because of their yield-enhancing effects [25]. Since the two starting precursors of riboflavin are originated from carbohydrate metabolism, the glyoxylate cycle in cooperation with gluconeogenesis plays an indispensable function in terms of growth and riboflavin production in *A. gossypii*. Therefore, in this metabolic pathway, enzymatic activity of isocitrate lyase, which catalyzes the cleavage of isocitrate to glyoxylate and succinate through a carbon-conserving pathway, is a key point. Recently, a correlation between isocitrate lyase activity and riboflavin production was studied [47] and the isocitrate lyase-encoding gene *AgICL1* was isolated from *A. gossypii* by heterologous complementation of *S. cerevisiae* *icl1d* mutant [144]. In contrast to *S. cerevisiae* *ICL1* that is located in the cytosol [145, 146], *AgICL1* is located in the peroxisomes. *AgICL1* was subject to glucose repression and derepressed by glycerol, and was also in part induced by ethanol and acetate and fully induced by soybean oil. These results agree well with the result that soybean oil is an enhanced carbon source in riboflavin production by *A. gossypii*.

On the other hand, using a molecular approach, a study of the structural gene of *A. gossypii* was also undertaken [50]. The structural gene *VMA1* encoding vacuolar ATPase subunit A was cloned. Disruption of this gene of *A. gossypii* could lead to the complete excretion of riboflavin into medium, instead of its retention in vacuolar compartments. Because riboflavin production by *A. gossypii* starts in the late growth phase when septa and vacuoles are formed in the hyphae and concomitantly, vacuoles begin to accumulate large amounts of riboflavin. Therefore *A. gossypii* lacking the intracellular vacuoles can produce more riboflavin without experiencing lethal effects, like *S. cerevisiae*. The same results were obtained with concanamycin A, which inhibits the vacuolar ATPase, which means that riboflavin accumulation to vacuoles depends on the activity of the ATPase.

Summarizing from the genetic viewpoint, the *AgICL1* encoding isocitrate lyase is thought to be a key gene in riboflavin production and the *VMA1* gene disruptant can cause the excretion of riboflavin extracellularly. The former has the ability to promote the main metabolic flux in the total metabolism, the latter has the ability to excrete the final product into the broth.

## CONCLUSION

Recently the riboflavin production is shifting from

chemical synthesis to microbial production. The microbial production of riboflavin was reviewed with the viewpoints of biosynthesis, key enzymes, and enzyme related genes. In the metabolic viewpoint for riboflavin overproduction the supplementation method of GTP, a limiting precursor should be developed. Especially in the case of fungi using vegetable oil as a carbon source, the GTP fluxes from glyoxylate cycle, in which case the key enzymes in the glyoxylate cycle need to be strengthened. The research on a gene that is related with key enzyme, *AgICL1* is carrying out to produce riboflavin efficiently. With the progress of these metabolic, biosynthetic, and genetic developments the riboflavin production will be completely replaced by microbial production in a near future.

## REFERENCES

- [1] Demain, A. L. (1999) [Online] Stunning achievements of industrial microbiology. <http://www.asmsa.org/memonly/asmnews/may99/feature7.html>.
- [2] Lago, B. D. and L. Kaplan (1981) Vitamin fermentations B<sub>2</sub> and B<sub>12</sub>, pp. 241-242. In: M. Moo Young (eds.). *Advances in Biotechnology*: vol. 3, Pergamon, Toronto, Canada.
- [3] Vandamme, E. J. (1992) Production of vitamins, coenzymes and related biochemicals by biotechnological processes. *J. Chem. Tech. Biotechnol.* 53: 313-327.
- [4] Stamann, K. -P., J. L. Revuelta, and H. Seulberger (2000) Three biotechnical processes using *Ashbya gossypii*, *Candida famate*, or *Bacillus subtilis* compete with chemical riboflavin production. *Appl. Microbiol. Biotechnol.* 53: 509-516.
- [5] Hickey, R. J. (1945) The inactivation of iron by 2,2'-bipyrimidine and its effect on riboflavin synthesis by *Clostridium acetobutylicum*. *Arch. Biochem.* 8: 439-447.
- [6] Tanner, F. W. Jr., Ch. Vajnovich, and J. Van Lanen (1945) Riboflavin production by *Candida species*. *Science* 101: 180-181.
- [7] Goodwin, T. and D. McEvoy (1959) General factors controlling flavinogenesis in the yeast *Candida flareri*. *Biochem. J.* 71: 742-748.
- [8] Burkholder, P. R. (1943) Synthesis of riboflavin by yeast. *Proc. Nat. Acad. Sci. USA.* 29: 166-172.
- [9] Burkholder, P. R. (1943) Influence of some environmental factors upon the production of riboflavin by yeast. *Arch. Biochem.* 3: 121-130.
- [10] Enari, T. M. (1955) Effect of cobalt and iron on riboflavin production by *Candida guilliermondia*. *Acta Chem. Scand.* 9: 1726-1727.
- [11] Takao, S (1964) Riboflavin biosynthesis by *Candida robusta*; Part III. Effect of metal ion and necessity of CaCO<sub>3</sub>. *Agr. Biol. Chem.* 28: 765-769.
- [12] Giri, K. V. and P. R. Krishnaswamy (1954) The synthesis of riboflavin by a mutant yeast, *Saccharomyces cerevisiae*. *J. Bacteriol.* 67: 309-332.
- [13] Levine, H., J. E. Oyaas, L. Wasserman, J. C. Hoogerheide, and R. M. Stern (1949) Riboflavin production by *Candida yeasts*. *Ind. Eng. Chem.* 41: 1665-1668.
- [14] Heefner, D., C. A. Weaver, M. J. Yarus, L. A. Burdzinski, D. C. Gyure, and E. W. Foster (1988) Riboflavin producing strains of microorganisms, method for selecting, and method for fermentation. Patent WO 88/09822.
- [15] Guilliermond, A., M. Fontaine, and A. Raffy (1935) Sur l'Existence dans l'*Eremothecium Ashbyii* d'un pigment jaune se rapportant au groupe des flavines. *C. R. Acad. Sci.* 201: 1077-1080.
- [16] McLaren, J. A. (1952) The effects of certain purines and pyrimidines upon the production of Riboflavin by *Eremothecium ashbyii*. *J. Bacteriol.* 63: 233-241.
- [17] Yaw, K. E. (1952) Production of riboflavin by *Eremothecium ashbyii* growth in a synthetic medium. *Mycologia.* 44: 307-317.
- [18] Hickey, R. J. (1953) Some nutritional requirements for biosynthesis of riboflavin by *Eremothecium ashbyii*. *J. Bacteriol.* 66: 27-33.
- [19] Beesch, S. C. and B. N. Frazer (1953) Riboflavin by microbiology fermentation. To publiker Industries Inc. US patent 2.647, 704.
- [20] Stepanov, A. I., M. Yu. Beburow, and V. G. Zhdanov (1974) Mutants of *Eremothecium ashbyii* resistant to 8-azaguanine; Communication I. Isolation of mutants and study of the level of riboflavin biosynthesis. *Sov. Genet.* 8: 729-733.
- [21] Kalingan, A. E. and M. R. V. Krishnan (1997) Application of agro-industrial by-products for riboflavin production by *Eremothecium ashbyii* NRRL 1363. *Appl. Microbiol. Biotechnol.* 47: 226-230.
- [22] Vaghy, T. (1971) Correlation between the morphological variability and riboflavin producing capacity of *Eremothecium ashbyii*. *Acta Microbiol. Acad. Sci. Hung.* 18: 75-80.
- [23] Stepanov, A. I., M. Yu. Beburow, and V. G. Zhdanov (1974) The use of mutagenic factors in the selection of the riboflavin producer *Eremothecium ashbyii*. *Sov. Genet.* 8: 745-749.
- [24] Özbas, T. and T. Kutsal (1986) Riboflavin production by *Eremothecium ashbyii* in a batch stirred tank fermentor. *Biotechnol. Lett.* 8: 441-444.
- [25] Özbas, T. and T. Kutsal (1986) Comparative study of riboflavin production by two organisms *Eremothecium ashbyii* and *Ashbya gossypii*. *Enzyme Microb. Technol.* 3: 593-596.
- [26] Kolonne, S., R. J. Seviour, and B. M. McDougall (1994) Effect of pH on exocellular riboflavin production by *Eremothecium ashbyii*. *Biotechnol. Lett.* 16: 79-84.
- [27] Bigelis, R. (1989) Industrial products of biotechnology: application of gene technology, pp. 243. In: H. J. Rehm, and G. Reed (eds.). *Biotechnology*, vol. 7b, Weinheim: VCH.
- [28] Batra, L. R. (1973) Nematosporeaceae (Hemiascomycetidae) taxonomy, pathogenicity distribution and vector relations. *USDA Technical Bull.* 1469. 8: 71-74
- [29] Moore, H. N., G. de Becze, and E. Schraffenberger (1947) Studies on *Eremothecium ashbyii*. *J. Bacteriol.* 53: 502.
- [30] Wickerham, L. J., M. H. Flickinger, and R. M. Johnston (1946) The production of riboflavin by *Ashbya gossypii*. *Arch. Biochem.* 9: 95-98.
- [31] Demain, A. L. (1972) Riboflavin oversynthesis. *Annu.*

- Rev. Microbiol.* 26: 369-388.
- [32] Goodman, J. J. and R. R. Ferrera (1954) Synthesis of riboflavin by *Ashbya gossypii* grown in a synthetic medium. *Mycologia*. 46: 556.
- [33] Smith, C. G., G. A. Smith, and Z. Papadoupoulou (1961) Effect of surface active agents on the biosynthesis of riboflavin by *Ashbya gossypii*. *Biochem. Biophys. Acta*. 47: 344-349.
- [34] Kaplan, L. and A. L. Demain (1970) Nutritional studies on riboflavin overproduction by *Ashbya gossypii*, pp. 137-159. In: D. G. Ahearn (eds.). *Recent Trends in Yeast Research*. Georgia State Univ., Atlanta, GA, USA.
- [35] Malzahn, R. C., R. F. Philips, and A. M. Hanson (1963) The production of riboflavin by *Ashbya gossypii*. *Bacteriol. Proc.* 21.
- [36] Hanson, A. M. (1967) Microbial production of pigments and vitamins. pp. 222-250. In: H. J. Pepler (eds.). *Microbial Technology*. Reinhold, USA.
- [37] Szczesniak, T., L. Karabin, M. Szczepankowska, and K. Wituch (1971) Biosynthesis of Riboflavin by *Ashbya gossypii*. Part I. The influence of fats of the animal origin on the riboflavin production. *Acta Microbiologica Polonica Ser. B*. 3: 29-34.
- [38] Chiao, J., Y. Gwan, Y. Shen, W. Chen, C. Lang, and H. Wang (1960) Studies on the biosynthesis of riboflavin by *Ashbya gossypii* I. Factors affecting the high yield of riboflavin. *Acta Biol. Exp. Sinica* 7: 161.
- [39] Szczesniak, T., L. Karabin, M. Szczepankowska, and K. Wituch (1971) Biosynthesis of Riboflavin by *Ashbya gossypii*. Part II. The influence of animal proteins on the riboflavin production. *Acta Microbiologica Polonica Ser. B*. 3: 91-95.
- [40] Plaut, G. W. E. (1961) Water-soluble vitamins, Part II. Folic acid, Riboflavin, Thiamine, Vitamin B<sub>12</sub>. *Ann. Rev. Biochem.* 30: 409-446.
- [41] Cerletti, P., R. Strom, M. G. Giordano, D. Barra, and S. Giovenco (1965) Flavin coenzymes, flavinogenesis and reproduction in *Ashbya gossypii*. *J. Biochem.* 57: 773-786.
- [42] Mukerji, K. G. (1968) C.M.I. Descriptions of pathogenic fungi and bacteria No. 185. *Commonwealth Agricultural Bureaux*. The Eastern Press, London, UK.
- [43] Obert, M., P. Peeifer, and M. Sernetz (1990) Microbial growth patterns described by fractal geometry. *J. Bacteriol.* 172: 1180-1185.
- [44] Stahmann, K.-P., C. Kupp, S. D. Feldmann, and H. Sahn (1994) Formation and degradation of lipid bodies found in the riboflavin-producing fungus *Ashbya gossypii*. *Appl. Microbiol. Biotechnol.* 42: 121-127.
- [45] Weber, D. J. (1980) Lipid metabolism during fungal development. pp. 300-325. In: J. D. Weete (eds.). *Lipid Biochemistry of Fungi and Other Organisms*, Plenum, NY, USA.
- [46] Schmidt, G., K.-P. Stahmann, and H. Sahn (1996) Inhibition of purified isocitrate lyase identified itaconate and oxalate as potential antimetabolites for the riboflavin overproducer *Ashbya gossypii*. *Microbiology* 142: 411-417.
- [47] Schmidt, G., K.-P. Stahmann, B. Kaesler, and H. Sahn (1996) Correlation of isocitrate lyase activity and riboflavin formation in the riboflavin overproducer *Ashbya gossypii*. *Microbiology*. 142: 419-426.
- [48] Stahmann, K.-P., T. Bötdecker, and H. Sahn (1997) Regulation and properties of a fungal lipase showing interfacial inactivation by gas bubbles, or droplets of lipid or fatty acid. *Eur. J. Biochem.* 244: 220-225.
- [49] Föster, C., S. Marienfeld, V. F. Wendisch, and R. Krämer (1998) Adaptation of the filamentous fungus *Ashbya gossypii* to hyperosmotic stress: different osmoresponse to NaCl and mannitol stress. *Appl. Microbiol. Biotechnol.* 50: 219-226.
- [50] Föster, C., S. Marienfeld, R. Wilhelm, and R. Krämer (1998) Organelle purification and selective permeabilisation of the plasma membrane: two different approaches to study vacuoles of the filamentous fungus *Ashbya gossypii*. *FEMS Microbiol. Lett.* 167: 209-214.
- [51] Bacher, A., M. Fisher, K. Kis, K. Kugelbrey, S. Mörtl, J. Scheuring, S. Weinkauff, S. Eberhardt, K. Schmidt-Bäse, R. Hurber, K. Ritsert, M. Cushman, and R. Ladenstein (1996) Biosynthesis of riboflavin: structure and mechanism of lumazine synthase. *Biochem. Soc. Trans.* 24: 89-94.
- [52] Sauer, U., V. Hatzimanikatis, H.-P. Hohmann, M. Manneberg, A. P. G. M. Loon, and J. E. Bailey (1996) Physiology and metabolic fluxes of wild-type and riboflavin-producing *Bacillus subtilis*. *Appl. Environ. Microbiol.* 62: 3687-3696.
- [53] Perkins, J. B., J. G. Pero, and A. Sloma (1991) Riboflavin overproducing strains of bacteria. European patent application 0-405-370-A1.
- [54] Contasti, V. and S. Bahar (1988) Riboflavin production by *Candida guilliermondii* from liquid brewery waste. *Acta Cient. Venez.* 39: 69-74.
- [55] Ghazlan, H. A. (1994) Utilization of beet molasses for riboflavin production by *Mycobacterium phlei*. *J. Basic Microbiol.* 34: 157-162.
- [56] Sabry, S. A., A. H. El-Refai, and S. Y. Gamati (1989) Riboflavin formation by mould fungi cultivated on hydrocarbon-containing media. *Microbios.* 57: 33-40.
- [57] Omar, S. H. and Z. A. Olama (1985) Utilization of hydrocarbons by some fungi isolated from El-Alameen crude oil refineries. *Commun. Sci. Devel. Res.* 12: 1.
- [58] Sabry, S. A., K. M. Ghanem, and H. A. Ghazlan (1993) Riboflavin production by *Aspergillus terreus* from beet-molasses. *Microbiologia* 9: 118-124.
- [59] Stryer, L. (1995) Biosynthesis of nucleotides: Biochemistry. 4th ed., pp. 739-740. W. H. Freeman and Co., NY, USA.
- [60] Ulane, R. and M. Ogur (1972) Genetic and physiological control of serine and glycine biosynthesis in *Saccharomyces*. *J. Bacteriol.* 109: 34-43.
- [61] McNeil, J. B., E. M. McIntosh, B. V. Taylor, F. Zhang, S. Tang, and A. L. Bognar (1994) Cloning and molecular characterization of three genes, including two genes encoding serine hydroxymethyltransferase, whose inactivation is required to render yeast auxotrophic for glycine. *J. Biol. Chem.* 269: 9155-9165.
- [62] Goodwin, T. W. and S. Pendlington (1954) Studies on the biosynthesis of riboflavin. Nitrogen metabolism and flavinogenesis in *Eremothecium ashbyii*. *Biochem. J.* 57: 631-641.

- [63] Goodwin, T. W. and O. T. G. Jones (1956) Studies on the biosynthesis of riboflavin. 3. The utilization of  $^{14}\text{C}$ -labelled serine for riboflavin biosynthesis by *Eremothecium ashbyii*. *Biochem. J.* 64: 9-13.
- [64] Goodwin, T. W. and A. A. Horton (1960) Studies on flavinogenesis. 6: The role of threonine in riboflavin biosynthesis in *Eremothecium ashbyii*. *Biochem. J.* 75: 53-57.
- [65] Bacher, A. and F. Lingens (1969) The structure of the purine precursor in riboflavin biosynthesis. *Angew. Chem. Int. Ed. Engl.* 8: 371-372.
- [66] Bacher, A. and B. Mailänder (1973) Biosynthesis of riboflavin. The structure of the purine precursor. *J. Biol. Chem.* 248: 6227-6231.
- [67] Shavlovskii, G. M., L. P. Strugovshchikova, and E. M. Logvinenko (1971) Flavinogenesis in guanine-dependent mutants of *Candida guilliermondii*. *Dokl. Akad. Nauk. SSSR.* 196: 701-704.
- [68] Baugh, C. M. and C. L. Krumdieck (1969) Biosynthesis of the purine precursor of riboflavin in *Corynebacterium*. *J. Bacteriol.* 98: 1114-1119.
- [69] Mailänder, B. and A. Bacher (1976) Biosynthesis of riboflavin. Structure of the purine precursor and origin of the ribityl side chain. *J. Biol. Chem.* 251: 3623-3628.
- [70] Mitsuda, H. and K. Nakajima (1973) Effects of 8-azaguanine on riboflavin production and on the nucleotide pools in non-growing cells of *Eremothecium ashbyii*. *J. Nutr. Sci. Vitaminol.* 19: 215-227.
- [71] Mitsuda, H. and K. Nakajima (1975) Guanosine nucleotide precursor for flavinogenesis of *Eremothecium ashbyii*. *J. Nutr. Sci. Vitaminol.* 21: 331-345.
- [72] Mitsuda, H. and K. Nakajima (1975) The immediate nucleotide precursor, guanosine triphosphate, in the riboflavin biosynthetic pathway. *J. Nutr. Sci. Vitaminol.* 23: 23-24.
- [73] Bresler, S. E., G. F. Gorinchuk, T. P. Chernik, and D. A. Perumov (1978) Operon of riboflavin biosynthesis in *Bacillus subtilis*. XV. A study of mutants related to the initial stages of biosynthesis. The origin of the ribityl chain of the riboflavin molecule. *Genetika* 14: 2082-2090.
- [74] Foor, F. and G. M. Brown (1975) Purification and properties of guanosine triphosphate cyclohydrolase II from *Escherichia coli*. *J. Biol. Chem.* 250: 3545-3551.
- [75] Bresler, S. E. and D. A. Perumov (1979) Study of the operon for riboflavin biosynthesis in *Bacillus subtilis*. Influence of genotype on regulating the synthesis of GTP-5-triphosphate cyclohydrolase. *Genetika* 15: 967-971.
- [76] Shavlovskii, G. M., E. M. Logvinenko, V. E. Kashchenko, L. V. Kolton, and A. E. Zakal'skii (1976) Determination of GTP-cyclohydrolase, the first enzyme of riboflavin biosynthesis from the yeast *Pichia guilliermondii*. *Dokl. Akad. Nauk. SSSR.* 230: 1485-1487.
- [77] Shavlovskii, G. M., V. E. Kashchenko, L. V. Kolton, E. M. Logvinenko, and A. E. Zakal'skii (1977) Regulation of synthesis of GTP-cyclohydrolase participating in yeast flavinogenesis by iron. *Mikrobiologiya* 46: 578-580.
- [78] Hayes, D. H. and G. R. Greenberg (1969) Conversion of C-8 of guanine into a one carbon unit during riboflavin biosynthesis. *Bull. Soc. Chim. Biol.* 51: 1187-1198.
- [79] Bacher, A. and F. Lingens (1970) Formation of 2,5-diamino-6-hydroxy-4-(1'-D-ribitylamino) pyrimidine in a riboflavin auxotroph. *J. Biol. Chem.* 245: 4647-4652.
- [80] Mitsuda, H., K. Nakajima, and T. Nadamoto (1976) Nucleotide precursor in riboflavin biosynthesis. *J. Nutr. Sci. Vitaminol.* 22: 477-480.
- [81] Mitsuda, H., K. Nakajima, and Y. Yamada (1977) Studies on the intermediates in the biosynthetic pathway of riboflavin. I. Identification of a green fluorescent compound, Compound G<sub>1</sub>, accumulated in non-growing cells of *Eremothecium ashbyii* by the addition of dimeric diacetyl. *J. Nutr. Sci. Vitaminol.* 23: 305-318.
- [82] Miersch, J., E. M. Logvinenko, A. E. Zakalsky, G. M. Shavlovskii, and H. Reinbothe (1978) Origin of the ribityl side-chain of riboflavin from the ribose moiety of guanosine triphosphate in *Pichia guilliermondii* yeast. *Biochim. Biophys. Acta.* 543: 305-312.
- [83] Logvinenko, E. M., G. M. Shavlovskii, and A. E. Zakalsky (1979) Product of second-stage riboflavin biosynthesis in *Pichia guilliermondii*. *Mikrobiologiya* 48: 756-758.
- [84] Nielsen, P. and A. Bacher (1981) Biosynthesis of riboflavin. Characterization of the product of the deaminase. *Biochim. Biophys. Acta* 662: 312-317.
- [85] Bacher, A., U. Eggers, and F. Lingens (1973) Genetic control of riboflavin synthetase in *Bacillus subtilis*. *Arch. Mikrobiol.* 89: 73-77.
- [86] Sadique, J., R. Shanmugasundaram, and E. R. B. Shanmugasundaram (1966) Isolation of 5-amino-4-ribitylamino-uracil from a riboflavinless mutant of *Aspergillus nidulans*. *Biochem. J.* 101: 2C.
- [87] Lingens, F., O. Oltmanns, and A. Bacher (1967) Über Zwischenprodukte der Riboflavin-Biosynthese bei *Saccharomyces cerevisiae*. *Z. naturforsch.* 22B: 755-758.
- [88] Mitsuda, H. and K. Nakajima (1976) Isolation of 4-ribitylamino-5-amino-2,6-dihydroxypyrimidine from a high flavinogenic mold *Eremothecium ashbyii*. *J. Nutr. Sci. Vitaminol.* 22: 307-312.
- [89] Mitsuda, H., K. Nakajima, and Y. Yamada (1978) Isolation of 4-(1'-D-ribitylamino)-5-amino-2,6-dihydroxypyrimidine from a riboflavin-adenine-deficient mutant of *Bacillus subtilis*. *J. Biol. Chem.* 253: 2238-2243.
- [90] Bacher, A. and F. Lingens (1970) Formation of 2,5-diamino-6-hydroxy-4-(1'-D-ribitylamino) pyrimidine in a riboflavin auxotroph. *J. Biol. Chem.* 245: 4647-4652.
- [91] Hollander, I. and G. M. Brown (1979) Biosynthesis of riboflavin: reductase and deaminase of *Ashbya gossypii*. *Biochem Biophys. Res. Commun.* 89: 759-763.
- [92] Burrows, R. B. and G. M. Brown (1978) Presence in *Escherichia coli* of a deaminase and a reductase involved in biosynthesis of riboflavin. *J. Bacteriol.* 136: 657-667.
- [93] Bresler, S. E., E. A. Glazunov, D. A. Perumov, and T. P. Chernik (1977) Riboflavin biosynthesis operon of *Bacillus subtilis*. XIII. Genetic and biochemical study of mutants with regard to intermediate stages of biosynthesis. *Genetika* 13: 2006-2016.
- [94] Goodwin, T. W. and D. H. Treble (1958) The incorporation of (2- $^{14}\text{C}$ ) acetylmethylcarbinol (acetone) into ring A of riboflavin by *Eremothecium ashbyii*; a new route for the biosynthesis of an aromatic ring. *Biochem. J.* 70: 14-15.
- [95] Katagiri, H., I. Takeda, and K. Imai (1959) Synthesis of riboflavin by microorganisms. VII. The enzymic ribofla-

- vin synthesis from 4-(*N*-ribitylamino)-5-aminouracil. *J. Vitaminol.* 5: 287-297.
- [96] Goodwin T. W. and A. A. Horton (1961) Biosynthesis of riboflavin in cell-free systems. *Nature.* 191: 772-774.
- [97] Jabasani, M. T. and U. A. S. Al-Khalidi (1975) Asymmetric labeling of the xylene ring in riboflavin. *Int. J. Biochem.* 6: 735-739.
- [98] Mitsuda, H., K. Nakajima, and Y. Ikeda (1978) Effects of various metabolites (sugars, carboxylic acids and alcohols) on riboflavin formation in non-growing cells of *Ashbya gossypii*. *J. Nutr. Sci. Vitaminol.* 24: 91-103.
- [99] Schlee, K. and H. Reinbothe (1970) Purinstoffwechsel und Riboflavin Bildung in Mikroorganismen. VI. Der Einfluß von L-Valin auf die Flavinogenese von *Candida guilliermondii*. *Z. Allg. Mikrobiol.* 10: 77-80.
- [100] Plaut, G. W. E. (1971) Metabolism of water-soluble vitamins. The biosynthesis of riboflavin. pp. 11-45. In: M. Florkin, and E. H. Stotz (eds.). *Comprehensive Biochemistry*, Vol. 21.
- [101] Ali, S. N. and U. A. S. Al-Khalidi (1966) The precursors of the xylene ring in riboflavin. *Biochem. J.* 98: 182-188.
- [102] Alworth, W. C., H. N. Bacher, M. F. Winkler, A. M. Keenan, G. W. Gokel, and F. L. Wood (1970) Biosynthesis of the 4,5-dimethyl-1,2-phenylene moiety of vitamin B<sub>12</sub>. *Biochem. Biophys. Res. Commun.* 40: 1026-1031.
- [103] Keller, P. J., Q. Le Van, A. Bacher, and H. G. Floss (1983) Biosynthesis of riboflavin. <sup>13</sup>C-NMR techniques for the analysis of multiply <sup>13</sup>C-labeled riboflavins. *Tetrahedron* 39: 3471-3481.
- [104] Floss, H. G., Q. Le Van, P. J. Keller, and A. Bacher (1983) Biosynthesis of riboflavin. An unusual rearrangement in the formation of 6,7-dimethyl-8-ribityllumazine. *J. Am. Chem. Soc.* 105: 2493-2494.
- [105] Nielsen, P., G. Neuberger, H. G. Floss, and A. Bacher (1984) Biosynthesis of riboflavin. Enzymatic formation of the xylene moiety from [<sup>14</sup>C]ribulose 5-phosphate. *Biochem. Biophys. Res. Commun.* 118: 814-822.
- [106] Le Van, Q., P. J. Keller, D. H. Bown, H. G. Floss, and A. Bacher (1985) Biosynthesis of riboflavin in *Bacillus subtilis*: origin of the four-carbon moiety. *J. Bacteriol.* 162: 1280-1284.
- [107] Volk, R. and A. Bacher (1990) Studies on the 4-carbon precursor in the biosynthesis of riboflavin. *J. Biol. Chem.* 265: 19479-19485.
- [108] Katagiri, H., I. Takeda, and K. Imai (1958) Synthesis of riboflavin by microorganisms. *J. Vitaminol.* 4: 278-284.
- [109] Plaut, G. W. E. (1960) Studies on the stoichiometry of the enzymic conversion of 6,7-dimethyl-8-ribityllumazine to riboflavin. *J. Biol. Chem.* 235: PC 41.
- [110] Beach, R. L. and G. W. E. Plaut (1970) Stereospecificity of the enzymatic synthesis of the *o*-xylene ring of riboflavin. *J. Am. Chem. Soc.* 92: 2913-2916.
- [111] Bacher, A., Q. Le Van, P. J. Keller, and H. G. Floss (1983) Biosynthesis of riboflavin. Incorporation of <sup>13</sup>C-labelled precursors into the xylene ring. *J. Biol. Chem.* 258: 13431-13437.
- [112] Sedlmaier, H., F. Miller, P. J. Keller, and A. Bacher (1987) Enzymatic synthesis of riboflavin and FMN specifically labeled with <sup>13</sup>C in the xylene ring. *Z. Naturforsch.* 42c: 425-429.
- [113] Wacker, H., R. A. Harvey, C. H. Winestock, and G. W. E. Plaut (1964) 4-(1'-D-ribitylamino)-5-amino-2,6-dihydroxypyrimidine, the second product of the riboflavin synthetase reaction. *J. Biol. Chem.* 239: 3493-3497.
- [114] Mitsuda, H., T. Nadamoto, and K. Nakajima (1976) Reutilization of by-product for riboflavin formation in the riboflavin synthetase reaction. *J. Nutr. Sci. Vitaminol.* 22: 67-70.
- [115] Mironov, V. N., A. E. Kraev, B. K. Chernov, A. V. Ul'yanov, Y. B. Golva, G. E. Pozmogova, M. L. Simonova, V. K. Gordeev, A. I. Stepanov, and K. G. Skryabin (1989) Riboflavin biosynthesis gene of *Bacillus subtilis* - complete primary structure and model of organization. *Dokl. Akad. Nauk. SSSR.* 305: 482-487.
- [116] Boretskii, Iu. R., Iu. S. Skoblov, O. M. Khodova, and P. M. Rabinovich (1992) Purification and properties of GTP-cyclohydrolase from *Bacillus subtilis*. *Biokhimiya* 57: 1021-1030.
- [117] Richter, G., H. Ritz, G. Katzenmeier, R. Volk, A. Kohnle, F. Lottspeich, D. Allendorf, and A. Bacher (1993) Biosynthesis of riboflavin: cloning, sequencing, mapping and expression of the gene encoding for GTP cyclohydrolase II in *Escherichia coli*. *J. Bacteriol.* 175: 4045-4051.
- [118] Richter, G., R. Volk, C. Krieger, H. W. Lahm, U. Rothlisberger, and A. Bacher (1992) Biosynthesis of riboflavin: cloning, sequencing, and expression of the gene encoding 3,4-dihydroxy-2-butanone 4-phosphate synthase of *Escherichia coli*. *J. Bacteriol.* 174: 4050-4056.
- [119] Revuelta, J. L., M. A. Santos, and J. J. Garcia-ramirez (1990) Biosynthesis of vitamin B<sub>2</sub> in yeast, pp. 117-122. In: Committee of Biotec-90 (eds.). *From Genes to Bio-Products*. DM PPU, Murcia, Spain.
- [120] Buitrago, M. -J., G. A. Gonzalez, J. E. Saiz, and J. L. Revuelta (1993) Mapping of the *RIB1* and *RIB7* genes involved in the biosynthesis of riboflavin in *Saccharomyces cerevisiae*. *Yeast* 9: 1099-1102.
- [121] Zakal'skii, A. E., M. L. Zlochevskii, I. Stasiv, E. M. Logvinenko, M. Beburow, and G. M. Shavlovskii (1990) Cloning of the gene coding for the enzyme of the first stage of flavinogenesis in the yeast *Pichia guilliermondii*, GTP cyclohydrolase, in *Escherichia coli* cells. *Genetika* 26: 614-620.
- [122] Liauta-Teglivets, O., M. Hasslacher, Iu. R. Boretskii, S. D. Kohlwein, and G. M. Shavlovskii (1995) Molecular cloning of the GTP-cyclohydrolase structural gene *RIB1* of *Pichia guilliermondii* involved in riboflavin biosynthesis. *Yeast* 11: 945-952.
- [123] Bacher, A. and F. Lingens (1971) Formation of 6-hydroxy-2,4,5-triaminopyrimidine in *RIB7* mutants of *Saccharomyces cerevisiae*. *J. Biol. Chem.* 246: 7018-7022.
- [124] Bacher, A. (1991) Biosynthesis of flavins, pp. 215-259. In: F. Muller (eds.). *Chemistry and biochemistry of flavins*. vol.1. Chemical Rubber Co., Boca Raton, Fla, USA.
- [125] Bandrin, S. V., P. M. Rabinovich, and A. I. Stepanov (1983) Three linkage groups of genes involved in riboflavin biosynthesis in *Escherichia coli*. *Sov. Genet.* 19: 1103-1109.
- [126] Perkins, J. B. and J. G. Pero (1993) Biosynthesis of riboflavin, biotin, folic acid, and cobalamin, pp. 319-334. In:

- A. Sonenshein (eds.). *Bacillus subtilis and Other Gram-positive Bacteria: Biochemistry, Physiology, and Molecular Genetics*. American Society for Microbiology, Washington, DC, USA.
- [127] Fischer, M., S. Eberhardt, G. Richter, C. Krieger, I. Gerstenschlager, and A. Bacher (1996) Biosynthesis of riboflavin. Bifunctional pyrimidine deaminase/reductase of *Escherichia coli* and *Bacillus subtilis*. *Biochem. Soc. Trans.* 24: 35S.
- [128] Tetsuya, T., C. Ueguchi, K. Shiba, and K. Ito (1992) Insertional disruption of the *nusB* (*ssyB*) gene leads to cold-sensitive growth of *Escherichia coli* and suppression of the *secY24* mutation. *Mol. Gen. Genet.* 234: 429-432.
- [129] Garcia-Ramirez, J. J., M. A. Santos, and J. L. Revuelta (1995) The *Saccharomyces cerevisiae* *RIB4* gene codes for 6,7-dimethyl-8-ribityllumazine synthase involved in riboflavin biosynthesis. *J. Biol. Chem.* 270: 23801-23807.
- [130] Bacher, A., R. Baur, U. Eggers, H. -D. Harders, M. K. Otto, and H. Schnepfle (1980) Riboflavin synthases of *Bacillus subtilis*. Purification and properties. *J. Biol. Chem.* 255: 632-637.
- [131] Bacher, A., H. C. Ludwig, H. Schnepfle, and Y. Ben-Shaul (1986) Heavy riboflavin synthase from *Bacillus subtilis*. Quaternary structure and reaggregation. *J. Mol. Biol.* 187: 75-86.
- [132] Ladenstein, R., B. Meyer, R. Huber, H. Labischinski, K. Bartels, H. -D. Bartunik, L. Bachmann, H. C. Ludwig, and A. Bacher (1986) Heavy riboflavin synthase from *Bacillus subtilis*. Particle dimensions, crystal packing and molecular symmetry. *J. Mol. Biol.* 187: 87-100.
- [133] Mirtl, S., M. Fischer, G. Richter, J. Tack, S. Weinkauff, and A. Bacher (1996) Biosynthesis of riboflavin. Lumazine synthase of *Escherichia coli*. *J. Biol. Chem.* 271: 33201-33207.
- [134] Schott, K., J. Kellermann, F. Lottspeich, and A. Bacher (1990) Riboflavin synthases of *Bacillus subtilis*. Purification and amino acid sequence of the  $\alpha$  subunit. *J. Biol. Chem.* 265: 4204-4209.
- [135] Harvey, R. A. and G. W. E. Plaut (1966) Riboflavin synthetase from yeast. Properties of complexes of the enzyme with lumazine derivatives and riboflavin. *J. Biol. Chem.* 241: 2120-2136.
- [136] Otto, M. K. and A. Bacher (1981) Ligand-binding studies on light riboflavin synthase from *Bacillus subtilis*. *Eur. J. Biochem.* 115: 511-517.
- [137] Tibbelin, G., S. Eberhardt, G. Richter, and A. Bacher (1996) Structure investigation on recombinant riboflavin synthase from *E. coli*. *Biochem. Soc. Trans.* 24: 34S.
- [138] Santos, M. A., J. J. Garcia-Ramirez, and J. L. Revuelta (1995) Riboflavin biosynthesis in *Saccharomyces cerevisiae*. Cloning, characterization, and expression of the *RIB5* gene encoding riboflavin synthase. *J. Biol. Chem.* 270: 437-444.
- [139] Volk, R. and A. Bacher (1991) Biosynthesis of riboflavin. Studies on the mechanism of L-3,4-dihydroxy-2-butanone 4-phosphate synthase. *J. Biol. Chem.* 266: 20610-20618.
- [140] Richter, G., C. Krieger, R. Volk, K. Kis, H. Ritz, E. Gotze, and A. Bacher (1997) Biosynthesis of riboflavin: 3,4-dihydroxy-2-butanone 4-phosphate synthase. *Methods Enzymol.* 280: 374-382.
- [141] Mehta, S. U., A. K. Matoo, and V. V. Modi (1972) Ribitol and flavinogenesis in *Eremothecium ashbyii*. *Biochem. J.* 130: 159-166.
- [142] Moschau, N., K.-P. Stahmann, H. Sahm, J. B. McNeil, and A. L. Bogner (1997) Identification of *Saccharomyces cerevisiae* *GLY1* as a threonine aldolase: a key enzyme in glycine biosynthesis. *FEMS Microbiol. Lett.* 150: 55-60.
- [143] Monschau, N., H. Sahm, and K.-P. Stahmann (1998) Threonine aldolase overexpression plus threonine supplementation enhanced riboflavin production in *Ashbya gossypii*. *Appl. Environ. Microbiol.* 64: 4283-4290.
- [144] Maeting, I., G. Schmidt, H. Sahm, J. L. Revuelta, Y.-D. Stierhof, and K.-P. Stahmann (1999) Isocitrate lyase of *Ashbya gossypii* – transcriptional regulation and peroxisomal location. *FEBS Lett.* 444: 15-21.
- [145] Taylor, K. M., C. P. Kaplan, X. Gao, and A. Baker (1996) Localization and targeting of isocitrate lyases in *Saccharomyces cerevisiae*. *Biochem. J.* 319: 255-262.
- [146] Chaves, R. S., P. Herrero, I. Ordiz, M. A. del Brio, and F. Moreno (1997) Isocitrate lyase localisation in *Saccharomyces cerevisiae* cells. *Gene* 198: 165-169.

[Received February 28, 2001; accepted April 24, 2001]