

b

Haemophilus influenzae
(Buccal Cellulitis) 1

. . *

= Abstract =

A Case of Buccal Cellulitis Caused by *Haemophilus influenzae*
Type b in an Immunocompetent Child

Jin A Lee, M.D., Dong Ho Kim, M.D.
Ja Wook Koo, M.D.* and Hoan Jong Lee, M.D.

Department of Pediatrics, College of Medicine, Seoul National University,
Department of Pediatrics, College of Medicine, Inje University, Seoul, Korea*

Buccal cellulitis which presents with high fever and a swelling of the cheek with violaceous hue in young children is most often caused by *H. influenzae*. Bacteremia is common in buccal cellulitis caused by *H. influenzae*, and a culture of cerebrospinal fluid should be obtained because meningitis may be present despite the lack of meningeal irritation signs. Although buccal cellulitis is considered to be one of the important manifestations of *H. influenzae* infection, only two cases have been reported in Korea yet. We experienced a case of buccal cellulitis with *H. influenzae* bacteremia in an immunocompetent girl of 18-month-old. She was presented with high fever followed by rapidly progressive swelling and tenderness of both cheeks with violaceous hue in four hours. The blood culture revealed *H. influenzae* type b. There was no concurrent otitis media, sinusitis, or meningitis and no portal of entry was identified. Fever subsided two days after starting intravenous cefotaxime. Intravenous cefotaxime was subsequently changed to oral cefixime, and antibiotics were administered for a total of two weeks. We report this case with a review of related literature.

Key Words : *Haemophilus influenzae*, Buccal cellulitis

(dermis)

, A

Tel : 02)760-3633 Fax : 02)745-4703
E-mail : hoanlee@plaza.snu.ac.kr

(Hib)

, b *Haemophilus influenzae*

Hib가 4 가 가

Hib 6 가 가

5 Hib 1

5 Hib 39 40

Hib (flucloxacillin + gentamicin)

1986 1996 *H. influenzae*

³⁾ 2 가 가

⁴⁾ Hib 1999 4 3

105/60 mmHg,

32 , 105 ,

37.0 . 11 kg(25 50) .

Hib 8 × 10

cm 가

(Fig. 1). 가

: , , 18

:

: 가

3.2 kg . BCG 1 , B 가

3 , DTaP/TOPV 4 , MMR 1

, Hib

가 : 1 1

4 ,

:

3

, 2

38.3



Fig. 1. Swelling of both buccal tissue and neck especially right side with violaceous hue in an 18-month-old girl.

: 1
 20,100/
 μL(16%, 65%, 14%) ,
 11.1 g/dL,
 32.3%, 19,700/ μL(16%,
 44%, 15%), 328,000/ μL
 . 15 mm/hour , C-
 19.1 mg/dL .
 6.2 g/dL, 3.4 g/dL, 5 mg/
 dL, 0.4 mg/dL, alanine aminotransferase
 18 U/L, aspartate aminotransferase 11 U/L, amylase
 33 U/L, creatine kinase 22 U/L, lactate dehydrogen-
 ase 209 U/L . 1 1
*H. influenzae*가
 , 2
 .
 1/ μL, 1/ μL(0/ μL, 1/

μL), 15 mg/dL, 56 mg/dL . , *H.*
influenzae

,
 . X- ,
 ,
 (spinal accessory lymph node)
 , . *H. in-*
fluenzae a f 6가
 (Denka Seiken, Tokyo, Japan)

b
 : nafcillin(100 mg/kg/day)
 cefotaxime(100 mg/kg/day)
 . 2 , 6
*H. influenzae*가
 nafcillin
 . 12
 cefixime(8 mg/kg/day) 3
 , 6

, Hib (PRP-T)
 .
 .
H. influenzae ,
 , , , ,
 , ,
 . a f
 6가 (nonencapsulated)
 , b
 ,
 .
 Hib 5 ,
 3 3 . Hib
 가 ,
 , , , ,
 .
 5
 Hib

, 50 65% , ⁸⁾ .
 10 30%, 5 9%, 5%, *H. influenzae*
 2 15% ^{1, 2)} Hib ,
 가 . 1986
 가 . 1996 11 *H.*
 5 100,000 16 69 *influenzae*
 Hib , 6 11 , 38
 275/100,000 Hib 25 가 , 8 가 , 3 가 , 1
⁵⁾ 5 200 1 가 , 1 가 ,
 , , Hib 1 가 , 9 가 , 2 가
 , 350 1 ⁶⁾ , ³⁾ .
 5 10% Hib
 가 Hib 가 5 .
 20 30/100,000 Hib 가 ,
 . Hib ,
 , Alaskan Eskimos, Canadian
 Inuts, Navajos Apaches , , Hib
 Apache Indian
 Hib 5 100,000 ⁷⁾ , Hib . ,
 600 , 150 가
 1 가 가
 가 . Hib ,
 , , 12 Hib
 가 5 100,000 10
 2 4 , Finland Hib 가 .
 , Hib 가 가 ⁴⁾ .
H. influenzae biotype II IV , 9
 2 ,
 ,
 Hib 5 28
 10 1.1 10 (orbital cellulitis) , 2
^{8 10)} Lau
 5 2.67/
 100,000 , 57 .
 , *H. influenzae* 2

가⁴⁾ (violaceous hue) Hib
^{11 13)} 1953 Alexander ,
¹⁴⁾ Hib Hib 가 . 2/3
가 , Ginsburg¹⁵⁾ ,
1970 1980 72 Hib . 18
. 72%가 1 Hib
. , .
13.8
. 68% 가
23 22 Hib가 15,000/ μL 가
. 86% , 7.6%¹⁹⁾ .
. Carter¹⁶⁾ 60 .
(portal of entry)가 가 가
21 39 ,
, *Staphylococcus au-* 0.1 mL 가
reus ,
26 19 *H.* . 55 86%
influenzae . Fleisher¹⁷⁾ 4.2 8.8%
78 ,
26 . 26^{15, 18, 19)} .
Hib , 19 Hib가
7 , 7 1 .
. Baker¹⁸⁾ 80 .
, 34²⁰⁾ , 72
, 46 (preseptal
cellulitis) . 67 Hib, 12
, 1 *Streptococcus pyogenes*가^{12, 13)} . Baker¹⁸⁾
. 7
6 Hib 1 , 5
(irritability) ,
. 2
, 1
, 1 . , 가
(pleocytosis), ,
. Hib 6 3 . Hib가
. 가
가 . 7 10

5 , 2 15%
가
Hib
18
cefotaxime(200 mg/kg/day) ceftriaxone(100 mg/kg/day)
Ampicillin chloramphenicol
Hib , 2 3
cephalosporin 1
가 methicillin(200 mg/kg/day)
ampicillin, chloramphenicol, cefotaxime
amoxicillin ampicillin
amoxicillin-clavulanate(40 mg/kg/day) 3
cephalosporin cefixime cefpodoxime
H. influenzae beta-lactamase ampicillin²¹⁾
가 50%
Hib 가
가 Hib
1989 5
가 100,000
41 1993 2²²⁾, 1997 0.4
Hib b
가
H. influenzae
가
Hib

- 1) Broadhurst LE, Erickson RL, Kelley PW. Decreases in invasive *Haemophilus influenzae* diseases in US Army children, 1984 through 1991. JAMA 1993;269:227-31.
- 2) Takala AK, Eskola J, Peltola H, et al. Epidemiology of invasive *Haemophilus influenzae* type b disease among children in Finland before vaccination with *Haemophilus influenzae* type b conjugate vaccine. Pediatr Infect Dis J 1989;8:297-302.
- 3) Lee HJ. Epidemiology of systemic *Haemophilus influenzae* disease in Korean children. Pediatr Infect Dis J 1998;17(Suppl):S185-9.
- 4) , , . *Haemophilus influenzae* Biotype II IV
2 . 1993;13: 631-6.
- 5) Wilfert CM. Epidemiology of *Haemophilus influenzae* type b infections. Pediatrics 1990;85: S631-5.
- 6) Cochi SL, Broome CV, Hightower AW. Immunization of U.S. children with *Haemophilus influenzae* type b polysaccharide vaccine : a cost effectiveness model of strategy assessment. JAMA 1985;54:521-9.
- 7) Clements DA. *Haemophilus influenzae* type b. In : Klugman S, Katz SL, Gershon AA, Wilfert CM, editors. Infectious diseases of children. 9th ed. St. Louis : Mosby, 1992:127-42.
- 8) Lau YL, Low LC, Yung R, et al. Invasive *Haemophilus influenzae* type b infections in

- children hospitalized in Hong Kong, 1986-1990. *Acta Paediatr* 1995;84:173-6.
- 9) Yang Y, Shen X, Jiang Z, et al. Study on *Haemophilus influenzae* type b diseases in China: the past, present and future. *Pediatr Infect Dis J* 1998;17(Suppl):S159-65.
- 10) Ishikawa T, Asano Y, Morishima T, et al. Epidemiology of bacterial meningitis in children: Aichi Prefecture, Japan, 1984-1993. *Pediatr Neurol* 1996;14:244-50.
- 11) Walker JS, Corcoran KJ. Buccal cellulitis. *Am J Emerg Med* 1990;8:542-5.
- 12) Rapkin RH, Bautista G. *Haemophilus influenzae* cellulitis. *J Dis Child* 1972;124:540-2.
- 13) Charnock DR, White T. Bruised cheek syndrome: *Haemophilus influenzae*, type b, cellulitis. *Otolaryngol Head Neck Surg* 1990;103:829-30.
- 14) Alexander HE. Influenza bacillus infections. In: Holt LE, McIntosh R, editors. *Pediatrics*. New York: Appleton-Century-Crofts, 1953:1200.
- 15) Ginsburg CM. *Haemophilus influenzae* type b buccal cellulitis. *J Am Acad Dermatol* 1981;4:661-4.
- 16) Carter S, Feldman WE. Etiology and treatment of facial cellulitis in pediatric patients. *Pediatr Infect Dis* 1983;2:222-4.
- 17) Fleisher G, Heeger P, Topf P. *Haemophilus influenzae* cellulitis. *Am J Emerg Med* 1983;1:274-7.
- 18) Baker RC, Bausher JC. Meningitis complicating acute bacteremic facial cellulitis. *Pediatr Infect Dis* 1986;5:421-3.
- 19) Chartrand SA, Harrison CJ. Buccal cellulitis reevaluated. *Am J Dis Child* 1986;140:891-3.
- 20) Nelson JD, Ginsburg CM. An hypothesis on the pathogenesis of *Haemophilus influenzae* buccal cellulitis. *J Pediatr* 1976;88:709-10.
- 21) Kwak YH, Jung HS, Park SE, et al. Serotypes and antimicrobial susceptibility in clinical isolates of *Haemophilus influenzae* from Korean children in prevaccination era. *J Korean Med Sci* 2000;15:616-22.
- 22) From the Centers for Disease Control and Prevention. Progress toward elimination of *Haemophilus influenzae* type b disease among infants and children - United States, 1987-1993. *MMWR Morb Mortal Wkly Rep* 1994;43:144-8.