

Abstract

Effects of Treadmill Training on Hyperextension of the Knee and Cadence in Patients With Hemiplegia

Park Chul-hong, B.H.Sc., P.T.

Dept. of Physical Therapy, Seoul Su Medical Center

Chung Bo-in, Ph.D.

Dept. of Rehabilitation Therapy, Collage of Health Science, Yonsei University

Institute of Health Science, Yonsei University

This study addresses the effects of treadmill training on hyperextended knee and cadence in patients with hemiplegia. A single subject research design with multiple baselines across individuals was used for the study. Two patients with hemiplegia participated in the experiment. The experiment consisted of interventions where the patients were asked to ambulate for 15 minutes at a comfortable walking speed on the treadmill with 11% slope grade and were allowed to rest for 10 minutes. Patients, then, were asked to ambulated 20 meters at walkway. The number of occurrences of knee hyperextension and the total number of steps were recorded. The results showed that the occurrence of knee hyperextension decreased by approximately 30% after the first session of the treadmill training and continued to gradually decrease during the following sets of treadmill training. Meanwhile, there was a slight increase in the cadence to a negligible extent. These results suggest that the gait training on the sloped treadmill may be helpful for correcting the knee hyperextension in patients with hemiplegia.

Key Words: Treadmill training; Hyperextended knee; Cadence.

(muscle weakness), (abnormal muscle tone) (abnormal movement pattern) (Sharp Brouwer, 1997). (Brandstater, 1983) (gait velocity), (stride length), (stride width) (Brandstater, 1983) (cadence), (step length), (Burridge, 1997). (disability)가 (Duncan, 1994; Langton-Hower, 1990). (stance phase) (Davis, 1985). , 0 15° (Hassid, 1997; Hesse, 1999; Visintin, 1998). Waafjord (1990) (toe-off)가 (Hesse, 1999) (Sawner LaVigne, 1991).

가 . 가 ,

1. 가 , 1 46

Scale G1 Modified Ashworth 2

45 8

가 2 G1 (1).

2. (single subject design) (multiple baseline across individuals design)

2000 11 20 12 16 2 3

가 . 가

가 6 가

가 20 m 가

500 m


Modified Ashworth 가 20 m

Scale G2


1.

	()	()	
1	46	9	G1
2	45	8	G1

15 가
 100%
 (%)
 3.
 (goniometer)
 가 180°
 micro-
 switch가 20 m
 1 m 가
 1
 m
 가



2. (A: , B:)



1. 2
 (A: , B:)

(sine) 100

11%

1

5

6

4.

Startrac
 series. USA, 1997)

(ETL Co. Tr 4000

8
 가

30

가 가

15
 가

가 가

10

100 m

, 10%

5.

가

100 m

10 m

1

(visual analysis)

가

. Hesse (1999)

1 2

,

가

99.0%,

1

67.8%

2

99.3%,

. Winter(1989)

76.4%

2 2

,

(stepping),

1

60.0

steps/min,

63.4 steps

/min

2

87.2 steps/min,

87.3

steps/min

,

가

가

(Hassid

Knuttson Richards(1979)

, 1997).

가

(reciprocal innervation)

가

(Davies, 1990).

Sherrington(1913)

. Cipriani (1995)

10%

가

(initial contact)

가

가 가

Sherrington(1913)

가
 . 2
 1
 60.0 steps/min , 가
 63.4 steps
 /min 가 가
 , 가
 가 가
 가

. 1999:44-46.

Brandstater ME, Bruin HD, Gowland C, et al. Hemiplegic gait: Analysis of temporal variables. Arch Phys Med Rehabil. 1983;64:583-587.

(sensitivity)

Burrige JH, Taylor PN, Hagan SA, et al. The effects of common peroneal stimulation on the effort and speed of walking: A randomized controlled trial with chronic hemiplegic patients. Clin Rehabil. 1997;11:201-210.

가

Cipriani DJ, Armstrong CW, Gaul S. Backward walking at three levels of treadmill inclination: An electromyographic and kinematic analysis. J Orthop Sports Phys Ther. 1995;3: 95-102.

가

Davies PM. Selective Trunk Activity in the Treatment of Adult Hemiplegia: Right in the middle. Potland, Book News Inc., 1990:53-57.

Davies PM. Steps to Follow. Berlin Heidelberg, Springer-Verlag, 1985.

Duncan PW. Stroke disability. Phys Ther.

- 1994;74:399-407.
- Hassid E, Guttry M, Dobkin BH, et al. Improved gait symmetry in hemiparetic stroke patients induced during body weight-supported treadmill stepping. *J Neuro Rehabil.* 1997;11:21-26.
- Hesse S, Konrad M, Uhlenbrock D. Treadmill walking with partial body weight support versus floor walking in hemiparetic subjects. *Arch Phys Med Rehabil.* 1999;80:421-427.
- Knuttson E, Richards C. Different types of disturbed motor control in gait of hemiplegic patients. *Brain.* 1979;102:405-430.
- Langton-Hower R. Rehabilitation after stroke. *Q J Med New Series.* 1990;76:659-674.
- Sawner KA, LaVigne JM. Brunnstrom's Movement Therapy in Hemiplegia. 2nd eds. 1991:131-144.
- Sharp SA, Brouwer BJ. Isokinetic strength training of the hemiparetic knee: Effects on function and spasticity. *Arch Phys Med Rehabil.* 1997;78:1231-1236.
- Sherrington CS. Reflex inhibition as a factor in the coordination of movements and postures. *Quart J Exp Physiol.* 1913;6:251.
- Visintin M, Barbeau H, Mayo NE, et al. A new approach to retrain gait in stroke patients through body weight support and treadmill stimulation. *Stroke.* 1998;29:1122-1128.
- Waagfjord J, Levangle PK, Certo CM. Effects of treadmill training on gait in a hemiparetic patient. *Phys Ther.* 1990;70:549-560.
- Winter DA. Biomechanics of normal and pathological gait: Implications for understanding human locomotor control. *J Motor Behavior.* 1989;21:337-356.