

A Study on Storage of a Major Herbal Medicine Material, Peucedani Radix

Seongkyu Choi*, Kyeong Won Yun and Byung-Sun Kwon

Department of Oriental Medicine Resources, Suncheon National University

*College of Agriculture and Life science, Suncheon National University, Suncheon 540-742, Korea.

ABSTRACT

To develop optimal storage method of root of Peucedani Radix, which has been grown as major cultural herbal medicine material, root of Peucedani Radix was stored for 10 months with different packing materials and sealing methods. The loss in dry weight as influenced by packing materials and sealing methods was the lowest at vacuum packing and followed by complete sealing methods with transparent polyethylene film. The ratio of root rot during the storage period was not significantly different between packing materials but was significantly different between sealing conditions. Conclusively vacuum packing and complete sealing with polyethylene film appears to be optimal for storage of Peucedani Radix.

Key Words : Peucedani Radix, storage, herbal medicine materials, polyethylene film

INTRODUCTION

Peucedamum japonicum Thunberg, a perennial herbal medicinal plant, has been grown through all the Korean Peninsula. The plant started to sprout early in May and bloom on July. *Peucedamum japonicum* is being cultivated at the middle southern area of Korea, and it needs to have long growing period(Choi, 1996; Kim et al, 1997).

Peucedamum japonicum harvested on the middle of November. After harvest, tertiary roots and shoot parts of the plant are removed by cutting and washed with fresh tap water. The root parts were dried by natural sunshine, which usually used in farmer household. In case of drying by sunshine, even though there is a little

difference according to weather, it takes 20 to 30 days to dry. After drying the materials were wrapped with polyethylene films, stored at well-ventilated place and finally shipped as major cultural herbal medicine materials.

Herbal medicine materials are generally stored until they were sold or used for medicine. During storage, temperature, humidity, and ventilation of room were very important factors to keep good quality of plants even they well dried and cleaned. The factors can influence damage in quality as well as aberration in the efficacy of chemicals(Choi, 1994; Choi et al, 1997). Several points must be considered during storage. It should be ventilated to lessen humidity, stored at cooling room, prevented from insects, and stored at airtight container. Choi (1994) reported in a previous

Table 1. Comparison of loss in weight as influenced by packing materials and sealing methods in root storage of *Peucedamum japonicum* Radix

Packing materials	Sealing degree	Investigating time					
		1999			1998		
		Dec. 15	Feb. 15	Apr. 15	Jun. 15	Aug. 15	Oct. 15
T. P. E ¹⁾	C.sealing ²⁾	1,000(g)	879(g)b ³⁾	869(g)b	855(g)b	854(g)b	849(g)b
	Punch ³⁾	1,000	865b	857b	838b	803c	785c
B. P. E	C.sealing	1,000	877b	886b	859b	855b	848b
	Punch	1,000	866b	858b	839b	801c	787c
G. P. E	C.sealing	1,000	875b	867b	857b	855b	849b
	Punch	1,000	866b	858b	837b	803c	785c
T. P. E	V. P. ⁴⁾	1,000	962a	949a	940a	932a	916a

³⁾ Mean separation within column by Duncan' s multiple range test, 5% level of significance

¹⁾ T. P. E : Transparent polyethylene film

B. P. E : Black color polyethylene film

G. P. E : Green color polyethylene film

²⁾ C. sealing : Complete sealing

³⁾ Punch : Punch(\varnothing 5mm) on polyethylene film

⁴⁾ V. P : Vacuum packing

study that herbal medicinal materials should be stored at the place where are well ventilated with low temperature, dry and prevented from insects.

Roots of *Peucedamum japonicum* have to be completely dried to prevent from decaying. Especially quality of goods might be lowered by fungus. This study was conducted to determine causes and their solution of loss in quantity as well as in quality during storage of *Peucedani Radix*. Loss in amount and rotting rate at different harvest time were examined according to storage of different packing materials and sealing conditions.

MATERIALS AND METHODS

Peucedani Radix was planted at herb garden of Department of Oriental Medicine Resources, Suncheon National University in November, 2000. Roots of one hundred plants were carefully sampled without damage by using agricultural spades at 60cm high from soil surface in November, 2001


The roots were washed with tap water, dried under natural sunshine for 24 hours, and dried again with cold

wind at 40°C for 7 days. After drying the roots were chopped into 10 cm in length of main root. Mean root weight a plant was about 15 grams.

The collected roots of *Peucedamum japonicum* were separately wrapped in different polyethylene film envelopes(15cm \times 30cm \times 0.05mm) with transparent, black, and green color. The envelopes containing samples were treated with vacuum packing, complete sealing, and punching method. Vacuum packing of transparent polyethylene film was using Vacuum Packer (JW-500XL). Punching on polyethylene film was made into 5mm in diameter and 10 punches an envelop were evenly given.

After packing the envelopes were dried and stored at cooling room until investigated. Loss in weight and rotten rate after storage were measured with 6 times at 2 month-intervals from December 15, 2000 to October 15, 2001. All assessment was made according to RDA Standard Methods for Field Experiment(Rural Development Administration, 1989).

RESULTS AND DISCUSSION



T.P.C(Transparent Polyethylene film Complete sealing)
 B.P.C(Black color Polyethylene film Complete sealing)
 G.P.C(Green color Polyethylene film Complete sealing)
 T.P.V(Transparent Polyethylene film Vacuum packing)

T.P.P(Transparent Polyethylene film Punch)
 B.P.P(Black color Polyethylene film Punch)
 G.P.P(Green color Polyethylene film Punch)

Fig. 1. The change of rot ratio as influenced by packing materials and sealing methods in root storage of *Peucedanum japonicum*

1. Dry Weight of *Peucedanum japonicum* Root during Storage

Dry weight of *Peucedanum japonicum* root during storage as affected by different packing materials and sealing methods was determined 6 times at 2 month-intervals in 2000 and 2001(Table 1).

No difference in dry weight among storage durations was observed, but complete sealing treatment affected dry weight of plant root. Dry weight of *Peucedanum japonicum* root was gradually reduced with time regardless of packing materials and sealing methods. In sealing treatment, dry weight of root stored with complete sealing with punch were more reduced than with vacuum packing.

It was thought that water from the dried roots could be naturally evaporated through complete sealing or punch with time during storage. Root dry weight of *Peucedanum japonicum* sealed with punch was more reduced than complete sealing without punch. In packing materials, the colors of polyethylene film did not affect root dry weight. This result showed that storage with complete sealing without punch could be

better than storage by sealing with punch for long term storage. Storage with vacuum packing did not affect root dry weight and was best among treatments. Dry weight of *Peucedanum japonicum* root was little bit changed by vacuum packing methods during storage. These results showed that storage methods with complete sealing or with vacuum did not affect dry weight of root because of restriction of air exchange.

Chung et al (1991) reported that complete sealing storage with polyethylene film sustained quality of kiwi-fruits for long term. Therefore, storage with complete sealing or with vacuum packing could be suitable for *Peucedanum japonicum* root, indicating no change in root weight, no humidity, and low infestation of insects and disease. However, more detail researches were needed for storage of other medicinal plant species.

2. Rotten Rate of *Peucedanum japonicum* Root during Storage

Rotten rate of *Peucedanum japonicum* root affected by different packing materials and sealing methods was

examined (Fig. 1).

There was no significance in rotten rates among colors of packing materials, transparent, black, and green polyethylene film. Rotten rate of root packed with transparent P.E. film was slightly lower than with black or green P.E. film. Rotten rate of root was about 12% at 7 months after storage but was 59% at 11 months after storage, increasing with time. At 11 months after storage, rotting rate of root with vacuum packing was the least(12%) and the highest with punch storage(50-60%). The results indicates that storage by vacuum packing or by complete sealing appeared low rotting rate of *Peucedamum japonicum* root. However, storage with punched polyethylene film allowed diseases and insects to enter into packing or sealing materials through air punches, and so increased rotten rates. More researches on kinds of pests and infestation rate were required to better understand rotting mechanism.

Seong et al (1996) suggested that dry roots must be fumigated with chloroform and methyl bromide when agricultural products were manufactured and stored to prevent from damaging of insects and weevil. The results of present study indicate that fumigates must be applied to control pests before storage with punched P.E. film for long term storage.

LITERATURE CITED

- Choi, K.J. 1994. A basic study on the storage of major Chinese Herbal *Medicine Materials*, *Moutan Radicis Cortex*. J. Oriental Bot. Res. 7:183~186.
- Choi, S .K. 1998. A basic study on the storage of major Cultural Herbal Medicine Materials, *Scutellarise Radix*. Journal of Basic Science. 9:101~106.
- Chung, D.S., Son, Y.K., Cho, K.D., Yoon, I.W. and Han, P.J. 1991. Studies on Safe Storage Methods of Kiwifruits Res. Rept. RDA(S., F.M., & F.P.U.). 33:39~45.
- Kim, J.C., Kim, J.H., Joung, K.R., Kim, K.J. and Park, H.P. 1997. Effects of Irrigation and Sowing Time on Growth and Yield of *Bupleurum falcatum* L. Korean J. Medicinal Crop Sci. 5::318~324.
- R. D. A.(Rual Development Administration). 1989. Research investigation standard of agriculture (medicinal crop)
- Seong, J.D., Park, Y.J., Kim, G.S., Kim, H.T., Suh, H.S. and Kim S.M. 1996. Effects of Topping on Growth and Root Yield in *Bupleurum falcatum* L. Korean J. Medicinal Crop Sci. 4:153~156.

(Received May 10, 2002)

(Accepted May 31, 2002)