

Synthesis and Anion Binding Properties of Bifunctional Urea Derivative of Calix[4]diquinone

Jung Min Oh, Eun Jin Cho, Byung Ju Ryu, Young Ju Lee, and Kye Chun Nam^{*}

Department of Chemistry and Institute of Basic Science, Chonnam National University, Gwangju 500-757, Korea

Received June 4, 2003

Key Words : Bifunctional receptor, Calix[4]diquinone, Urea derivative, Anion binding

The simultaneous binding of cationic and anionic guest species by ditopic receptors is a rapidly developing new field for ion pair recognition of environment and biological importance. By taking advantage of calix[4]arene framework, a few neutral bifunctional receptors¹⁻¹¹ were developed and showed the enhanced binding constants of hydrophilic anions in the presence of cations in organic media. As a consequence of their upper and lower rim of topologies the calixarenes are attractive host molecules to modify and so create unique geometries for the multifunctional receptors. The combination of two binding sites, urea for anions and two podand groups for cations, yields receptor that is capable of binding anions and alkali metal cations simultaneously. In addition to ion binding ability, two quinone moieties are introduced for the redox switchable receptors.⁵⁻⁷ We report here the synthesis of a new bifunctional receptor calix[4]-diquinone **4** containing two urea moieties and two podand groups and the simultaneous binding properties of cation and anion guests were studied.


Results and Discussion

By taking advantage of a selective 1,3-alkylation, dialkylated urea derivative **3** was prepared by the reaction of *p-t*-butylcalix[4]arene **1** and 2-[(2-(*N*'-phenylureido)ethoxy)ethanol *p*-toluenesulfonate **2** in the presence of K₂CO₃.⁸ In order to introduce the long urea podand chain into calixarene

effectively, **2** was prepared separately from the tosylation of 2-[(2-(*N*'-phenylureido)ethoxy)ethanol which was then synthesized by the reaction of 2-(2-aminoethoxy)ethanol with phenylisocyanate (Scheme 1). In order to increase the cooperative binding properties, cone conformers of receptor **4** are desired and under the condition applied the high yield cone products were obtained. Bifunctional urea derivative calix[4]diquinone **4** was prepared from the oxidation of **3** with TFA (thallium trifluoroacetate) in TFA (trifluoroacetic acid).

The ¹H NMR spectrum of **3** shows a pair of doublets at δ 4.39 and 3.39 for the bridged methylene protons, two singlets at δ 0.98 and 1.31 for *t*-butyl protons, a singlet at δ 8.01 for OH protons and a singlet and a triplet at δ 7.10 and 6.52 for N-H protons, indicating that **3** exists as a cone conformation. The cone conformation was confirmed by the ¹³C NMR spectrum.⁹ The ¹H NMR spectrum of bifunctional calix[4]diquinone urea derivative **4** shows a pair of doublets at δ 3.21 and 4.01 for the bridged methylene protons, a singlet at δ 1.08 for *t*-butyl protons, a broad singlet and a triplet at δ 6.38 and 7.88 for N-H protons.

The anion binding properties were examined by ¹H NMR titration experiments in CDCl₃. Substantial down field shift of urea NH signal was observed when tetrabutylammonium acetate, dihydrogen phosphate, hydrogen sulfate, chloride and bromide salts were added, indicating that anion binding is taking place at the urea vicinity. Figure 1 showed the ¹H



Scheme 1. Synthesis of bifunctional receptors.




Figure 1. ^1H NMR spectra of **4** (a) without cation and anion. (b) with 1 equivalent of NaClO_4 . (c) with 5 equivalents of TBACl. (d) with 1 equivalent of NaClO_4 and 5 equivalents TBACl in CDCl_3 .

NMR spectra of **4** in the presence sodium and chloride ions. When sodium ions were added, two N-H protons signals at δ 6.38 and 7.88 was not changed much, but a slight upfield shift of two aromatic singlets at δ 6.80 and 6.71 and a widening move of a pair of doublets at δ 3.21 and 4.01 for the bridge methylene protons were observed, indicating that sodium ions could be bound at the podand oxygen ether sites³ and fixed the conformation of calixarene. Further addition of sodium ions did not change the spectrum, indicating that an 1:1 complex with sodium and **4** could be formed. On the other hand, when chloride ions were added, a large downfield shift of N-H protons were observed without changing aromatic signals as shown in spectrum (c) in Figure 1. In order to investigate the binding enhancement of anions in the presence of alkali metal cations, anion titrations were conducted in the presence of sodium ion. Obviously a further downfield shift of N-H proton signals were observed when sodium ions were bound as shown in spectrum (d) in Figure 1. Stability constants (Table 1) were calculated from the titration results using EQNMR¹⁰ for complexation with anions. A significant increase in the strength of anion binding is observed when sodium metal ions are bound simultaneously. Chloride binding strength increase more than 20 fold and bromide binding seven fold in the presence of sodium ion. This positive cooperative binding of the chloride and bromide in the presence of sodium could be attributed through electrostatic effects of the complexed sodium ions. The electrochemical property of **4** was investigated using cyclic voltammetry. The most significant result was observed with H_2PO_4^- anion. The addition of dihydrogen phosphate anion causes a 27 mV cathodic shift.

In conclusion, bifunctional receptor **4** can bind alkali metal cations and halide anions simultaneously with positive cooperativity. This compound has a potential for practical application, and is currently being investigated.

Table 1. Stability constants (K_a) of **4** in CDCl_3 and cathodic shifts in the presence of anions in DMF

Metal	Anion	$K/\text{dm}^3 \text{mol}^{-1}$	ΔE (mV)
	H_2PO_4^- ^a	2340	27
	HSO_4^-	1100	14
	CH_3CO_2^-	890	0
	Cl^-	110	0
	Br^-	308	0
Na^+	H_2PO_4^-	— ^b	
	HSO_4^-	— ^b	
	CH_3CO_2^-	— ^b	
	Cl^-	2200	
	Br^-	2200	

^aTetrabutylammonium salts. Errors estimated to be <10%. ^bDue to the precipitate, stability constants can not be obtained.

Experimental Section

5,11,17,23-Tetra-tert-butyl-25,27-bis(cyanopropoxy)-26,28-dihydroxycalix[4]arene (1). was prepared by the known procedure.¹¹ mp ≥ 300 °C.

2-[2-(N'-Phenylureido)ethoxy]ethanol p-toluenesulfonate (2). To a solution of 2.0 g (8.9 mmol) of 2-[2-(N'-phenylureido)ethoxy]ethanol in 100 mL CH_2Cl_2 , 5.1 g (26.7 mmol) of *p*-toluenesulfonyl chloride was added. After the reaction flask was immersed into an ice-water bath, 3.7 mL of triethylamine was added slowly and stirred for 12 h at room temperature. Evaporation and extraction with ether provided crude products which was purified by column chromatography (eluent, CHCl_3 : acetone = 10 : 1) to yield 1.85 g (55%) of **2**. ^1H NMR (CDCl_3) δ 7.80 (d, 2H, $J = 6.5$ Hz, ArH), 7.38-7.25 (m, 7H, ArH and NH), 7.05 (t, 1H, $J = 7.2$ Hz, ArH), 6.90 (br t, 1H, NH), 4.24-3.43 (m, 8H, $-\text{CH}_2-$), 2.45 (s, 3H, $-\text{CH}_3$).

5,11,17,23-Tetra-tert-butyl-25,27-bis[2-{2-(N'-phenylureido)ethoxy}ethyl]oxy-26,28-dihydroxycalix[4]arene (3). To a solution of 0.50 g (0.77 mmol) of **1** and 1.1 g K_2CO_3 in 100 mL acetonitrile, 0.64 g (1.7 mmol) of **2** was added. The reaction mixture was refluxed for two days under nitrogen condition. After removing the solvents, the crude products were purified by column chromatography (eluent, CHCl_3 : *n*-hexane : ethyl acetate = 2 : 1 : 1) to yield 0.8 g (90%) of **3**. ^1H NMR (CDCl_3) δ 8.01 (s, 2H, OH), 7.22-7.18 and 6.96 (m, 10H, ArH), 7.14 and 6.83 (two s, 4H, ArH), 7.10 (s, 2H, NH), 6.52 (br t, 2H, NH), 4.39 and 3.39 (a pair of d, 8H, $J = 13.2$ Hz, ArCH_2Ar), 4.16-3.59 (m, 16H, $-\text{CH}_2-$), 1.31 and 0.98 (two s, 36H, $\text{C}(\text{CH}_3)_3$); ^{13}C NMR (CDCl_3) δ 156.1 ($-\text{NHCONH}-$), 149.7, 149.14, 147.5, 143.1, 139.6, 132.5, 128.8, 128.4, 125.84, 125.79, 122.0 and 118.8 (Ar), 75.8 ($-\text{OCH}_2-$), 70.1 and 70.0 ($-\text{CH}_2\text{OCH}_2-$), 40.5 ($-\text{CH}_2\text{N}-$), 34.0, 31.64, 31.59, 31.0 and 29.3 ($-\text{C}(\text{CH}_3)_3$ and ArCH_2Ar). MS (MALDI TOF) m/z 1084.5 (M-Na).

5,17-Di-tert-butyl-26,28-bis[2-{2-(N'-phenylureido)ethoxy}ethyl]oxy-26,28[4-25,27-diquinone (4). To a solution of 0.5 g (0.46 mmol) of **3** in 10 mL trifluoroacetic acid, 0.75 g (1.37 mmol) of $\text{Ti}(\text{CF}_3\text{CO}_2)_3$ was added under

nitrogen condition. After 24 h stirring the solvents were removed. The crude products were purified by column chromatography (eluent. CHCl₃ : acetone = 1 : 1) to yield 0.16 g (35%) of **4**. ¹H NMR (CDCl₃) δ 7.89 (s, 2H, NH), 7.44 (d, 4H, *J* = 7.7 Hz, ArH), 7.27 (t, 4H, *J* = 6.9 Hz, ArH), 6.99 (t, 2H, *J* = 7.3 Hz, ArH), 6.80 and 6.71 (two s, 8H, ArH), 6.38 (br t, 2H, NH), 4.01 and 3.21 (a pair of d, 8H, *J* = 13.0 Hz, ArCH₂Ar), 3.81-3.48 (m, 16H, -CH₂-), 1.08 (s, 9H, C(CH₃)₃); ¹³C NMR (CDCl₃) δ 189.2 and 185.7 (-CO), 156.1 (-NHCONH-), 153.2, 148.0, 146.7, 139.9, 132.8, 129.8, 129.3, 128.9, 126.5, 125.5, 122.2, 118.9 and 118.1 (Ar) 73.7 (-OCH₂-), 71.0 and 70.6 (-CH₂OCH₂-), 40.5 (-CH₂N-), 34.1, 31.3 and 29.3 (-C(CH₃)₃ and (ArCH₂Ar). MS (MALDI TOF) *m/z* 999.5 (M + Na).

Acknowledgment. This work was supported by Grant No. R01-2000-00047-0 from the Basic Research Program of the Korea Science & Engineering Foundation.

References

- Scheerder, J.; van Duynhoven, J. P. M.; Engbersen, J. F. J.; Reinhoudt, D. N. *Angew. Chem. Int. Ed. Eng.* **1996**, *35*, 1090.
- Beer, P. D.; Cooper, J. B. *Chem. Commun.* **1998**, 129.
- Kang, S. O.; Nam, K. C. *Bull. Korean Chem. Soc.* **2000**, *21*, 461.
- Kang, S. O.; Nam, K. C. *Bull. Korean Chem. Soc.* **2002**, *23*, 640.
- (a) Reddy, P. A.; Kashap, R. P.; Qatson, W. H.; Gutsche, C. D. *Israel J. Chem.* **1992**, *32*, 89. (b) Reddy, P. A.; Gutsche, C. D. *J. Org. Chem.* **1993**, *58*, 3245. (c) Gomez-Kaifer, M.; Reddy, P. A.; Gutsche, C. D.; Echegoyen, L. *J. Am. Chem. Soc.* **1994**, *115*, 3580.
- (a) Beer, P. D.; Chen, Z.; Drew, M. G. B.; Gale, P. A. *J. Chem. Soc., Chem. Comm.* **1994**, 2207. (b) Chen, Z.; Gale, P. A.; Beer, P. D. *J. Chem. Soc. Faraday Trans.* **1994**, *90*, 2207. (a) Beer, P. D.; Chen, Z.; Gale, P. A. *Tetrahedron* **1994**, *50*, 931.
- (a) Nam, K. C.; Kang, S. O.; Jeong, H. S.; Jeon, S. *Tetrahedron Lett.* **1999**, *40*, 7343. (b) Cho, E. J.; Hwang, S. S.; Oh, J. M.; Lee, H. K.; Jeon, S.; Nam, K. C. *Bull. Korean Chem. Soc.* **2001**, *22*, 782. (c) Kang, S. O.; Oh, J. M.; Yang, Y. S.; Chun, J. C.; Jeon, S.; Nam, K. C. *Bull. Korean Chem. Soc.* **2002**, *23*, 145. (d) Jeong, H. S.; Choi, E. M.; Kang, S. O.; Nam, K. C.; Jeon, S. *J. Electroanal. Chem.* **2000**, *485*, 154.
- Collins, E. M.; McKervey, M. A.; Madigan, E.; Moran, M. B.; Owens, M.; Ferguson, G.; Harris, S. J. *J. Chem. Soc. Perkin Trans. 1* **1991**, 3137.
- Jaime, C.; Mendoza, J. D.; Parados, P.; Nieto, P. M.; Sanchez, C. J. *Org. Chem.* **1991**, *56*, 3372.
- Hynes, M. J. *J. Chem. Soc. Dalton Trans.* **1993**, 311.
- Gutsche, C. D.; Iqbal, M.; Stewart, D. *J. Org. Chem.* **1986**, *51*, 742.