

Kinetics and Mechanism of the Addition of Benzylamines to Benzylidene Meldrum's Acids in Acetonitrile

Hyuck Keun Oh,[†] Tae Soo Kim,[†] Hai Whang Lee,^{*} and Ikchoon Lee^{*}

[†]Department of Chemistry, Chonbuk National University, Jeonju 560-756, Korea

Department of Chemistry, Inha University, Incheon 402-751, Korea


Received November 14, 2002

Nucleophilic addition reactions of benzylamines (XC₆H₄CH₂NH₂) to benzylidene Meldrum's acids (BMA; YC₆H₄CH=C(COO)₂C(CH₃)₂) have been investigated in acetonitrile at 20.0 °C. The rates of addition are greatly enhanced due to the abnormally high acidity of Meldrum's acid. The magnitudes of the Hammett (ρ_X and ρ_Y) and Brønsted (β_X) coefficients are rather small suggesting an early transition state. The sign and magnitude of the cross-interaction constant, ρ_{XY} ($= -0.33$), and kinetic isotope effects ($k_H/k_D \cong 1.5$ -1.7) involving deuterated benzylamine nucleophilites (XC₆H₄CH₂ND₂) are indicative of hydrogen-bonded cyclic transition state. The activation parameters, $\Delta H^\ddagger \cong 4$ kcal mol⁻¹ and $\Delta S^\ddagger \cong -37$ e.u., are also in line with the proposed mechanism.


Key Words : Benzylidene Meldrum's acid, Nucleophilic addition reaction, Cross-interaction constant

Introduction

The nucleophilic attack to carbon-carbon double bonds has been the subject of numerous investigations.¹ Nucleophilic vinylic substitution reactions activated by electron-withdrawing groups (Z,Z') are known to proceed by the two-step mechanism, eq. (1), where LG = leaving group and Nu = nucleophile.¹ The primary addition step, k_a , leads to an association complex or intermediate which for amine base is a zwitterion, T[±].




In the simple nucleophilic vinylic additions, however, when the amines are primary or secondary, intramolecular proton transfer may immediately follow or even simultaneously upon formation of T[±].^{1b,2,3} eq. (2). Previous works




have shown that in aqueous solution^{1b,1} the mechanism of nucleophilic addition to activated olefins is complex involving group separated products formation following the rate-limiting addition step, k_a , with imbalanced transition states in which the development of resonance to the activating groups (Z,Z') lags behind charge transfer or N-C bond formation. In contrast, the amine addition in acetonitrile³ was found to be a relatively simple one, *i.e.*, one step addition.

In this work we report kinetic study of the nucleophilic

addition of benzylamines to benzylidene Meldrum's acids (BMAs) in acetonitrile at 20.0 °C, eq. (3). Meldrum's acid 1,



has attracted considerable attention due to its high acidity,⁴ higher ($\text{p}K_a = 7.3$) than its acyclic analogues such as dimethyl malonate (2; $\text{p}K_a = 15.9$) or even the diketone analogue (3; $\text{p}K_a = 11.2$).^{4b} The nucleophilic addition reactions to BMA in aqueous solution have been investigated extensively by Bemasoni *et al.*⁵ They reported some unusual kinetic behaviors in the addition reaction of BMA.



The primary purpose of this study is to elucidate the mechanism of primary amine addition to BMA in acetonitrile. Since the Meldrum's acid in BMA is known to have unusually strong power of stabilizing the anion formed in the primary attack,^{2-4,5} we are interested especially in the rates of addition (k_2 step in eq. (3)) and in the degree of bond formation in the TS by determining the Hammett (ρ_X) and Brønsted coefficients (β_X), and cross-interaction constant,⁶ ρ_{XY} in eqs. (4), where X and Y are the substituents in the nucleophile (benzylamine) and substrate (BMA), respectively.

$$\log(k_{XY}/k_{HH}) = \rho_X\sigma_X + \rho_Y\sigma_Y + \rho_{XY}\sigma_X\sigma_Y \quad (4a)$$

$$\rho_{XY} - \partial\rho_Y/\partial\sigma_X - \partial\rho_X/\partial\sigma_Y \quad (4b)$$

In aqueous solution, the magnitude of $\beta_X(\beta_{\text{inc}})$ determined with piperidine and morpholine pair has been shown to be exceptionally small ($\beta_X = 0.07$)^{5a} but that with primary amines is somewhat larger with $\beta_X = 0.22$.^{5c} It is of interest to find out whether such unusual behaviors in aqueous solution persist in acetonitrile or not.

Results and Discussion


A clean second-order kinetics according to eqs. (5) and (6) was obtained throughout this work, where k_2 is the second-order rate constant for the benzylamine (BnA) addition to

$$\text{Rate} = k_{\text{obsd}} [\text{BMA}] \quad (5)$$

$$k_{\text{obsd}} = k_2 [\text{BnA}] \quad (6)$$

the benzylidene Meldrum's acid (BMA) under pseudo-first-order condition (k_{obsd}) with large excess of amine nucleophiles. Plots of k_{obsd} against [BnA] were linear for *ca.* 10-fold increase in [BnA]. The k_2 values determined from the slopes of these plots are summarized in Table 1. The rates are compared with those involving other activating groups, Z,Z', in Table 2. The rates in acetonitrile are much slower than those ($\log k_0$) in aqueous solution. Excepting for the

amine addition to BMA, the reactivity in aqueous solution increases with the (polar) electron-withdrawing power ($\Sigma\sigma$)⁷ of the activating groups whereas that in acetonitrile depends on the through conjugative electron-withdrawing strength ($\Sigma\sigma^-$)⁷ of the activating groups. We note that the reactivity of BMA is exceptionally large in aqueous solution as well as in acetonitrile. The $\Sigma\sigma^- 0.75$ should place the reactivity of BMA to the lowest place but the $\log k_0$ value is the second after that of BMN for which $\Sigma\sigma^- 1.32$. Again the $\Sigma\sigma^- 1.38$ should give the reactivity of BMA in acetonitrile less than that of BID ($\Sigma\sigma^- 2.08$) but the reactivity of BMA in acetonitrile is the largest. The enhanced reactivity of the amine addition to BMA no doubt reflects the unusually strong acidity of the Meldrum's acid due to strong stabilizing power of the Meldrum's acid anion,^{2,4,5} in which the anionic charge is delocalized, **1A**.



1A

The anomalously high acidity of **1** has been a subject of extensive theoretical as well as experimental investigations.

Table 1. The second-order rate constants, k_2 ($\text{M}^{-1}\text{s}^{-1}$), and selectivity parameters^a for the addition of X-benzylamines to Y-benzylidene Meldrum's acids in acetonitrile at 20.0 °C

X	Y				ρ_{XY}^b
	<i>p</i> -OMe	<i>p</i> -Me	H	<i>p</i> -Cl	
<i>p</i> -OMe	74.4 48.7 ^c 32.4 ^d	86.4	101	132 85.8 56.3	0.49 ± 0.02
<i>p</i> -Me	71.9	80.3	93.5	117	0.42 ± 0.01
H	67.2	73.9	86.6	106	0.40 ± 0.01
<i>p</i> -Cl	62.6 42.1 28.4	68.0	76.6	91.0 60.3 40.6	0.32 ± 0.01
<i>m</i> -Cl	59.3	62.1	70.4	78.9	0.26 ± 0.02
ρ_X^c	-0.15 ± 0.01	-0.21 ± 0.01	-0.24 ± 0.01	-0.33 ± 0.02	ρ_{XY}^d
β_X^e	0.15 ± 0.01	0.21 ± 0.01	0.23 ± 0.05	0.32 ± 0.02	= -0.33 ± 0.03

^aThe σ values were taken from ref. 7. The $\text{p}K_a$ values were taken from ref. 8. The $\text{p}K_a = 9.67$ was used for X-*p*-ClH₃O (ref. 9). ^bCorrelation coefficients were better than 0.996 in all cases. ^cAt 10.0 °C. ^dAt 0.0 °C. ^eCorrelation coefficients were better than 0.994 in all cases. ^fCorrelation coefficients were better than 0.993 in all cases. ^gCorrelation coefficient was 0.998.

Table 2. Comparisons of reactivity parameter for the addition reaction, $\text{YC}_6\text{H}_4\text{CH}=\text{CZ}' + \text{XC}_6\text{H}_4\text{CH}_2\text{NH}_2$, in acetonitrile


Entry	Z,Z'	$k_2^a/\text{M}^{-1}\text{s}^{-1}$	$\log k_0^b$	ρ_X^c	ρ_Y^c	ρ_{XY}^d	$\Sigma\sigma^e$	$\Sigma\sigma^f$
1 (BMA) ^g	(COO) ₂ CMe ₂	86.6 (20 °C)	≈4.10	-0.24	0.40	-0.33	0.75	1.38
2 (BMN) ^h	CN, CN	1.48 ⁱ	4.94	-1.62	-0.55	-0.31	1.32	2.00
3 (BID) ^j	(CO) ₂ C ₆ H ₄	1.48	—	-1.10	0.41	-0.33	0.83	2.08
4 (NS) ^k	NO ₂ , H	2.63×10^{-2}	2.55	-1.22	1.73	-0.40	0.78	1.27
5 (NSB) ^l	NO ₂ , C ₆ H ₅	2.69×10^{-2}	1.42	-0.82	1.27	-0.52	0.77	1.27 ^m
6 (CNS) ⁿ	CN, <i>p</i> -NO ₂ C ₆ H ₄	1.26×10^{-1} (30 °C)	≈3.35	-1.15	1.10	-0.67	0.92	1.00 ^o

^aFor X=Y=H at 25.0 °C unless otherwise noted in parentheses. ^bIntrinsic rate constants, k_0 , for carbanion forming reactions (k_2 in eq. 2) in 50% DMSO-50% H₂O at 20.0 °C with amines (refs. 1b and 5d). ^cFor Y=H and X=H, respectively. ^dCorrelation coefficients were better than 0.997 in all cases. ^eNormal Hammett substituent constant (σ_p). ^fExalted substituent constant (σ_p^-) for direct conjugation with anionic functional center (ref. 7). ^gThis work. ^hBenzylidenemalononitrile (ref. 3b). ⁱBenzylidene-1,3-indandione (ref. 3c). ^j β -Nitrostyrene (ref. 3a). ^k β -Nitrostilbene and β -cyano- β -4'-nitrostilbene (ref. 3d). ^lExtrapolated value. ^mThe value of Ph group is excepted (ref. 3d). ⁿThe value of *p*-NO₂Ph group is excepted (ref. 3d).

Arnett *et al.*^{4a} measured acidity constants of esters and lactones that are relevant to the Meldrum's acid and attributed the strong acidity of **1** to restricted rotation which faces the ester groups to adopt unfavorable anti conformation in Meldrum's acid. Wiberg *et al.*^{4c} studied the acidity difference between syn and anti rotamers of methyl acetate theoretically at the MP3/6-31G**//HF/6-31G* level¹⁰ and concluded that the change in charge distribution that accompanies the more facile loss of proton from the anti rotamer (by 4.7 kcal mol⁻¹) leads to the unusual acidity of Meldrum's acid, having a bis (anti) conformation. Wang and Houk^{4b} also reported ab initio calculations on the acidity of methyl acetate in syn and anti conformations at the HF/6-31+G**//HF/3-21G level.¹⁰ They attributed the strong acidity of the anti form (Meldrum's acid has a bis (anti) form) to greater dipole-dipole electrostatic repulsion interaction between the ether and carbonyl oxygen atoms in the anti conformation, which is significantly relieved in the corresponding enolate anion.

The Hammett ($\rho_X = -0.15 \sim -0.33$) and Brønsted coefficients ($\beta_X = 0.15-0.32$) in Table 1 are similar to those obtained in the corresponding reactions with primary amines in aqueous solution ($\rho_X = -0.22$).^{5c} The β_X values are greater than that obtained with secondary amines ($\beta_X = 0.07$ ^{5a} by two points correlation using piperidine and morpholine) in aqueous solution, but are rather smaller still than those for other benzylamine addition reactions.³ (although the β_X values are not listed in Table 2, they are numerically quite similar to the ρ_X values in Table 2 due to the near unity slope in the correlation between the pK_a 's of benzylamine and Hammett substituent constants, σ ; *i.e.*, $\delta pK_a/\delta\sigma \cong 1.06$ for benzylamines.¹¹) The larger β_X value with primary amines than with secondary and tertiary amines seems to represent a common pattern, albeit there are some exceptions.^{5c} For example, Skoog and Jencks¹² reported $\beta_X = 0.19$ and 0.17 for the reactions of primary amines and pyridines with phosphorylated 3-methoxypyridine and $\beta_X = 0.28$ and 0.22 for the corresponding reactions with phosphorylated 4-morpholinopyridine, respectively. The aminolysis of isoquinoline *N*-phosphonate with primary amines and piperidine-morpholine pair gave $\beta_X = 0.15$ and 0.0, respectively.¹³ For the aminolysis of *S*-(2,4-dinitrophenyl) *O*-ethyl carbonate, EtOC(=O)SDNP, $\beta_X = 0.56$ ¹⁴ and 0.48¹⁵ are obtained with secondary alicyclic amines and quinuclidines, respectively. The ρ_Y values are also small so that charge transfer from the benzylamine nucleophiles to the vinylic carbon in the TS may be small.

On the other hand, the cross-interaction constant, ρ_{XY} ($= -0.33$), is negative as in other bond forming processes⁶ and the magnitude is similar to those for other relatively faster addition reactions with stronger activating groups, entries 2 (BMN: $\rho_{XY} = -0.31$) and 3 (BID: $\rho_{XY} = -0.33$) in Table 2. The modest size of the ρ_{XY} value despite the relatively early TS with a lower degree of bond making (lower β_X values) may point to a possibility of intramolecular hydrogen bonding¹⁶ to an oxygen of the (COO)₂-C(CH₃)₂ moiety, **6**, or to the central carbon atom, **7**, since in these types of hydrogen bonded TS structures the interaction



between X and Y can be through two routes so that the intensity of interaction, *i.e.*, the magnitude of ρ_{XY} , can have a modest value even though the C-N bond making is small.¹⁶ Normally the amine nucleophiles approach toward an end carbon of the vinylic double bond from almost directly above (or below) the carbon so that distancewise the H...C in **7** may be nearer than H...O in **6**.

The kinetic isotope effects involving deuterated benzylamine nucleophiles (XC₆H₄CH₂ND₂) in Table 3 are all greater than unity ($k_H/k_D = 1.5-1.7$) and indeed supports such hydrogen bonded TS structure, **6** and **7**. The size of k_H/k_D increases with a stronger electron acceptor substituent in the substrate (Y) as well as in the nucleophile (X) which is in accord with a greater degree of bond making in the TS with electron acceptor Y (with a greater β_X value) and the enhanced acidity of the NH proton with an electron-acceptor substituent X in the nucleophile. The overall size of the k_H/k_D values ranging 1.5-1.7 are, however, smaller than those of the corresponding values for the substrates with weaker electron acceptor Z,Z' exhibiting slower addition rates, *e.g.* for entries 4, 5, and 6 in Table 2 the k_H/k_D values are larger ranging 2.2-3.1.^{3a,d} This is in line with the greater degree of bond formation in the TS for those substrates (entries 4-6) with larger magnitude of the cross-interaction constants, ρ_{XY}

Table 3. Kinetic isotope effects on the second-order rate constants for the addition of deuterated X-benzylamines (XC₆H₄CH₂ND₂) to Y-benzylidene Meldrum's acids in acetonitrile at 20.0 °C

X	Y	$k_H/M^{-1}s^{-1}$	$k_D/M^{-1}s^{-1}$	k_H/k_D
<i>p</i> -OMe	<i>p</i> -OMe	74.4 (± 0.1)	51.3 (± 0.5)	1.45 ± 0.02 ^a
<i>p</i> -OMe	<i>p</i> -Me	86.4 (± 0.9)	57.6 (± 0.5)	1.50 ± 0.02
<i>p</i> -OMe	H	101 (± 1)	64.7 (± 0.8)	1.56 ± 0.03
<i>p</i> -OMe	<i>p</i> -Cl	132 (± 2)	81.5 (± 0.9)	1.62 ± 0.03
<i>p</i> -Cl	<i>p</i> -OMe	62.6 (± 0.8)	40.9 (± 0.4)	1.53 ± 0.03
<i>p</i> -Cl	<i>p</i> -Me	68.0 (± 0.9)	42.8 (± 0.4)	1.59 ± 0.03
<i>p</i> -Cl	H	76.6 (± 0.9)	45.6 (± 0.4)	1.68 ± 0.02
<i>p</i> -Cl	<i>p</i> -Cl	91.0 (± 0.1)	52.3 (± 0.5)	1.74 ± 0.02

^aStandard deviations.

Table 4. Activation parameters^a for the addition of X-benzylamines to Y-benzylidene Meldrum's acids in acetonitrile

X	Y	$\Delta H^\ddagger/kcal\ mol^{-1}$	$-\Delta S^\ddagger/cal\ mol^{-1}K^{-1}$
<i>p</i> -OMe	<i>p</i> -OMe	3.9	37
<i>p</i> -OMe	<i>p</i> -Cl	4.1	36
<i>p</i> -Cl	<i>p</i> -OMe	3.8	38
<i>p</i> -Cl	<i>p</i> -Cl	3.8	37

^aCalculated by the Eyring equation. The maximum errors calculated (by the method of Wiberg, K. B. ref. 17) are = 0.5 kcal mol⁻¹ and = 2 e.u. for ΔH^\ddagger and ΔS^\ddagger , respectively.

As expected from a hydrogen bonded cyclic TS, the enthalpies of activation are low, $\Delta H^\ddagger \cong 4 \text{ kcal mol}^{-1}$, and the entropies of activation are large negative $\Delta S^\ddagger \cong -37 \text{ e.u.}$ (Table 4). The addition step is facilitated by the simultaneous hydrogen bonding (low ΔH^\ddagger) but the TS is constrained (large negative ΔS^\ddagger) in a cyclic structure.

Conclusion

The reactions of benzylamine additions to benzylidene Meldrum's acids (BMA) are unusually fast in acetonitrile, which can be ascribed to the abnormally high acidity of the Meldrum's acid. The reactions proceed by simultaneous addition of amine and proton to the vinylic double bond with hydrogen bonded cyclic transition state. Relatively low β_X , ρ_X and ρ_Y values are interpreted to indicate an early transition state. The normal kinetic isotope effects ($k_H/k_D = 1.5\text{--}1.7$) involving deuterated benzylamine nucleophiles, a modest size of the negative cross-interaction constant, $\rho_{XY} = -0.33$, and activation parameters support the proposed mechanism.

Experimental Section

Materials. Merck GR acetonitrile was used after three distillations. The benzylamine nucleophiles, Aldrich GR, were used after recrystallization. Toluene, Meldrum's acid and piperidine were Aldrich reagents.

Preparation of Benzylidene Meldrum's Acids. The benzylidene Meldrum's acids were prepared by the literature method of Schuster *et al.*¹⁸ A solution of Meldrum's acid (3 mmol) and benzaldehyde (3 mmol) in toluene was treated with a few drops of piperidine and refluxed for 3 h. The solution was cooled, some of the toluene was evaporated, and the dark-colored solid was removed by filtration to obtain (85%) crude material. This was recrystallized from toluene. Melting points, IR and ¹H and ¹³C NMR (400 MHz) data were found to agree well with the literature values.¹⁸

Kinetic Measurements. The reaction was followed spectrophotometrically by monitoring the decrease in the concentration of benzylidene Meldrum's acids, [BMA], at λ_{max} of the substrate to over 80% completion. The reaction was studied under pseudo-first-order condition, [BMA] $\cong 6.0 \times 10^{-5} \text{ M}$ and [BnA] = $\sim 10^{-3} \text{ M}$ at $20.0 \pm 0.1 \text{ }^\circ\text{C}$. The pseudo first-order rate constant, k_{obsd} , was determined from the slope of the plot ($r > 0.993$) of $\ln [\text{BMA}]$ vs time. Second-order rate constants, k_2 , were obtained from the slope of a plot ($r > 0.995$) of k_{obsd} vs [BnA] with more than five concentrations of benzylamine and were reproducible to within $\pm 3\%$.

Product Analysis. The analysis of final products was difficult due to partial decomposition during product separation and purification. We therefore analyzed the addition reaction of *p*-MeOC₆H₄CH₂NH₂ to *p*-ClC₆H₄CH = C(COO)₂C(CH₃)₂ by ¹³C NMR (400 MHz) at appropriate intervals under exactly the same reaction conditions as the kinetic measurement in CD₃CN at 30.0 $^\circ\text{C}$. Initially we found a peak for CH in the

reactant, *p*-ClC₆H₄CH = C(COO)₂C(CH₃)₂, at 7.62 ppm, which was gradually reduced, and two new peaks for CH-CH in the product, *p*-ClC₆H₄(*p*-MeOC₆H₄CH₂NH)CH-CH(COO)₂-C(CH₃)₂, grew at 3.93 and 4.69 ppm as the reaction proceeded. No other peaks or complications were found during the reaction except the 3 peak height changes indicating that the reaction proceeds with no other side reactions.

Acknowledgment. This work was supported by Korea Research Foundation Grant (KRF-2002-070-C00061).

References

- (a) Patai, S.; Rappoport, Z. In *The Chemistry of Alkenes*; Patai, S., Ed.; Interscience: New York, 1964; p 469. (b) Bernasconi, C. F. *Tetrahedron* **1989**, *45*, 4017. (c) Rappoport, Z.; Ladhani, D. *Chem. Scr.* **1974**, *5*, 124. (d) Rappoport, Z. *Adv. Phys. Org. Chem.* **1969**, *7*, 1. (e) Modena, G. *Acc. Chem. Res.* **1971**, *4*, 73. (f) Müller, S. I. *Tetrahedron* **1977**, *33*, 1211. (g) Rappoport, Z. *Acc. Chem. Res.* **1981**, *14*, 7. (h) Rappoport, Z. *Acc. Chem. Res.* **1992**, *25*, 474. (i) Bernasconi, C. F. *Acc. Chem. Res.* **1987**, *20*, 301.
- Schreiber, B.; Martinek, H.; Wolscham, P.; Schuster, P. *J. Am. Chem. Soc.* **1979**, *101*, 4708.
- (a) Oh, H. K.; Yang, J. H.; Sung, D. D.; Lee, I. *J. Chem. Soc., Perkin Trans. 2* **2000**, 101. (b) Oh, H. K.; Yang, J. H.; Lee, H. W.; Lee, I. *J. Org. Chem.* **2000**, *65*, 2188. (c) Oh, H. K.; Yang, J. H.; Lee, H. W.; Lee, I. *J. Org. Chem.* **2000**, *65*, 5391. (d) Oh, H. K.; Kim, T. S.; Lee, H. W.; Lee, I. *J. Chem. Soc., Perkin Trans. 2* **2002**, 282.
- (a) Arnett, E. M.; Harrelson, J. A. Jr. *J. Am. Chem. Soc.* **1987**, *109*, 809. (b) Wang, X.; Houk, K. N. *J. Am. Chem. Soc.* **1988**, *110*, 1870. (c) Wiberg, K. B.; Laidig, K. E. *J. Am. Chem. Soc.* **1988**, *110*, 1872.
- (a) Bernasconi, C. F.; Fornarini, S. *J. Am. Chem. Soc.* **1980**, *102*, 5329. (b) Bernasconi, C. F.; Leonarduzzi, G. D. *J. Am. Chem. Soc.* **1982**, *104*, 5133. (c) Bernasconi, C. F.; Murray, C. J. *J. Am. Chem. Soc.* **1986**, *108*, 5251. (d) Bernasconi, C. F.; Panda, N. *J. Org. Chem.* **1987**, *52*, 3042. (e) Bernasconi, C. F.; Ketner, R. J. *J. Org. Chem.* **1998**, *63*, 6266.
- (a) Lee, I. *Adv. Phys. Org. Chem.* **1992**, *27*, 57. (b) Lee, I.; Lee, H. W. *Collect. Czech. Chem. Commun.* **1999**, *64*, 1529. (c) Lee, I. *Chem. Soc. Rev.* **1990**, *19*, 317.
- Hansch, C.; Leo, A.; Taft, R. W. *Chem. Rev.* **1991**, *91*, 165.
- Blackwell, L. F.; Fischer, A.; Miller, I. J.; Topsom, R. D.; Vaughan, J. J. *J. Chem. Soc.* **1964**, 3588.
- Oh, H. K.; Lee, J. Y.; Lee, I. *Bull. Korean Chem. Soc.* **1998**, *19*, 1198.
- (a) Hehre, W. J.; Radom, L.; Schleyer, P. v. R.; Pople, J. A. *Ab Initio Molecular Orbital Theory*; Wiley: New York, 1986; Chapter 4. (b) Foresman, J. B.; Frisch, E. *Exploring Chemistry with Electronic Structure Methods*, 2nd Ed.; Gaussian Inc.: Pittsburgh, 1996; Part 2.
- Page, M.; Williams, A. *Organic and Bio-organic Mechanisms*; Longman: Harlow, 1997; p 259.
- Skoog, M. T.; Jencks, W. P. *J. Am. Chem. Soc.* **1984**, *106*, 7597.
- Bourne, N.; Williams, A. *J. Am. Chem. Soc.* **1984**, *106*, 7591.
- Castro, E. A.; Ibanez, F.; Salas, M.; Santos, J. G. *J. Org. Chem.* **1991**, *56*, 4819.
- Castro, E. A.; Munoz, P.; Santos, J. G. *J. Org. Chem.* **1999**, *64*, 8298.
- Lee, I. *Chem. Soc. Rev.* **1995**, *24*, 223.
- Wiberg, K. B. *Physical Organic Chemistry*; Wiley: New York, 1964; p 378.
- Schuster, P.; Polansky, O. E.; Wessely, F. *Monatsh. Chem.* **1963**, *95*, 53.