

## Antioxidant Principles of *Nelumbo nucifera* Stamens

Hyun Ah Jung, Jung Eun Kim, Hae Young Chung<sup>1</sup>, and Jae Sue Choi

Faculty of Food Science and Biotechnology, Pukyong National University, Busan 608-737, Korea and <sup>1</sup>College of Pharmacy, Pusan National University, Busan 609-735, Korea

(Received January 22, 2003)

In our ongoing study to identify antioxidants from natural sources, the antioxidant activity of *Nelumbo nucifera* stamens was evaluated for their potential to scavenge stable 1,1-diphenyl-2-picrylhydrazyl (DPPH) free radicals, inhibit total reactive oxygen species (ROS) generation, in kidney homogenates using 2',7'-dichlorodihydrofluorescein diacetate (DCHF-DA), and scavenge authentic peroxyntrites (ONOO<sup>-</sup>). A methanol (MeOH) extract of the stamens of *N. nucifera* showed strong antioxidant activity in the ONOO<sup>-</sup> system, and marginal activity in the DPPH and total ROS systems, so were therefore fractionated with several organic solvents, such as dichloromethane (CH<sub>2</sub>Cl<sub>2</sub>), ethyl acetate (EtOAc) and *n*-butanol (*n*-BuOH). The EtOAc soluble fraction, which exhibited strong antioxidant activity in all the model systems tested, was further purified by repeated silica gel and Sephadex LH-20 column chromatographies. Seven known flavonoids [kaempferol (**1**), kaempferol 3-O-β-D-glucuronopyranosyl methylester (**2**), kaempferol 3-O-β-D-glucopyranoside (**3**), kaempferol 3-O-β-D-galactopyranoside (**4**), myricetin 3',5'-dimethylether 3-O-β-D-glucopyranoside (**5**), kaempferol 3-O-α-L-rhamnopyranosyl-(1→6)-β-D-glucopyranoside (**6**) and kaempferol 3-O-β-D-glucuronopyranoside (**7**)], along with β-sitosterol glucopyranoside (**8**), were isolated. Compound **1** possessed good activities in all the model systems tested. Compounds **2** and **7** showed scavenging activities in the DPPH and ONOO<sup>-</sup> tests, while compounds **3** and **4** were only active in the ONOO<sup>-</sup> test. Conversely, compound **8** showed no activities in any of the model systems tested.

**Key words:** *Nelumbo nucifera*, Nymphaeaceae, Flavonoids, Antioxidant, Free radical, 1,1-Diphenyl-2-picrylhydrazyl (DPPH) radical, Reactive oxygen species (ROS), Peroxyntrite (ONOO<sup>-</sup>)

### INTRODUCTION

Reactive oxygen and nitrogen species (ROS and RNS, respectively), such as superoxide anion radical (·O<sub>2</sub><sup>-</sup>), hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>), hydroxyl radical (·OH), singlet oxygen (¹O<sub>2</sub>), alkoxyl radical (RO·), peroxy radical (ROO·) and peroxyntrite (ONOO<sup>-</sup>), are generated under normal and/or pathological conditions. These ROS and RNS are likely to damage several cellular molecules, such as lipids, proteins, nucleic acids and DNA, due to oxidation and nitration, and cause inflammation, or lesions, of various organs (Beckman *et al.*, 1990). Also, these reactive species are associated with various degenerative diseases, such as cancer (Dreher *et al.*, 1996), aging (Sohal, 2002), and arteriosclerosis and rheumatism (Griffiths and Lunec, 1996; Squadrito and Pryor, 1998; Choi *et al.*, 2002).

There has been much research on the powerful, nontoxic natural antioxidants from edible or medicinal plants to prevent not only these oxidative stress-induced disorders in humans, but also the oxidative deterioration of food by auto-oxidation and lipid peroxidation. Many naturally occurring compounds from plant sources have also been reported to retard the oxidation process, in their natural environment, by acting as free radical scavengers, reactive oxygen scavengers or as reducing agents *in vitro* (Pratt, 1994; Larson, 1988).

In our ongoing investigation to identify natural antioxidants, we have dedicated much time to the study of *Nelumbo nucifera* Gaertner, from the Nymphaeaceae family. This edible, medicinal plant is an aquatic, perennial herb, which is distributed in all over the world, including Korea (Van Bergen *et al.*, 1997; Kim, 1996). Due to its various therapeutic effects, it has been used as a traditional folk medicine for the treatment of diarrhea, gastritis, insomnia, nervous prostration and as a haemostatic in Korea and other countries, such as China and India (Kim, 1997; La Cour *et al.*, 1995). The biological activities of *N. nucifera*,

Correspondence to: Jae Sue Choi, Faculty of Food Science and Biotechnology, Pukyong National University, Busan 608-737, Korea  
Tel: 82-51-620-6335. Fax: 82-51-620-6330  
E-mail: choijs@pknu.ac.kr

such as anti-diarrheic (Mukjerjee *et al.*, 1995), psychopharmacological (Mukjerjee *et al.*, 1996), hypoglycemic (Mukjerjee *et al.*, 1997), hypolipemic (La Cour *et al.*, 1995), anti-pyretic (Sinha *et al.*, 2000) and antioxidant activities (Liou *et al.*, 1999; Guo *et al.*, 2000) have all previously been reported.

In the present study, the antioxidant activity of the methanol (MeOH) extract, and its organic solvent soluble fractions, such as those from dichloromethane (CH<sub>2</sub>Cl<sub>2</sub>), ethyl acetate (EtOAc), *n*-butanol (*n*-BuOH), and the water (H<sub>2</sub>O) layer, from the stamens of *N. nucifera*, were evaluated by the 1,1-diphenyl-2-picrylhydrazyl (DPPH) radical, total ROS and ONOO<sup>-</sup> scavenging/inhibitory tests. The isolation and identification of seven known flavonoids (1-7), together with compound 8 from the active EtOAc fraction of *N. nucifera* stamens was also investigated. Furthermore, the antioxidant properties of these isolated flavonoids (1-4 and 7) were measured by three model systems, those being the DPPH radical, total ROS and ONOO<sup>-</sup> model systems.

## MATERIALS AND METHODS

### Plant materials

The stamens of *N. nucifera* were purchased in April 2000 from the herbal medicine co-operative association of Busan Province, Korea, and authenticated by Hae Young Chung, professor of the College of Pharmacy, Pusan National University. A voucher specimen has been deposited in the Herbarium of the Medicinal Plant Garden, the College of Pharmacy, Pusan National University.

### Chemicals

The DPPH, L-ascorbic acid, Trolox and DL-penicillamine (DL-2-amino-3-mercapto-3-methylbutanoic acid) were purchased from Sigma Chemical Company (St. Louis, MO, USA). The high quality DCFH-DA and DHR 123 (dihydrorhodamine 123), and ONOO<sup>-</sup> were purchased from Molecular Probes (Eugene, Oregon, USA) and Cayman (Ann Arbor, MI, USA), respectively.

### General experimental procedures

The <sup>1</sup>H- and <sup>13</sup>C-NMR spectra were recorded on a Varian UNITY-400 spectrometer. Chemical shifts were referenced to the respective residual solvent peaks, and the values recorded in δ. The HMQC and HMBC spectra were recorded using pulsed field gradients, rather than phase cycling, for coherence pathway selection. The multiplicities of the <sup>1</sup>H- and <sup>13</sup>C-NMR signals are indicated as s (singlet), d (doublet) and m (multiplet). The column chromatography was performed with silica gel (Merck, 70-230 mesh), and the TLC on precoated Merck Kieselgel 60 F<sub>254</sub> plate (0.25 mm), and spots detected using 50% H<sub>2</sub>SO<sub>4</sub> reagent.

### Isolation of compounds 1-8

The dried *Nelumbo nucifera* stamens (3.4 kg) were refluxed with MeOH for three hours. The total filtrate was concentrated to dryness, *in vacuo* at 40°C, to render the MeOH extract (560 g). This extract was suspended in H<sub>2</sub>O and then successively partitioned with CH<sub>2</sub>Cl<sub>2</sub>, EtOAc and *n*-BuOH, to afford the CH<sub>2</sub>Cl<sub>2</sub> (330 g), EtOAc (20 g), and *n*-BuOH fractions (66 g), and the H<sub>2</sub>O residue (140 g). The EtOAc extract (20 g) was fractionated with Sephadex LH-20, using MeOH, to obtain 5 fractions (Fr. 1-Fr. 5). Fractions 1 (3.5 g) and 2 (5.7 g) were separately purified by Sephadex LH-20, with MeOH, to obtain compounds 8 (100 mg) and 5 (25 mg). Fraction 3 (9.7 g) was subjected to column chromatography on a Si gel column, with gradient elution, using an EtOAc:MeOH eluent (10:1, gradient to MeOH), to give compounds 2 (20 mg), 3 (450 mg), 4 (180 mg) and 6 (45 mg). Fractions 4 (1.5 g) and 5 (2.0 g) were recrystallized to obtain compounds 7 (80 mg) and 1 (260 mg), respectively.

#### Kaempferol (1)

<sup>1</sup>H-NMR (400 MHz, DMSO-*d*<sub>6</sub>) δ: 6.19 (1H, d, *J* = 2.0 Hz, H-6), 6.44 (1H, d, *J* = 2.0 Hz, H-8), 6.92 (2H, d, *J* = 8.0 Hz, H-3', 5'), 8.06 (2H, d, *J* = 8.0 Hz, H-2', 6'), 12.47 (1H, brs, 5-OH); <sup>13</sup>C-NMR (100 MHz, DMSO-*d*<sub>6</sub>) δ: 175.9 (C-4), 163.9 (C-7), 160.7 (C-5), 159.2 (C-4'), 156.2 (C-9), 146.8 (C-2), 135.7 (C-3), 129.5 (C-2', 6'), 121.7 (C-1'), 115.4 (C-3', 5'), 103.0 (C-10), 98.2 (C-6), 93.5 (C-8).

#### Kaempferol 3-O-β-D-glucuronopyranosyl methyl ester (2)

<sup>1</sup>H-NMR (400 MHz, DMSO-*d*<sub>6</sub>) δ: 3.57 (1H, s, OMe), 5.47 (1H, d, *J* = 7.4 Hz, H-1''), 6.21 (1H, d, *J* = 2.0 Hz, H-6), 6.43 (1H, d, *J* = 2.0 Hz, H-8), 6.89 (2H, d, *J* = 8.0 Hz, H-3', 5'), 8.02 (2H, d, *J* = 8.0 Hz, H-2', 6'), 12.51 (1H, brs, 5-OH); <sup>13</sup>C-NMR (100 MHz, DMSO-*d*<sub>6</sub>) δ: 177.2 (C-4), 169.0 (C-6''), 164.3 (C-7), 161.2 (C-5), 160.2 (C-4'), 156.6 (C-9), 156.4 (C-2), 133.0 (C-3), 130.9 (C-2', 6'), 120.6 (C-1'), 115.1 (C-3', 5'), 104.0 (C-10), 101.4 (C-1''), 98.9 (C-6), 93.8 (C-8), 75.6 (C-5''), 75.5 (C-3''), 73.9 (C-2''), 71.5 (C-4''), 51.9 (OMe).

#### Kaempferol 3-O-β-D-glucopyranoside (3)

<sup>1</sup>H-NMR (400 MHz, DMSO-*d*<sub>6</sub>) δ: 5.47 (1H, d, *J* = 7.3 Hz, H-1''), 6.21 (1H, *J* = 2.0 Hz, H-6), 6.44 (1H, d, *J* = 2.0 Hz, H-8), 6.88 (2H, d, *J* = 8.0 Hz, H-3', 5'), 8.05 (2H, d, *J* = 8.0 Hz, H-2', 6'), 12.62 (1H, brs, 5-OH); <sup>13</sup>C-NMR (100 MHz, DMSO-*d*<sub>6</sub>) δ: 177.6 (C-4), 164.2 (C-7), 161.3 (C-5), 160.0 (C-4'), 156.5 (C-9), 156.4 (C-2), 133.3 (C-3), 131.0 (C-2', 6'), 121.0 (C-1'), 115.2 (C-3', 5'), 104.1 (C-10), 100.9 (C-1''), 98.8 (C-6), 93.7 (C-8), 77.6 (C-5''), 76.5 (C-3''), 74.3 (C-2''), 70.0 (C-4''), 60.9 (C-6'').

**Kaempferol 3-O- $\beta$ -D-galactopyranoside (4)**

$^1\text{H-NMR}$  (400 MHz,  $\text{DMSO-}d_6$ )  $\delta$ : 5.41 (1H, d,  $J = 7.7$  Hz, H-1''), 6.21 (1H,  $J = 2.0$  Hz, H-6), 6.43 (1H, d,  $J = 2.0$  Hz, H-8), 6.86 (2H, d,  $J = 8.0$  Hz, H-3', 5'), 8.08 (2H, d,  $J = 8.0$  Hz, H-2', 6'), 12.63 (1H, brs, 5-OH);  $^{13}\text{C-NMR}$  (100 MHz,  $\text{DMSO-}d_6$ )  $\delta$ : 177.6 (C-4), 164.2 (C-7), 161.2 (C-5), 160.0 (C-4'), 156.4 (C-9), 156.4 (C-2), 133.2 (C-3), 131.0 (C-2', 6'), 120.9 (C-1'), 115.1 (C-3', 5'), 104.0 (C-10), 101.7 (C-1''), 98.7 (C-6), 93.7 (C-8), 75.8 (C-5''), 73.1 (C-3''), 71.2 (C-2''), 67.9 (C-4''), 60.2 (C-6'').

**Myricetin 3',5'-dimethylether 3-O- $\beta$ -D-glucopyranoside (5)**

$^1\text{H-NMR}$  (400 MHz,  $\text{DMSO-}d_6$ )  $\delta$ : 3.84 (6H, s,  $\text{OCH}_3$ ), 5.59 (1H, d,  $J = 7.3$  Hz, H-1''), 6.22 (1H,  $J = 2.1$  Hz, H-6), 6.50 (1H, d,  $J = 2.1$  Hz, H-8), 7.49 (2H, s, H-2', 6'), 9.17 (1H, brs, OH), 10.87 (1H, brs, OH), 12.60 (1H, brs, 5-OH);  $^{13}\text{C-NMR}$  (100 MHz,  $\text{DMSO-}d_6$ )  $\delta$ : 177.4 (C-4), 164.1 (C-7), 161.2 (C-5), 156.3 (C-9), 156.3 (C-2), 147.4 (C-3', 5'), 138.6 (C-4'), 133.1 (C-3), 119.8 (C-1'), 106.9 (C-2', 6'), 104.0 (C-10), 100.7 (C-1''), 98.7 (C-6), 93.8 (C-8), 77.4 (C-5''), 76.4 (C-3''), 74.3 (C-2''), 69.8 (C-4''), 60.5 (C-6''), 56.2 ( $\text{CCH}_3$ ).

**Kaempferol 3-O- $\alpha$ -L-rhamnopyranosyl-(1 $\rightarrow$ 6)- $\beta$ -D-glucopyranoside (6)**

$^1\text{H-NMR}$  (400 MHz,  $\text{DMSO-}d_6$ )  $\delta$ : 0.97 (3H, s,  $J = 6.2$  Hz,  $\text{CH}_3$ ), 4.38 (1H, s, H-1'''), 5.30 (1H, d,  $J = 7.4$  Hz, H-1''), 6.23 (1H,  $J = 2.1$  Hz, H-6), 6.45 (1H, d,  $J = 2.1$  Hz, H-8), 6.89 (2H, d,  $J = 8.8$  Hz, H-3', 5'), 7.97 (2H, d,  $J = 8.8$  Hz, H-2', 6'), 10.22 (1H, brs, OH), 11.01 (1H, brs, OH), 12.55 (1H, brs, 5-OH);  $^{13}\text{C-NMR}$  (100 MHz,  $\text{DMSO-}d_6$ )  $\delta$ : 177.4 (C-4), 164.2 (C-7), 161.1 (C-5), 160.0 (C-4'), 156.8 (C-9), 156.4 (C-2), 133.2 (C-3), 130.8 (C-2', 6'), 120.8 (C-1'), 115.1 (C-3', 5'), 103.9 (C-10), 101.3 (C-1''), 100.7 (C-1'''), 98.7 (C-6), 93.7 (C-8), 75.1 (C-5''), 76.4 (C-3''), 74.1 (C-2''), 71.8 (C-4''), 70.6 (C-2'''), 70.3 (C-3'''), 69.9 (C-4'''), 68.2 (C-5'''), 66.9 (C-6''), 17.7 (C-3''').

**Kaempferol 3-O- $\beta$ -D-glucuronopyranoside (7)**

$^1\text{H-NMR}$  (400 MHz,  $\text{DMSO-}d_6$ )  $\delta$ : 5.43 (1H, d,  $J = 7.2$  Hz, H-1''), 6.15 (1H, d,  $J = 2.0$  Hz, H-6), 6.36 (1H, d,  $J = 2.0$  Hz, H-8), 6.86 (2H, d,  $J = 8.0$  Hz, H-3', 5'), 8.02 (2H, d,  $J = 8.0$  Hz, H-2', 6'), 12.48 (1H, brs, 5-OH);  $^{13}\text{C-NMR}$  (100 MHz,  $\text{DMSO-}d_6$ )  $\delta$ : 177.4 (C-4), 171.4 (C-6''), 164.5 (C-7), 161.1 (C-5), 160.1 (C-4'), 156.4 (C-9), 156.4 (C-2), 133.2 (C-3), 131.0 (C-2', 6'), 120.7 (C-1'), 115.1 (C-3', 5'), 103.8 (C-10), 101.3 (C-1''), 98.8 (C-6), 93.8 (C-8), 76.1 (C-5''), 75.2 (C-3''), 73.9 (C-2''), 71.8 (C-4'').

**Determination of the scavenging effect on DPPH radicals:**

The DPPH radical scavenging effect was evaluated

according to the method first employed by Blois (1958). A hundred and sixty microliters ( $\mu\text{L}$ ) of a MeOH solution of various sample concentrations (10-320  $\mu\text{L/mL}$ ) was added to 40  $\mu\text{L}$  of a DPPH methanol solution ( $1.5 \times 10^{-4}$  M). After mixing gently, and leaving for 30 min at room temperature, the optical density was measured at 520 nm using a microplate reader spectrophotometer VERSAmax (Molecular Devices, CA, USA). The antioxidant activity of each sample was expressed in terms of  $\text{IC}_{50}$  ( $\mu\text{g/mL}$  or  $\mu\text{M}$  required to inhibit DPPH radical formation by 50%) and calculated from the log-dose inhibition curve.

**Measurement of the inhibition of the total ROS generation**

Rat kidney homogenates, prepared from the kidneys of freshly killed male Wistar rats, weighing 150-200 g, were mixed with or without a suspension of extracts/or compounds, then incubated with 12.5  $\mu\text{M}$  DCFH-DA, at 37°C for 30 min. Fifty mM of pH 7.4 phosphate buffer was used. DCFH-DA is a stable compound, which easily diffuses into cells, and is hydrolyzed by intracellular esterase to yield a reduced non-fluorescent compound, DCFH, which is trapped within cells. The ROS produced by cells oxidize the DCFH to the highly fluorescent 2',7'-dichlorodihydrofluorescein (DCF). The fluorescence intensity of the oxidized DCF was monitored on a microplate fluorescence spectrophotometer (Bio-Tek Instruments Inc., Winooski, VT), with excitation and emission wavelengths of 460 and 530 nm, respectively (Label and Bondy, 1990).

**Measurement of the ONOO<sup>-</sup> scavenging activity**

The ONOO<sup>-</sup> scavenging activity was measured by monitoring the oxidation of DHR 123, by modifying the method of Kooy *et al.* (1994). The DHR 123 (5 mM), in dimethylformamide, was purged with nitrogen, stored at -80°C and used as a stock solution. This solution was then placed on ice, and kept from exposure to light, prior to the study. The buffer used consisted of 90 mM sodium chloride, 50 mM sodium phosphate, 5 mM potassium chloride, at pH 7.4, and 100  $\mu\text{M}$  diethylenetriaminepentaacetic acid (DTPA), each of which were prepared with high quality deionized water, and purged with nitrogen. The final concentration of the DHR 123 was 5  $\mu\text{M}$ . The background and final fluorescent intensities were measured 5 min after treatment, both with and without the addition of authentic ONOO<sup>-</sup>. The DHR 123 was oxidized rapidly by authentic ONOO<sup>-</sup>, and its final fluorescent intensity remained unchanged over time. The fluorescence intensity of the oxidized DHR 123 was measured using a microplate fluorescence reader FL 500 (Bio-Tek Instruments Inc.), with excitation and emission wavelengths of 480 and 530 nm, respectively. The results were expressed as the mean  $\pm$  standard error ( $n=3$ ) of the final fluorescence

intensity minus the background fluorescence. The effects were expressed as the percentage of inhibition of the DHR 123 oxidation.

### Statistical analysis

All values were expressed as the mean±standard error of three or five replicate experiments.

## RESULTS AND DISCUSSION

The formation of reactive species, such as  $H_2O_2$ ,  $O_2^-$ ,  $\cdot OH$ ,  $NO\cdot$  and  $ONOO^-$ , in the body, as a consequence of aerobic metabolism, play roles in the pathogenesis of human degenerative diseases (Pincemail, 1995; Beckman et al., 1990), and are also likely to damage intracellular components, such as nucleic acids, proteins and lipids (Sagar et al., 1992; Ames et al., 1993).

In our ongoing investigations to identify powerful, nontoxic and natural antioxidants, we evaluated general antioxidants for their potential to scavenge authentic  $ONOO^-$  and DPPH radicals as well as to inhibit the total ROS generation of the MeOH extract, along with its solvent soluble fractions, including the  $CH_2Cl_2$ , EtOAc and *n*-BuOH fractions, as well as the  $H_2O$  layer derived from *N. nucifera* stamens. As summarized in Table I, the EtOAc, *n*-BuOH and  $H_2O$  fractions possessed good scavenging activities for the DPPH radical, with 50% inhibition concentration ( $IC_{50}$ ) values of 13.5, 34.5 and 37.5  $\mu g/mL$  respectively, compared to the 6.2  $\mu g/mL$  of the well-known antioxidant, L-ascorbic acid. In the total ROS system, the  $IC_{50}$  values of these three fractions were  $7.81\pm 0.91$ ,  $42.25\pm 0.33$  and  $64.64\pm 3.35$   $\mu g/mL$ , respectively, with the EtOAc fraction especially exhibiting more potent inhibitory activity than the positive

control Trolox ( $IC_{50}$   $28.42\pm 1.21$   $\mu g/mL$ ). The MeOH extract, and all the fractions from the stamens of *N. nucifera*, showed marked scavenging activity on the authentic  $ONOO^-$  in the following order: EtOAc fraction ( $IC_{50}$   $0.11\pm 0.06$   $\mu g/mL$ ) > *n*-BuOH fraction ( $IC_{50}$   $3.41\pm 0.26$   $\mu g/mL$ ) > MeOH extract ( $IC_{50}$   $18.90\pm 2.14$   $\mu g/mL$ ) >  $H_2O$  layer ( $IC_{50}$   $28.01\pm 1.21$   $\mu g/mL$ ) >  $CH_2Cl_2$  fraction ( $IC_{50}$   $35.52\pm 1.01$   $\mu g/mL$ ). The  $ONOO^-$  scavenging activities of the EtOAc and *n*-BuOH fractions were about eight and two time as good, respectively, as that of the positive control (penicillamine), which has an  $IC_{50}$  value of  $8.27\pm 0.06$   $\mu g/mL$ . These results suggest that there are likely to be many antioxidants in the EtOAc soluble fraction, so much attention should be given to the isolation of the antioxidative compounds from this fraction.


The EtOAc soluble part of the MeOH extract, from the stamens of *N. nucifera*, repeatedly underwent chromatography on the Si gel and Sephadex LH-20 to yield compounds 1-8. The structural identifications of these compounds were elucidated from the 1D ( $^1H$ - and  $^{13}C$ -NMR) and 2D NMR (HMQC and HMBC) spectral data, and by comparison with published spectral data (Park et al., 1991; Mpalantinos et al., 1998; Agrawal, 1992; Pauli, 2000; Ismail and Alam, 2001; Sung et al., 1997). The isolated compounds were readily elucidated as: kaempferol (1), kaempferol 3-O- $\beta$ -D-glucuronopyranosyl methylester (2), kaempferol 3-O- $\beta$ -D-glucopyranoside (3), kaempferol 3-O- $\beta$ -D-galactopyranoside (4), myricetin 3',5'-dimethylether 3-O- $\beta$ -D-glucopyranoside (5), kaempferol 3-O- $\alpha$ -L-rhamnopyranosyl-(1 $\rightarrow$ 6)- $\beta$ -D-glucopyranoside (6), kaempferol 3-O- $\beta$ -D-glucuronopyranoside (7) and  $\beta$ -sitosterol glucopyranoside (8). The chemical structures of compounds 1-8, from the stamens of *N. nucifera*, are shown in Fig. 1. Each chemical shift, corresponding to the aglycone of seven compounds 1-4, 6 and 7 were similar to those of the  $^1H$ - and  $^{13}C$ -NMR spectra of kaempferol. This indicated that the compounds 2-4, 6 and 7 were kaempferol glycosides and their structures were further confirmed from the HMQC and HMBC spectroscopies, as well as by comparison with the published NMR spectral data, of several monoglycosides (Agrawal, 1992; Pauli, 2000).

The  $^1H$ -NMR data for compound 5 suggested the presence of three hydroxyl signals (1H, brs), in the downfield region, at  $\delta_H$  9.17,  $\delta_H$  10.87,  $\delta_H$  12.60, along with two characteristic *meta*-coupled doublets ( $J = 2.1$  Hz) at  $\delta_H$  6.22 (H-6) and  $\delta_H$  6.50 (H-8). Additionally, there was a two-proton singlet at  $\delta_H$  7.49 (H-2', H-6'), a six-proton singlet at  $\delta_H$  3.84, corresponding to two methoxyl groups at the *meta*-position in the B ring, as well as one anomeric proton at  $\delta_H$  5.59 (1H,  $J = 7.3$  Hz). The above  $^1H$ -NMR data and values of the coupling constants of 5 led us to conclude it was the glycoside of myricetin dimethylether, which further supported by the  $^{13}C$ -NMR data. In the

**Table I.** Antioxidant activities of the MeOH extract and its solvent soluble fractions of the stamens of *N. nucifera*

Samples	DPPH <sup>a</sup> $IC_{50}$ ( $\mu g/mL$ )	Total ROS <sup>b</sup> $IC_{50}$ ( $\mu g/mL$ ) Mean $\pm$ SE <sup>d</sup>	$ONOO^-$ <sup>c</sup> $IC_{50}$ ( $\mu g/mL$ ) Mean $\pm$ SE <sup>d</sup>
MeOH	273.5	> 100	$18.90\pm 2.14$
$CH_2Cl_2$	279.0	> 100	$35.52\pm 1.01$
EtOAc	13.5	$7.81\pm 0.91$	$0.11\pm 0.06$
<i>n</i> -BuOH	34.5	$42.25\pm 0.33$	$3.41\pm 0.26$
$H_2O$	37.5	$64.64\pm 3.35$	$28.01\pm 1.21$
L-Ascorbic acid	6.2		
Trolox		$28.42\pm 1.21$	
Penicillamine			$8.27\pm 0.06$

<sup>a</sup>DPPH is the free radical scavenging activity ( $IC_{50}$ :  $\mu g/mL$ ). <sup>b</sup>Total ROS is the inhibitory activity of the total free radical generation in the kidney postmicrosomal fraction ( $IC_{50}$ :  $\mu g/mL$ ). <sup>c</sup> $ONOO^-$  is the inhibitory activity of authentic peroxyntirite ( $IC_{50}$ :  $\mu g/mL$ ). <sup>d</sup>Values of total ROS and  $ONOO^-$  were expressed as the mean±standard error of three or five experiments.



Kaempferol (1):  $R_1=H, R_2=H, R_3=H$

Kaempferol 3-O- $\beta$ -D-glucuronopyranosyl methyl ester (2):  $R_1=Gln-Me, R_2=R_3=H$

Kaempferol 3-O- $\beta$ -D-glucopyranoside (3):  $R_1=Glu, R_2=R_3=H$


Kaempferol 3-O- $\beta$ -D-galactopyranoside (4):  $R_1=Gal, R_2=R_3=H$

Myricetin 3',5'-dimethylether 3-O- $\beta$ -D-glucopyranoside (5):  $R_1=Glu, R_2=R_3=OCH_3$

Kaempferol 3-O- $\alpha$ -L-rhamnopyranosyl-(1 $\rightarrow$ 6)- $\beta$ -D-glucopyranoside (6):

$R_1=Flt\alpha-(1\rightarrow6)-Glu, R_2=R_3=H$

Kaempferol 3-O- $\beta$ -D-glucuronopyranoside (7):  $R_1=Gln, R_2=R_3=H$



$\beta$ -sitosterol glucopyranoside (8)

Fig. 1. Chemical structures of compounds 1-8 from the stamens of *N. nucifera*.

HMBC, the proton signal for two *meta*-methoxyl groups, at  $\delta_H$  3.84 (6H, s), was correlated with that of the myricetin C-2' and 6', at  $\delta_C$  115.4. Also, an anomeric glucose proton signal, at  $\delta_H$  5.59 (1H,  $J = 7.3$  Hz), showed a cross peak with the myricetin C-3, at  $\delta_C$  133.1. These results confirmed that 5 was myricetin 3',5'-dimethylether 3-O- $\beta$ -D-glucopyranoside.

The  $^1H$ -NMR data for compound 6 showed signals for two *meta*-coupled protons at  $\delta_H$  6.23 (1H,  $J = 2.1$  Hz, H-6) and  $\delta_H$  6.45 (1H, d,  $J = 2.1$  Hz, H-8), together with two *ortho*-coupled systems at  $\delta_H$  6.89 (2H, d,  $J = 8.8$  Hz, H-3', 5') and  $\delta_H$  7.97 (2H, d,  $J = 8.8$  Hz, H-2', 6'), indicating a kaempferol derivative (Harborne, 1980; Agrawal, 1992). From the analysis of the  $^1H$ -NMR and coupling constants of 6, there were characteristic signals of rhamnopyranoside and glucopyranoside, at  $\delta_H$  4.38 (1H, s, H-1'') and at  $\delta_H$  5.30 (1H, d,  $J = 7.4$  Hz, H-1''), respectively (Pauli, 2000). These were further supported by the HMBC spectrum; the signals at  $\delta_H$  4.38 and  $\delta_H$  5.30 correlated with those of glucose C-6, at  $\delta_C$  66.9, and kaempferol C-3, at  $\delta_C$  133.2, respectively. These results suggested that the order of sugars was  $\alpha$ -L-rhamnopyranosyl-(1 $\rightarrow$ 6)- $\beta$ -D-glucopyranoside (Agrawal, 1992), which confirmed 6 to be kaempferol 3-O- $\alpha$ -L-rhamnopyranosyl-(1 $\rightarrow$ 6)- $\beta$ -D-glucopyranoside.

Table II shows the antioxidative activities of the flavonoids 1-4 and 7, as kaempferol, and its derivatives isolated from the active EtOAc fraction of *N. nucifera* stamens, on the DPPH radical, total ROS and ONOO<sup>-</sup> systems. Flavonoids,

Table II. Antioxidant activities of isolated compounds 1-4 and 7 from the stamens of *N. nucifera*

Compounds	DPPH <sup>a</sup> IC <sub>50</sub> ( $\mu$ M)	Total ROS <sup>b</sup> IC <sub>50</sub> ( $\mu$ M) Mean $\pm$ SE <sup>d</sup>	ONOO <sup>-c</sup> IC <sub>50</sub> ( $\mu$ M) Mean $\pm$ SE <sup>d</sup>
1	13.8	26.84 $\pm$ 2.05	3.00 $\pm$ 0.07
2	61.4	> 100	10.51 $\pm$ 0.77
3	> 100	> 100	6.98 $\pm$ 0.37
4	> 100	> 100	16.59 $\pm$ 3.99
7	20.1	> 100	3.18 $\pm$ 0.39
L-Ascorbic acid	11.5		
Trolox		6.09 $\pm$ 1.29	
Penicillamine			5.11 $\pm$ 0.23

<sup>a</sup>DPPH is the free radical scavenging activity (IC<sub>50</sub>:  $\mu$ M). <sup>b</sup>Total ROS is the inhibitory activity of the total free radical generation in the kidney postmicrosomal fraction (IC<sub>50</sub>:  $\mu$ M). <sup>c</sup>ONOO<sup>-</sup> is the inhibitory activity of authentic peroxynitrite (IC<sub>50</sub>:  $\mu$ M). <sup>d</sup>Values of total ROS and ONOO<sup>-</sup> were expressed as the mean $\pm$ standard error of three or five experiments.

a family of diphenylpropanes (C<sub>6</sub>-C<sub>3</sub>-C<sub>6</sub>), which are naturally occurring compounds that are widely distributed in the plant kingdom. In particular, flavonoids have been found to owe their antioxidant activity to both their excellent metal chelating and radical scavenging properties (Morel *et al.*, 1993; Salah *et al.*, 1995; Van Acker *et al.*, 1998; Rice-Evans *et al.*, 1996). The differences in flavonoids are dependent on their basic structure, as well as the number, and arrangement of the functional groups, such as hydroxyl groups, methoxyl groups and glycosidic units (Rice-Evans *et al.*, 1997). Of the five flavonoids tested, compound 1 possessed good activities in the three model systems tested. Compounds 2 and 7 showed scavenging activities in the DPPH and ONOO<sup>-</sup> tests, while compounds 3 and 4 were only active in the ONOO<sup>-</sup> test. Because compounds 5 and 6 were isolated as minor components, their antioxidant activities were not determined. Conversely, compound 8 exhibited no activity in any of the model systems tested, although the data is not shown. These results were in accordance with the other research on the structure-antioxidant activity relationship of flavonoids. As shown in Table II, the differences in the structures of flavonoids affected their antioxidant activity in the three model systems tested. When the hydroxyl groups at the C-3 positions are masked by a glycoside, and other factors, the IC<sub>50</sub> values were dramatically decreased, although differences according to the diversity of the systems used were shown. The above results suggest that the hydroxyl groups at the C-3 positions are a structural requirement for the antioxidant activities of the flavonoids in the model systems tested (Van Acker, S.A.B.E. *et al.*, 1996; Cos *et al.*, 1998; Nagao *et al.*, 1999; Haenen *et al.*, 1997; 2001;

Choi *et al.*, 2002).

The present work found that the antioxidant capacities of the MeOH extract, its various fractions and active components, from the *N. nucifera* stamens, could be useful for the treatment of oxidative damage. Furthermore, it will be interesting to investigate the antioxidative activity of these natural compounds still further for the prevention of various radical-mediated injuries in pathological situations *in vivo*. The investigation of further antioxidant principles, and the biochemical mechanisms of the flavonoids remain to be elucidated in future studies.

## ACKNOWLEDGEMENT

This research was supported by a grant (PE002201-07 and -08) from the Plant Diversity Research Center of 21<sup>st</sup> Century Frontier Research Program, funded by the Ministry of Science and Technology of the Korean government.

## REFERENCES

- Agrawal, P. K., NMR spectroscopy in the structural elucidation of oligosaccharides and glycosides. *Phytochem.*, 31, 3307-3330 (1992).
- Ames, B. N., Shigenaga, M. K., and Hage, T. M., Oxidants, antioxidants and the degenerative diseases of aging. *Proc. Natl. Acad. Sci. USA*, 90, 7915-7922 (1993).
- Beckman, J. S., Beckman, T. W., Chen, J., Marshall, P. A., and Freeman, B. A., Apparent hydroxyl radical production by peroxynitrite: implications for endothelial injury from nitric oxide and superoxide. *Proc. Natl. Acad. Sci., USA*, 87, 1620-1624 (1990).
- Blois, M. S., Antioxidant determination by the use of a stable free radical. *Nature*, 181, 1199-1202 (1958).
- Choi, J. S., Chung, H. Y., Kang, S. S., Jung, M. J., Kim, J. W., No, J. K., and Jung, H. A., The structure-activity relationship of flavonoids as scavengers of peroxynitrite. *Phytother. Res.*, 16, 232-235 (2002).
- Cos, P., Ying, L., Calomme, M., Hu, J. P., Cimanga, K., Poel, B. V., Pieters, L., Vlietinck, A. J., and Berghe, D., Structure-activity relationship and classification of flavonoids as inhibitors of xanthine oxidase and superoxide scavengers. *J. Nat. Prod.*, 61, 71-76 (1998).
- Dreher, D. and Junod, F., Role of oxygen free radicals in cancer development. *Eur. J. Cancer*, 32A(1), 30-38 (1996).
- Griffiths, H. R. and Lunec, J., The C1q binding activity of IgG is modified *in vitro* by reactive oxygen species: implications for rheumatoid arthritis. *FEBS Lett.*, 388, 161-164 (1996).
- Guo, C., Wei, J., Ge, Y., and Yang, J., ROS scavenging activity of nuts. *Shipin Kexue* (Beijing) 21, 42-44 (2000).
- Haenen, G. R. M. M., Parquay, J. B. G., Korthouwer, R. E. M. and Bast, A., Peroxynitrite scavenging by flavonoids. *Biochem. Biophys. Res. Commun.*, 236, 591-593 (1997).
- Harborne, J. B., *The flavonoids-Advances in Research since 1980*. Chapman and Hall, London, (1988).
- Heijnen, C. G. M., Haenen, G. R. M. M., Vekemans, J. A. J. M. and Bast, A., Peroxynitrite scavenging of flavonoids: structure activity relationship. *Environ. Toxicol. Pharmacol.*, 10, 199-206 (2001).
- Ismail, N. and Alam, M., A novel cytotoxic flavonoid glycoside from *Physalis angulata*. *Fitoterapia*. 72, 676-679 (2001).
- Kim, J. G., *Illustrated Natural Drugs Encyclopedia* (color edition). Namsandang, Korea, Vol. 2, pp. 95, (1997).
- Kim, T. J., *Korean Resources Plants*. Seoul National University Publisher, Seoul, Korea, Vol. II, pp. 218-219, (1996).
- Kooy, N. W., Royall, J. A., Ischiropoulos, H., and Beckman, J. S., Peroxynitrite-mediated oxidation of dihydrorhodamine 123. *Free Radic. Biol. Med.*, 16, 149-156 (1994).
- La Cour, B., Mølgaard, P. and Yi, Z., Traditional Chinese medicine in treatment of hyperlipidaemia. *J. Ethnopharmacol.*, 46, 125-129 (1995).
- Label, C. P. and Bondy, S. C., Sensitive and rapid quantitation of oxygen reactive species formation in rat synaptosomes. *Neurochem. Int.*, 17, 435-441 (1990).
- Larson, R. A., The antioxidants of higher plants. *Phytochem.*, 27, 969-978 (1988).
- Liou, B.-K., Chen, H.-Y., and Yen, G.-C., Antioxidant activity of the methanolic extracts from various traditionally edible plants. *Zhonggou Nongye Huaxue Huizhi*, 37, 105-116 (1999).
- Mpalantinos, M. A., De Moura, R. S., Parente, J. P., and Kuster, R. M., Biologically active flavonoids and kava pyrones from the aqueous extract of *Alpinia zerumbet*. *Phytother. Res.*, 12, 442-444 (1998).
- Morel, I., Lescoat, G., Cognel, P., Sergent, O., Padelop, N., Brissot, P., Cillard, P., and Cillard, J., Antioxidant and iron-chelating activities of the flavonoids catechins, quercetin and diosmetin on iron-loaded rat hepatocyte cultures. *Biochem. Pharmacol.*, 45, 13-19 (1993).
- Mukherjee, P. K., Das, J., Balasubramanian, R., Saha, K., Pal, M., and Saha, B. P., Antidiarrhoeal evaluation of *Nelumbo nucifera* rhizome extract. *Indian J. Ethnopharmacol.*, 27, 262-264 (1995).
- Mukherjee, P. K., Saha, K., Balasubramanian, R., Pal, M., and Saha, B. P., Studies on psychopharmacological effects of *Nelumbo nucifera* Gaertn. rhizome extract. *J. Ethnopharmacol.*, 54, 63-67 (1996).
- Mukherjee, P. K., Saha, K., Pal, M., and Saha, B. P., Effect of *Nelumbo nucifera* rhizome extract on blood sugar level in rats. *J. Ethnopharmacol.*, 58, 207-213 (1997).
- Nagao, A., Seki, M., and Kobayashi, H., Inhibition of xanthine oxidase by flavonoids. *Biosci. Biotechnol. Biochem.*, 63, 1787-1790 (1999).
- Park, H. J., Young, H. S., Park, K. Y., Rhee, S. H., Chung, H. Y., and Choi, J. S., Flavonoids from the whole plants of *Orostachys japonicus*. *Arch. Pharm. Res.*, 14, 167-171 (1991).
- Pauli, G. F., Higher order and substituent chemical shift effects

- in the proton NMR of glycosides. *J. Nat. Prod.*, 63, 834-838 (2000).
- Pincus, J. J., Free radicals and antioxidants in human diseases, In Favier, A. E., Cadet, J., Kalyanaraman, B., Fontecave, M., Pierre, J.-L. (Eds.). *Analysis of Free radicals in Biological Systems*. Birkhauser Verlag, Berlin, pp.83-98, (1995).
- Pratt, D. E., Natural anti-oxidants from plant material, In *Phenolic compounds and their effects on health II*, ACS symposium series 547, American Chemical Society, Washington DC, pp. 54-71, (1994).
- Rice-Evans, C. A., Miller, N., and Paganga, G., Antioxidant properties of phenolic compounds. *Trends in Plant Science*, 2, 152-159 (1997).
- Rice-Evans, C. A., Miller, N., and Paganga, G., Structure-antioxidant activity relationships of flavonoids and phenolic acids. *Free Radic. Biol. Med.*, 20, 933-956 (1996).
- Saga, S., Kalló, I. J., Kaul, N., Ganguly, N. K., and Sharma, B. K., Oxygen free radicals in essential hypertension. *Mol. Cell Biochem.*, 111, 103-108 (1992).
- Salah, N., Miller, N. J., Paganga, G., Tijburg, L., Bolwell, G. P., and Rice-Evans, C. A., Polyphenolic flavanols as scavengers of aqueous phase radicals and as chain-breaking antioxidants. *Arch. Biochem. Biophys.*, 322, 339-346 (1995).
- Sinha, S., Mukherjee, P. K., Mukherjee, K., Pal, M., Mandal, S. C., and Saha, K., Evaluation of antipyretic potential of *Nelumbo nucifera* stalk extract. *Phytother. Res.*, 14, 272-274 (2000).
- Sohal, R. S., Role of oxidative stress and protein oxidation in the aging process. *Free Radic. Biol. Med.*, 33(1), 37-44 (2002).
- Squadrito, G. L. and Pryor, W. A., Oxidative chemistry of nitric oxide: the role of superoxide, peroxy nitrite, and carbon dioxide. *Free Radic. Biol. Med.*, 25, 392-403 (1998).
- Sung, S. H., Kwon, S. H., Cho, N. J., and Kim, Y. C., Hepatoprotective flavonol glycosides of *Saururus chinensis* herbs. *Phytother. Res.*, 11, 500-503 (1997).
- Van Acker, S. A. B. E., Van Balen, G. P., Van den Berg, D. J., Bast, A., and Van der Vijgh, W. J. F., Influence of iron chelation on the antioxidant activity of flavonoids. *Biochem. Pharmacol.*, 56, 935-943 (1998).
- Van Bergen, P. E., Hatcher, P. G., Boon, J. J., Collinson, M. E., and De Leeuw, J. W., Macromolecular composition of the propagule wall of *Nelumbo nucifera*. *Phytochem.*, 45, 601-617 (1997).