

Cloning and Sequencing Analysis of the Repressor Gene of Temperate Mycobacteriophage L1

Subrata Sau*, Partho Chattoraj, Tridib Ganguly, Chia Yen Lee[†] and Nitai Chandra Mandal*

Department of Biochemistry, Bose Institute, P1/12-CIT Scheme VII M, Calcutta 700 054, India

[†]Department of Microbiology, Molecular Genetics and Immunology, KU Medical Center, 3901 Rainbow Blvd., Kansas City, KS 66103, USA

Received 24 June 2003, Accepted 13 August 2003

The wild-type and *temperature-sensitive (ts)* repressor genes were cloned from the temperate mycobacteriophage L1 and its mutant L1cIts391, respectively. A sequencing analysis revealed that the 131st proline residue of the wild-type repressor was changed to leucine in the *ts* mutant repressor. The 100% identity that was discovered between the two DNA regions of phages L1 and L5, carrying the same sets of genes including their repressor genes, strengthened the speculation that L1 is a minor variant of phage L5 or vice versa. A comparative analysis of the repressor proteins of different mycobacteriophages suggests that the mycobacteriophage-specific repressor proteins constitute a new family of repressors, which were possibly evolved from a common ancestor. Alignment of the mycobacteriophage-specific repressor proteins showed at least 7 blocks (designated I-VII) that carried 3-8 identical amino acid residues. The amino acid residues of blocks V, VI, and some residues downstream to block VI are crucial for the function of the L1 (or L5) repressor. Blocks I and II possibly form the turn and helix 2 regions of the HTH motif of the repressor. Block IV in the L1 repressor is part of the most charged region encompassing amino acid residues 72-92, which flanks the putative N-terminal basic (residues 1-71) and C-terminal acidic (residues 93-183) domains of L1 repressor.

Keywords: Helix-turn-helix (HTH) motif, *M. smegmatis*, Mycobacteriophage L1, Repressor gene, Temperature-sensitive (*ts*)

Introduction

The phages of mycobacteria are extremely diverse in nature and carry highly mosaic genomes (Pedulla *et al.*, 2003). Several molecular tools have been developed from mycobacteriophages during the last ten years. They are very useful for mycobacterial research and the diagnosis of mycobacterial infections (Hatfull, 2000). Among the mycobacteriophages, L5, Bxb1, I3 (Hatfull, 2000), Ms6 (Garcia *et al.*, 2002), and L1 (Chaudhuri *et al.*, 1993) were studied to some extent at the molecular level. Both the *cis*- and *trans*-acting regulatory elements that are involved in the integration of L5 into its host genome were identified and characterized at length (Hatfull, 2000). Several promoters of I3 (Ramesh and Gopinathan, 1995), L5 (Nesbit *et al.*, 1995), Bxb1 (Jain and Hatfull, 2000), and Ms6 (Garcia *et al.*, 2002) were reported. The repressors of both L5 and Bxb1 negatively regulate the expression of their respective early promoters by binding at the cognate operators (Nesbit *et al.*, 1995; Jain and Hatfull, 2000). Currently, the molecular mechanism of the interaction between the repressor of any mycobacteriophage and its operator DNA is poorly understood, though it has immense potential in deciphering the gene regulation in both mycobacteriophage and mycobacterial systems. Also, the information could lead to the construction of the tightly regulated expression vector (for the mycobacterial system) by assembling the early promoter and the repressor gene of temperate mycobacteriophage.

Mycobacteriophage L1, a sister homoimmune phage of L5, has a 50-kb double-stranded DNA genome. The genes that regulate both the lysogenic and lytic development of L1 were mapped and some were to some extent characterized (Chaudhuri *et al.*, 1993). The *G27* gene of L1 was shown to be an early positive regulator as it controls the expression of both the delayed early and late genes at the transcriptional level (Datta and Mandal, 1998). A few promoters of L1 have been cloned in a promoter-cloning vector having β -galactosidase as

*To whom correspondence should be addressed.

Fax: 91-33-2337-9416

E-mail: sau@bic.boseinst.ernet.in or ncmandal@boseinst.ernet.in

the reporter gene, but further characterization of these promoters is yet to be reported (Barletta *et al.*, 1992). Recently, two early promoters of L1 were cloned, which are regulated by its own repressor (Chattopadhyay *et al.*, 2003). In this communication, as a prerequisite to developing suitable mycobacteria-specific expression vectors and studying the molecular nature of the DNA-protein interaction in the mycobacterial system, we report the cloning and preliminary characterization of the repressor genes of both wild-type L1 and its *cIts391* mutant derivative (Chaudhuri *et al.*, 1993).

Materials and Methods

Bacterial and phage strains, vectors and growth conditions *E. coli* DH5 α cells were grown in Luria broth. *M. smegmatis* mc²155 and LR222, obtained from Dr. Anil Tyagi (University of Delhi, South Campus, New Delhi, India), were routinely grown in Middlebrook 7H9 medium (Chaudhuri *et al.*, 1993). The wild-type mycobacteriophage L1, its mutants L1cI, L1cIts391, and its lysogen were described in Chaudhuri *et al.* (1993). The *E. coli*-*M. smegmatis* shuttle vector pSD5S30, containing a mycobacteria-specific promoter *S30*, was obtained from Dr. Anil Tyagi (Bashyam *et al.*, 1996; Jain *et al.*, 1997).

DNA isolations and manipulations All of the molecular biological and recombinant DNA techniques were used as described by Sambrook *et al.* (1989). Mycobacteriophage L1 DNA was isolated as described by Chaudhuri *et al.* (1993). Isolation of plasmid DNA from *M. smegmatis* and the electrotransformation of *M. smegmatis* LR222 were performed according to the standard methods of Das Gupta *et al.* (1993).

Cloning of the wild-type repressor gene of mycobacteriophage L1 L1 DNA was digested with *Pst*I and ligated to the identical site of pSD5S30. The ligated DNA was electroporated into competent *M. smegmatis* LR222 (L1cIts391), then the culture was plated on Luria agar containing kanamycin (25 μ g/ml). After incubation for 20 h at 32°C, when the transformants were just visible, the plates were shifted to 42°C and further incubated for 20–24 h. The colonies, which grew larger at 42°C, were selected and purified. The ones that showed immunity to superinfection by L1cI were selected and purified. Plasmids were isolated from three of these transformants and digested with *Pst*I. The fragments were analyzed by 1% agarose gel electrophoresis.

Construction of a limited genomic library of L1cIts391 The genomic DNA of L1cIts391 was digested together with *Bam*HI & *Sal*I. The DNA fragments with a size range from 1,000 to 2,000 bp were eluted from low melting gel. Fragments were ligated into identical sites of pBluescript SK. DNAs were then transformed to *E. coli* XL1 Blue. Several transformants, which appeared as colorless colonies on LA-Amp-X-gal-IPTG, were selected and purified. Plasmids, isolated from twenty such colorless colonies, were digested by *Bam*HI & *Sal*I and analyzed by agarose gel electrophoresis (data not shown). Those carrying the 1.4 kb *Bam*HI-*Sal*I fragments were selected and purified.

DNA sequencing and analysis DNA sequencing of the L1 inserts were carried out by ABI automated sequencer according to the manufacturer's protocol.

DNA accession number Upon deposition of the 1.4 kb L1 DNA sequence (of pCP2) carrying the *ts* mutant repressor gene of phage L1, the accession number AY303696 was obtained from GenBank.

Results and Discussion

Cloning of the DNA fragment carrying the wild-type repressor gene of phage L1 In temperate mycobacteriophage L1, the gene encoding repressor has been designated *cI* (Chaudhuri *et al.*, 1993). The *cIts391* mutation, which makes the repressor phenotypically temperature sensitive, causes the induction of L1cIts391 lysogen in *M. smegmatis* at 42°C, thereby killing the bacterium (Chaudhuri *et al.*, 1993). It was assumed that the wild-type repressor of L1 that was supplied in *trans* from a plasmid in a lysogen of L1cIts391 could prevent the induction (killing) of the latter at 42°C. Therefore, for cloning the repressor gene of phage L1, a *Pst*I genomic library of L1 DNA was prepared by ligating *Pst*I-digested L1cI DNA fragments with the same enzyme-digested plasmid DNA and transformed into *M. smegmatis* LR222 lysogenic for L1cIts391. The resulting transformants, which survived and grew on a kanamycin plate at 42°C, were selected and purified. A restriction analysis of the plasmids from three of these transformants showed that all of them carried a common 6 kb *Pst*I fragment of L1 DNA (data not shown). This plasmid was designated pLC1 (Table 1). The L1cI phage did not plate on the LR222 cells carrying the pLC1 plasmid. This suggests that the 6 kb insert in the pLC1 plasmid possibly carries the wild-type *cI* gene of L1.

Deletion analysis of 6 kb *Pst*I fragment To identify the shorter segment of DNA within the previously mentioned 6 kb *Pst*I fragment carrying the putative wild-type repressor gene of L1, the gel purified 6 kb *Pst*I fragment was digested with *Bam*HI. The resulting DNA fragments were ligated with *Bam*HI-digested and *Bam*HI & *Pst*I double-digested pSD5S30 DNA. The ligated DNAs were separately transformed into LR222 (L1cIts391) lysogen. The transformants, which formed colonies, both at 32° and 42°C, were selected. A couple of the transformants that showed heat-stable properties were obtained from the set where *Bam*HI and *Pst*I double-digested pSD5S30 DNA was used. A restriction analysis of the plasmids that were isolated from 3 such transformants showed that they all carried a 2.1 kb *Bam*HI-*Pst*I fragment (data not shown). This plasmid was designated pLC2 (Table 1). LR222 (pLC2) also showed immunity to superinfection by L1cI, which suggests that the 2.1 kb insert of pLC2 carries the wild-type *cI* repressor gene of L1 (Table 1). Further deletion of the 2.1 kb L1 DNA insert by *Sau*3AI partial digestion resulted in the loss of immunity to

Table 1. Plating of L1cI⁻ phages on *M. smegmatis* carrying different plasmids and phages

<i>M. smegmatis</i> strains	Size of insert (kb)	Percent e.o.p. ^a of L1cI ⁻	Diameter of Plaques (mm)	
			32°C	42°C
LR222	-	100	2.12	3.10
LR222 (L1cI ⁻)	-	<10 ⁻⁷	-	-
LR222 (pLC1)	6	<10 ⁻⁷	-	-
LR222 (pLC2)	2.1	<10 ⁻⁷	-	-
mc ² 155 (pSD5S30)	-	100	2.00	3.00
mc ² 155 (pCP2)	1.4	36	0.40	1.00

^aPercent efficiency of plating (e.o.p) on all bacteria were determined relative to *M. smegmatis* at 32°C. Abbreviations: +, yes; -, no. See text for details.




Fig. 1. Alignment of the amino acid sequences of mycobacteriophage-specific repressors. The amino acid sequences of the repressor proteins of Bxb1 (ac. no. AAG59774), L1, MS6 (ac. no. AAG48322), and Bxz2 (ac. no. AAN01828) are aligned by the Clustal W program. Blocks carrying 3-8 identical amino acids are shaded and numbered I to VII. The 131st amino acid proline, which is changed to leucine in the repressor gene of L1cI^s391, has been made bold. See text for details.

superinfection by L1cI⁻ (data not shown).

Identification of the repressor gene of L1 Since the previously mentioned 2.1 kb L1 DNA fragment carries the repressor gene of L1, it was sequenced to know more about this negative regulatory gene and other neighboring genes. A sequencing analysis revealed five intact open-reading frames, designated ORF1 to ORF5, in the 2.1 kb L1 DNA (data not shown). All of the ORFs were transcribed in one direction. To detect which of these ORFs encodes the repressor protein CI of L1, databases were searched against the amino acid sequence of each of the ORFs. Surprisingly, ORF5 was found to be 100% identical with that of the repressor gene 71 of the

homoimmune phage L5 (Hatfull, 2000). Therefore, ORF5 in the 2.1 kb L1 DNA was designated the cI repressor gene of L1. The ORF1, ORF2, ORF3, and ORF4 were also found to be perfectly identical to gp75, gp74, gp73, and gp72 of L5, respectively. The entire 2.1 kb L1 DNA sequence is in fact identical to the 43,933-46,039 bp coordinates of the L5 genome (Hatfull, 2000). It was reported earlier that both L1 and L5 have an identical digestion pattern for over 20 restriction enzymes (Lee *et al.*, 1991; our unpublished data), and the former does not yield plaque at 42°C (Lee *et al.*, 1991). A variant of L1 was isolated and grew well at 42°C (Chaudhuri *et al.*, 1993). Recently, a 430 bp DNA fragment carrying the early promoters of L1 was also shown to be

100% identical to the DNA region corresponding to the 51,694-52,123 bp coordinates of L5 (Chattopadhyay *et al.*, 2003). All of the data has in fact strengthened the speculation that L1 is a minor variant of L5 or vice versa.

Other proteins, which show about 40-50% identities over almost the entire length of L1 repressor are the gp6, gp74, and gp69 of mycobacteriophage MS6, Bxz2, and Bxb1, respectively (Fig. 1). The gp74 is the smallest among these orthologous proteins. The most notable finding is that there are about 31 amino acid residues at the N-terminal end in gp74 that are missing (Fig. 1). Interestingly, the gp6 of Ms6 was more identical (about 96%) to the repressor of Bxb1 than to either L1 or Bxz2 (Fig. 1). On the other hand, gp74 showed about a 58% identity to both gp6 and gp69. Apart from these orthologs of the CI protein of L1, a couple of DNA binding proteins also showed a very good degree of similarity within a narrow region (carrying amino acid residues 34-53) of the L1 repressor (discussed below). These results, therefore, suggest that CI of L1, gp6, gp69, and gp74 constitute a new family of repressor proteins of mycobacteriophage origin that possibly evolved from a common ancestor.

Identification of the temperature-sensitive mutation in the repressor gene of L1cIts391 Several *ts* mutant repressor proteins of phage λ were used extensively to elucidate the structure and function of its repressor (Chattopadhyaya and Ghosh, 2003) and to construct a couple of expression vectors (Sambrook *et al.*, 1989). For a similar purpose, a limited library of L1cIts391 genomic DNA was constructed according to the procedure described in Materials and Methods. The plasmids carrying the 1.4 kb *Bam*HI-*Sal*I fragment were selected for cloning the *ts* repressor gene of L1cIts391. These plasmids were preferentially selected since the restriction map of the above 2.1 kb *Bam*HI-*Pst*I L1 DNA fragment showed that the repressor gene of L1 was located within its 1.4 kb *Bam*HI-*Sal*I fragment (corresponding to 43933-45335 bp co-ordinates of L5). The enzyme *Bg*III cleaved the latter 1.4 kb DNA fragment into two smaller fragments, sizes 871 and 532 bp, respectively (data not shown). Among the 1.4 kb insert carrying plasmids, the ones that produced these two fragments upon digestion with *Bg*III were, therefore, selected (data not shown). One of them was picked up for further work and designated pBL1. The *Sal*I site of pBL1 was modified by the *Bam*HI linker DNA. The resulting plasmid was designated pBL2. The 1.4 kb *Bam*HI fragment of pBL2 was gel-purified and cloned to the identical site of pSD5S30. The resulting plasmid, designated pCP2, was transformed into *M. smegmatis* mc²155. The efficiency of plating of L1cI Γ on mc²155 carrying pCP2 was about 36% when compared to that on mc²155 carrying pSD5S30 or nonlysogen at 32°C (Table 1). Moreover, the plaque sizes that were produced by L1cI Γ on the lawn of mc²155 carrying pCP2 were reduced 5 and 3 folds at 32°C and 42°C, respectively (Table 1). Also, L1cI Γ produced turbid plaques on the lawn of mc²155 carrying pCP2 at 32°C, but not at 42°C. These results suggest that the *ts* repressor gene is indeed present in pCP2, but

	Helix 1	turn	Helix 2
Bxb1 gp69	E A D I A R M Y	G V T P Q A	V S W H V H
Ms6 gp6	E A D I A R M Y	G V T P Q A	V S W H V H
L1 cI	Q T E I A E L Y	G V T R Q A	V S W H K K
434 Cro	Q T E L A T K A	G V K Q Q S	I Q L I E A
P22 Cro	Q R A V A K A L	G I S D A A	V S Q W K E
434 R	Q A E L A Q K V	G T T Q Q S	I E Q L E N
λ R	Q E S V A D K M	G M G Q S G	V G A L F N
λ Cro	Q T K T A K D L	G V Y Q S A	I N K A I H
P22 R	Q A A L G K M V	G V S N V A	I S Q W E R
CAP	R Q E I G Q I V	G C S R E T	V G R I L K
LexR	R A E I A Q R L	G F R S P N	A A E E H L
LacR	L Y D V A Y E A	G V S Y Q T	V S R V V N
Bxz2 gp74	Q S E I A E M Y	G V T R Q Y	V S W I K H

Fig. 2. Alignment of helix-turn-helix motifs of proteins. The helix-turn-helix (HTH) motifs of some bacterial and phage-specific regulatory proteins are aligned. Highly conserved residues of HTH motifs are shaded. All of the sequences are available in GenBank. Helix 1, turn, and helix 2 regions of HTH motifs are shown by three rectangles at the top of the figure. The non-assigned amino acid residues of helix 1 of the HTH motif of the putative repressor of Bxz2 are underlined.

its expression is inadequate to completely block the vegetative growth of L1cI Γ .

A DNA sequence analysis of the 1.4 kb mutant L1 DNA fragment revealed that the 131st codon, CCC, in the wild-type cI gene of L1 was changed to CTC in the *ts* repressor gene of L1cIts391 (data not shown). The corresponding amino acid change occurred from proline to leucine in the *ts* repressor protein (Fig. 1). Interestingly, the deoxynucleotide change in the repressor gene of L1cIts391 created a novel restriction endonuclease site, *Sac*I, which is absent in wild-type L1. The *Sac*I also cleaved the 1.4 kb *Bam*HI-*Sal*I mutant DNA fragment into two smaller pieces with sizes of 846 and 557 bp, respectively (data not shown).

Identification of the putative domains and motifs in L1 repressor Specific regions of the repressor proteins of λ and other phages were shown to direct the specific functions, such as binding with their cognate operators through their HTH motifs, interaction with RNA polymerase, and the interaction between the monomers to form oligomers, etc. (Chattopadhyaya and Ghosh, 2003). To look for these regions in the L1 repressor, if any, its primary sequence was analyzed by several computer programs.

Alignment of gp6, gp69, gp74, and CI protein sequences by Clustal W showed at least 7 blocks (designated I-VII) carrying 3-8 identical amino acid residues (Fig. 1). These identical blocks were possibly assigned for similar work in all 4 proteins. Blocks I and II carried the 'turn' and 'helix 2' (recognition helix) region of the putative helix-turn-helix (HTH) motif, respectively. Alignment of this motif with similar motifs of the proteins of other phages revealed a couple of distinct features, which were not previously identified (Fig. 2). When the putative HTH motifs of the repressors of both L1 and Bxb1 were compared, twelve amino

acids were found common. The helix 2 and turn regions were more homologous than those of the helix 1 (Fig. 2). The HTH motifs of gp69 of Bxb1 and gp6 of Ms6 were 100% identical, although they are not 100% similar over the entire length of their amino acid sequences. Interestingly, the gp74 of Bxz2, which produced turbid plaques on *M. smegmatis* (Pedula *et al.*, 2003) and showed a good degree of identity with the repressors of Bxb1 and L1, carried only the helix 2 and turn region of the HTH motif (Fig. 2). A DNA-protein translation program showed that there were 14 extra amino acid residues at the upstream of the N-terminal methionine residue of Bxz 2 gp74 (data not shown). Six of these 14 amino acid residues may potentially form the helix 1 of HTH motif (Fig. 2). It would be interesting to see whether Bxz2 really synthesizes these 14 amino acids prior to the synthesis of gp74 from methionine. However, a further analysis revealed that the helix 1 and turn of the HTH motif of CI of L1 was about 55% identical to the same phage 434 Cro protein; whereas, helix 2 is about 44% identical to the same P22 Cro protein (Fig. 2). The most striking feature is that there are three highly charged basic residues in the helix 2 of the HTH motif of CI of L1. Genetic studies showed that substitutions of the acidic or amide derivatives of the acidic amino acids of helix 2 of 434 CI, CAP, lexR to basic or non polar amino acids greatly reduced the binding affinity to their cognate operators (Ebright *et al.*, 1984; Wharton and Ptashne, 1987; Thliveris and Mount, 1992). On the contrary, replacement of the glycine residue of the recognition helix with asparagine increases the affinity of the λ repressor to its operator (Hochschild *et al.*, 1986). Furthermore, it was reported that the affinity of the L5 repressor to its operator DNA is low compared to that of the λ (Brown *et al.*, 1997) and Bxb1 repressors (Jain and Hatfull, 2000). Taken together, it suggests that three basic amino acids (especially, arginine at position 12 of the helix 2 in the HTH motif) possibly affect the binding of the L1 repressor to the bases at the major groove of its operator.

Block VI and its neighboring sequences strongly control the function of the L1 repressor since the mutations at positions 123, 124, and 131 made it biologically inactive at 42°C (Hatfull, 2000; our observation, see above). Similarly, the amino acid residues encompassing block V are crucial for the repressor proteins function since the L5 repressor, carrying a point mutation at position 108, forms a clear plaque (Hatfull, 2000).

The hydrophobicity plot of CI of L1 revealed that the region carrying block IV and its upstream 13 amino acid residues was much more polar than the flanking regions (data not shown). It was also discovered that the two flanking regions of the L1 repressor, carrying amino acid residues 1-71 and 93-183, are distinctly basic (pI 10.56) and acidic (pI 4.23) in nature, respectively. The Bxb1 repressor also carries both domains at the N- and C-terminal ends, but they are much weaker than those found in the L1 repressor (data not shown). These basic N-terminal and acidic C-terminal domains are also prevalent in the 434 and P22 repressor proteins.

The amino acid residues at the downstream of position 131 in the L1 repressor must be vital for its function since the mapping of the different *cIts* mutations showed that mutations 29 and 578 are located at the right side of mutation 391 (Chaudhuri *et al.*, 1993; data not shown). The function of the block III amino acid residues might also be important, because it was shown that the point mutation in lambda and other phage repressor proteins greatly affect their structure and function (Chattopadhyay and Ghosh, 2003).

Acknowledgments We thank Drs. B. Bloom and Anil Tyagi for the phage and bacterial strains (such as mycobacteriophage L1, *M. smegmatis*, and plasmid like pSD5S30). The technical assistance of Mr. N. C. Datta is thankfully acknowledged. This research was partly supported by a research grant from C. S. I. R., Govt. of India, New Delhi.

References

- Barletta, R. G., Kim, D. D., Snapper, S. B., Bloom, B. R. and Jacobs, W. R., Jr. (1992) Identification of expression signals of the mycobacteriophage Bxb1, L1 and TM4 using the Escherichia-Mycobacterium shuttle plasmids pYUB75 and pYUB76 designed to create translational fusions to the *lacZ* gene. *J. Gen. Microbiol.* **138**, 23-30.
- Bashyam, M. D., Kaushal, D., Dasgupta, S. K. and Tyagi, A. K. (1996) A study of the mycobacterial transcriptional apparatus: identification of novel features in promoter elements. *J. Bacteriol.* **178**, 4847-4853.
- Brown, K. L., Sarkis, G. J., Wadsworth, C. and Hatfull, G. F. (1997) Transcriptional silencing by the mycobacteriophage L5 repressor. *EMBO J.* **16**, 5914-5921.
- Chaudhuri, B., Sau, S., Datta, H. J. and Mandal, N. C. (1993) Isolation, characterization and mapping of *temperature-sensitive* mutations in the genes essential for lysogenic and lytic growth of the mycobacteriophage L1. *Virology.* **194**, 166-172.
- Chattopadhyaya, R. and Ghosh, K. (2003) A comparative three-dimensional model of the carboxy-terminal domain of the lambda repressor and its use to build intact repressor tetramer models bound to adjacent operator sites. *J. Struct. Biol.* **141**, 103-114.
- Chattopadhyay, C., Sau, S. and Mandal, N. C. (2003) Cloning and characterization of the promoters of temperate mycobacteriophage L1. *J. Biochem. Mol. Biol.* **36**, 586-592.
- Dasgupta, S. K., Bashyam, M. D. and Tyagi, A. K. (1993) Cloning and assessment of mycobacterial promoters by using a plasmid shuttle vector. *J. Bacteriol.* **175**, 5186-5192.
- Datta, H. J. and Mandal, N. C. (1998) Identification of an early positive regulatory gene of mycobacteriophage L1. *J. Gen. Virol.* **79**, 205-210.
- Ebright, R. H., Cossart, P., Gicquel-Sanzey, B. and Beckwith, J. (1984) Mutations that alter the DNA sequence specificity of the catabolite gene activator protein of *E. coli*. *Nature* **311**, 232-235.
- Garcia, M., Pimentel, M. and Moniz-pereira, J. (2002) Expression

- of mycobacteriophage Ms6 lysis genes is driven by two σ 70-like promoters and is dependent on a transcription termination signal present in the leader RNA. *J. Bacteriol.* **184**, 3034-3043.
- Hatfull, G. F. (2000) *Molecular Genetics of Mycobacteria*. Hatfull, G. F. and Jacob, W. R., Jr. (eds.), pp. 37-54. ASM Press. Washington D.C., USA.
- Hochschild, A., Douhan, J. III. and Ptashne, M. (1986) How λ Repressor and λ Cro distinguish between O_{R1} and O_{R3}. *Cell* **47**, 807-816.
- Jain, S., Kaushal, D., DasGupta, S. K. and Tyagi, A. K. (1997) Construction of shuttle vectors for genetic manipulation and molecular analysis of Mycobacteria *Gene* **190**, 37-44.
- Jain, S. and Hatfull, G. F. (2000) Transcriptional regulation and immunity in mycobacteriophage Bxb1. *Mol. Microbiol.* **38**, 971-985.
- Lee, M. H., Pascopella, L., Jacobs, W. R. Jr. and Hatfull, G. F. (1991) Site-specific integration of mycobacteriophage L5: integration-proficient vectors for *Mycobacterium smegmatis*, *Mycobacterium tuberculosis*, and bacille Calmette-Guerin. *Proc. Natl. Acad. Sci. USA* **88**, 3111-3115.
- Nesbit, C. E., Levin, M. E., Donnelly-Wu, M. K. and Hatfull, G. F. (1995) Transcriptional regulation of repressor synthesis in mycobacteriophage L5. *Mol. Microbiol.* **17**, 1045-1056.
- Pedulla, M. L., Ford, M. E., Houtz, J. M., Karthikeyan, T., Wadsworth, C., Lewis, J. A., Jacobs-Sera, D., Falbo, J., Gross, J., Pannunzio, N. R., Brucker, W., Kumar, V., Kandasamy, J., Keenan, L., Bardarov, S., Kriakov, J., Lawrence, J. G., Jacobs, W. R. Jr., Hendrix, R. W. and Hatfull, G. F. (2003) Origins of Highly Mosaic Mycobacteriophage Genomes. *Cell*. **113**, 171-182.
- Ramesh, G. and Gopinathan, K. P. (1995) Cloning and characterization of mycobacteriophage I3 promoters. *Indian J. Biochem. Biophys.* **32**, 361-367.
- Sambrook, J., Fritsch, E. F. and Maniatis, T. (1989) *Molecular Cloning: A Laboratory Manual*. 2nd ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York, USA.
- Thliveris, A. T. and Mount, D. W. (1992) Genetic identification of the DNA binding domain of *Escherichia coli* LexA protein. *Proc. Natl. Acad. Sci. USA* **89**, 4500-4504.
- Wharton, R. P. and Ptashne, M. (1987) A new-specificity mutant of 434 repressor that defines an amino acid-base pair contact. *Nature* **326**, 888-891.