

# Decreasing effect of an anti-Nfa1 polyclonal antibody on the in vitro cytotoxicity of pathogenic *Naegleria fowleri*

Seok-Ryoul JEONG<sup>1)</sup>, Su-Yeon KANG<sup>1)</sup>, Sang-Chul LEE<sup>1)</sup>, Kyoung-Ju SONG<sup>1)</sup>,  
Kyung-il IM<sup>2)</sup> and Ho-Joon SHIN<sup>1)</sup>\*

<sup>1)</sup>Department of Microbiology, Ajou University School of Medicine, Suwon 442-749,

<sup>2)</sup>Department of Parasitology and Institute of Tropical Medicine, Yonsei University College of Medicine, Seoul 121-752, Korea

**Abstract:** The *nfa1* gene was cloned from a cDNA library of pathogenic *Naegleria fowleri* by immunoscreening; it consisted of 360 bp and produced a 13.1 kDa recombinant protein (rNfa1) that showed the pseudopodia-specific localization by immunocytochemistry in the previous study. Based on the idea that the pseudopodia-specific Nfa1 protein mentioned above seems to be involved in the pathogenicity of *N. fowleri*, we observed the effect of an anti-Nfa1 antibody on the proliferation of *N. fowleri* trophozoites and the cytotoxicity of *N. fowleri* trophozoites on the target cells. The proliferation of *N. fowleri* trophozoites was inhibited after being treated with an anti-Nfa1 polyclonal antibody in a dose-dependent manner for 48 hrs. By a light microscope, CHO cells co-cultured with *N. fowleri* trophozoites (group I) for 48 hrs showed severe morphological destruction. On the contrary, CHO cells co-cultured with *N. fowleri* trophozoites and anti-Nfa1 polyclonal antibody (1:100 dilution) (group II) showed less destruction. In the LDH release assay results, group I showed 50.6% cytotoxicity, and group II showed 39.3%. Consequently, addition of an anti-Nfa1 polyclonal antibody produced a decreasing effect of in vitro cytotoxicity of *N. fowleri* in a dose-dependent manner.

**Key words:** *Naegleria fowleri*, cytotoxicity, proliferation, *nfa1* gene

## INTRODUCTION

*Naegleria fowleri*, a free-living amoeba widespread in moist soil, water and sediment, exists as a virulent pathogen causing fatal primary amoebic meningo-

encephalitis (PAME) in experimental animals and humans (Culbertson, 1971; Marciano-Cabral, 1988; Ma et al., 1990; Kollars and Wilhelm, 1996). To obtain antigenic molecules to be used as diagnostic agents in pathogenic *N. fowleri* infection, we previously cloned an antigenic gene (called *nfa1*) from a cDNA library of *N. fowleri* by immunoscreening, using infected and immune mouse sera (Shin et al., 2001). We reported that the *nfa1* gene had the coding nucleotide sequence of 360 bp, producing a recombinant protein (rNfa1) of 13.1 kDa (Shin et al., 2001). An anti-Nfa1 polyclonal antibody obtained from mice immunized with a rNfa1

• Received 16 January 2004, accepted after revision 9 February 2004.

• This study was supported by a grant (No. R01-2000-000-00077-0) from the Korea Science & Engineering Foundation and in part by a grant from the Ajou University School of Medicine in 2000.

\*Corresponding author (e-mail: hjshin@ajou.ac.kr)

protein was used in immunocytochemistry, showing the Nfa1 protein as an indication of the pseudopodia-specific immunolocalization on a trophozoite of *N. fowleri* (Cho et al., 2003).

We produced an anti-Nfa1 polyclonal antibody based on the hypothesis that Nfa1 protein involved in pseudopodia activity may be concerned with the pathogenicity of *N. fowleri*. And then we obtained a preliminary data of the anti-Nfa1 antibody causing a decreasing effect on the cytotoxicity of *N. fowleri* trophozoites against CHO (Chinese hamster ovary) cells, much like treating an anti-Nfa1 antibody on a co-cultivating system (Cho et al., 2003).

In this research, for a more detailed analysis, we observed whether an anti-Nfa1 polyclonal antibody effects the proliferation of *N. fowleri* trophozoites in cultivation and on the in vitro cytotoxicity of pathogenic *N. fowleri* in a time- or a dose-dependent manner.

## MATERIALS AND METHODS

### Cultivation of *N. fowleri* and CHO cells

*N. fowleri* trophozoites (Cater NF69 strain, ATCC No.30215) were axenically cultured at 37°C in Nelson's media (Willaert, 1971). CHO cells were cultured with EMEM (Earle's Minimum Essential Medium) containing 10% fetal bovine serum (complete EMEM) at 37°C in a 5% CO<sub>2</sub> incubator (Im and Shin, 2003).

### Expression of the *nfa1* gene and production of a rNfa1 protein

To obtain a rNfa1 fusion protein, the expression and purification of the *nfa1* gene product were performed accordingly by the method mentioned in the previous paper (Shin et al, 2001). The purified DNA (5 µg/µl) obtained from PCR-T7/NT TOPO expression vector (Invitrogen, Grohingen, Netherlands) containing the *nfa1* gene was subsequently transformed into the BL21(DE3)pLysS *E. coli* strain by the heat-shock method. Cells were cultured at 37°C in the LAC (Luria-Bertani media containing 100 µg/ml of ampicillin and 34 µg/ml of chloramphenicol) plates for selection. A transformed-colony was selected and cul-

tured at 37°C in the LAC broth until the absorbance reached 0.5-0.8 at 600 nm; then 1 mM IPTG was added to the media. After 4 hrs of incubation, the cells were harvested by centrifugation (6,000 × g for 15 min). Cell extracts were compared with those of non-transformed BL21(DE3)pLysS by SDS-PAGE, and the presence of the expressed gene product was confirmed by Western blotting using both the immune and the infected sera, and anti-His and Xpress antibodies (Invitrogen).

### Production of an anti-Nfa1 polyclonal antibody

For the production of anti-Nfa1 polyclonal serum, the rNfa1 protein (50 µg/mouse) was mixed with an equal volume of Freund's complete adjuvant (Sigma), and the mixture was injected intraperitoneally into an 8-week-old female BALB/c mice (purchased from the Korea Institute of Science and Technology, Daejeon, Korea). The mouse was boosted biweekly for another 4 weeks with the rNfa1 protein (25 µg/mouse) containing an equal volume of incomplete Freund's adjuvant (Sigma). After the third boosting, the rNfa1 protein (5 µg/mouse) was injected intravenously without the adjuvant. Four days later, anti-Nfa1 polyclonal serum was collected from the mouse blood by centrifuging at 2,500 × g for 30 min at 4°C. ELISA was performed with a purified Nfa1 protein (5 µg/ml) and with a rabbit anti-mouse whole immunoglobulin (1:10,000 dilution) conjugated with alkaline phosphate (Sigma). Western blotting for the recombinant Nfa1 protein was performed according to the method in a previous paper (Cho et al., 2003).

### Proliferation of *N. fowleri* trophozoites

*N. fowleri* trophozoites (1 × 10<sup>4</sup> cells) were put in triplet culture tubes containing 1 ml of Nelson's medium. Then an anti-Nfa1 polyclonal antibody (1:200, 1:100 and 1:50 dilution) obtained from mice was added in each tube. Cultivating in a 37°C incubator, the number of trophozoites in each tube was counted with an haematocytometer up to 48 hrs post-incubation.

### Microscopic findings and LDH release assay for the cytotoxicity of *N. fowleri*

*N. fowleri* trophozoites containing CHO cells cultured in complete EMEM were put in each well of a 96-well plate (group I). For the experimental group (group II), an anti-Nfa1 polyclonal antibody (1:200, 1:100 and 1:50 dilution) obtained from mice was added in the triplet wells refer to the 96-well plate. After incubating at 37°C for 24 hrs and 48 hrs, respectively, the effect of the anti-Nfa1 polyclonal antibody was observed with a light microscope and LDH (lactate dehydrogenase) release assay. The cytotoxicity from the LDH release assay was observed with the CytoTox 96™ Non-Radioactive Cytotoxicity Assay Kit (Promega, WI, USA). About  $1 \times 10^4$  CHO cells were prepared with 100  $\mu$ l of EMEM in a round-bottomed 96-well plate. The amoeba trophozoites ( $1 \times 10^5$ ) and/or an anti-Nfa1 polyclonal antibody (1:200, 1:100 and 1:50 dilution) were added to each well, and five sets of triplet wells for controls including spontaneous releasing from medium and naive cells were prepared. The casted plate was incubated in a humidified chamber at 37°C and in a 5% CO<sub>2</sub> incubator for 45 min consecutively, and then centrifuged at 250  $\times$  g for 4 min. The supernatant (50  $\mu$ l) from each well was transferred to a fresh flat-bottomed 96-well plate. Equal volumes of the reconstituted Substrate Mix were added to each well. The plate was incubated at room temperature for 30 min. Fifty  $\mu$ l of Stop Solution was added, and after 1 hr, the absorbance was recorded at 490 nm.

$$\text{Cytotoxicity (\%)} = \frac{\text{Absorbance of (experimental group - control group)}}{\text{Absorbance of control group}} \times 100$$

## RESULTS

### Characterization of an anti-Nfa1 polyclonal antibody

To obtain a purified recombinant Nfa1 protein, a fusion protein of 17 kDa from the BL21 *E. coli* strain containing the *nfa1* gene was purified using a nickel resin column and enzyme digestion. Finally the purified protein eluted from the gels had a molecular

weight of 13.1 kDa, which corresponded well to the size in the previous research (Cho et al., 2003) (data not shown). An anti-Nfa1 polyclonal antibody was collected from mice immunized with a recombinant Nfa1 protein, and its specific reactivity against the Nfa1 protein was observed with ELISA and Western blotting. Values of anti-Nfa1 polyclonal antibodies by ELISA ranged from  $0.212 \pm 0.021$  (1:10,000 dilution, M  $\pm$  SD, N=5) to  $1.213 \pm 0.024$  (1:200 dilution), which showed a difference from the  $0.112 \pm 0.005$  (1:200 dilution) of the normal sera and also the  $0.109 \pm 0.005$  of the PBS control.

### Proliferation of *N. fowleri* trophozoites with an anti-Nfa1 antibody

*N. fowleri* trophozoites ( $1 \times 10^4$  cells) cultured from the beginning with an anti-Nfa1 polyclonal antibody (1:200 dilution) proliferated to  $1.1 \times 10^4$  cells in 8 hrs,  $2.9 \times 10^4$  cells in 24 hrs,  $3.9 \times 10^4$  cells in 32 hrs and  $15.1 \times 10^4$  cells in 48 hr post-incubation (Fig. 1). When an anti-Nfa1 antibody (1:100 dilution) was treated in cultivating system, *N. fowleri* trophozoites ( $1 \times 10^4$  cells) proliferated to  $1.0 \times 10^4$  cells in 8 hrs,  $2.5 \times 10^4$  cells in 24 hrs,  $3.2 \times 10^4$  cells in 32 hrs and  $6.7 \times 10^4$  cells in 48 hrs post-incubation (Fig. 1). In the control group, without the anti-Nfa1 antibody, the amoeba proliferated to  $1.7 \times 10^4$  cells in 8 hrs,  $4.9 \times 10^4$  cells in 24 hrs,  $8.5 \times 10^4$  cells in 32 hrs and  $21.7 \times 10^4$  cells in 48




Fig. 1. Proliferation of *Naegleria fowleri* trophozoites cocultured with an anti-Nfa1 polyclonal antibody. Box indicates the dilution ratio of the anti-Nfa1 polyclonal antibody.




Fig. 2. Microscopic findings of CHO cells (arrows) incubated in EMEM (A), with *Naegleria fowleri* trophozoites (arrow heads) (B), or with *N. fowleri* and an anti-Nfa1 antibody (C) 48 hrs post-cultivation ( $\times 200$ ).

hrs post-incubation (Fig. 1). Consequently, the addition of an anti-Nfa1 polyclonal antibody showed an inhibition on the proliferation of *N. fowleri* in a dose-dependent manner ( $p < 0.05$ ).




Fig. 3. A graphic view showing the addition of the anti-Nfa1 polyclonal antibody inhibited the cytotoxicity of *Naegleria fowleri* against CHO cells. Box indicates the time post-incubation.

#### Effects of the anti-Nfa1 antibody on the cytotoxicity of *N. fowleri*

CHO cells co-cultured with *N. fowleri* trophozoites (group I) for 24 hrs showed severe morphological destruction (Fig. 2B). On the contrary, CHO cells cocultured with *N. fowleri* trophozoites and anti-Nfa1 polyclonal antibody (group II) showed less destruction than above experimental group I (Fig. 2C). In the LDH release assay results, when an anti-Nfa1-antibody (1:200 dilution) was added in the co-cultivating system, group I showed 55.3% cytotoxicity and group II showed 43.7% in the LDH release assay 24 hrs post-incubation. After 48 hrs of incubation, group I showed 59.0% cytotoxicity, and group II showed 48.0% (Table 1). When an anti-Nfa1-antibody (1:100 dilution) was added in the co-cultivating system, group I showed 50.6% cytotoxicity 24 hrs post-incubation, and group II showed 39.3% (Table 1). After 48 hr of incubation, group I showed 53.3% cytotoxicity and group II 41.3%, respectively (Table 1). When an anti-Nfa1 antibody (1:50 dilution) was added in the co-cultivating system, group I showed 54.3% cytotoxicity 24 hrs post-incubation, and group II showed 34.0%. After 48 hrs of incubation, group I showed 66.3% cytotoxicity, and group II showed 42.0% (Table 1). Consequently, the addition of an anti-Nfa1 polyclonal antibody

**Table 1.** Cytotoxicity (%) of *Naegleria fowleri* against CHO cells by LDH release assay

Groups	1:200 dilution <sup>a)</sup>		1:100 dilution		1:50 dilution	
	24 hrs <sup>b)</sup>	48 hrs	24 hrs	48 hrs	24 hrs	48 hrs
CHO cells + trophozoites	55.3 ± 0.57	59.0 ± 2.00	50.6 ± 4.16	53.3 ± 3.51	54.3 ± 3.21	66.3 ± 5.51
CHO cells + trophozoites + anti-Nfa1 antibody	43.7 ± 1.52	48.0 ± 1.00	39.3 ± 3.05	41.3 ± 3.21	34.0 ± 3.61	42.0 ± 6.24

<sup>a)</sup>(v/v) ratio of anti-Nfa1 polyclonal antibody added in 200 µl of culture medium.

<sup>b)</sup>times post-cultivation.

showed a decreasing effect of in vitro cytotoxicity of *N. fowleri* in a dose-dependent manner (Fig. 3).

## DISCUSSION

PAME caused by *N. fowleri* is an acute progressive illness with the olfactory neuroepithelium as the major route of invasion for the *N. fowleri* (John DT, 1982). Concerned with host-tissue invasion, the adherence of the amoeba to host cells is the most important step in the mechanism of pathogenicity of *N. fowleri*, a specific pseudopodial projection, called an amoebastome, is formed (Marciano-Carbral, 1988). Other cytotoxic toxins and cytolytic proteinases have been proposed that amoeba destroy target cells (John, 1982; Marciano-Cabral, 1988; Ma et al., 1990).

In the previous research, we cloned an antigenic gene, *nfa1*, by using the immune and the infected sera and produced a recombinant 13.1 kDa protein that had a strong immunoreactivity against both the immune and the infected sera (Shin et al., 2001). The amino acid sequence and the secondary structure of the *nfa1* gene were related to those of the hemerythrin (Hr) or myohemerythrin (myoHr) gene of some marine invertebrates (Negri et al., 1994). The *nfa1* gene encoding myoHr-like protein was firstly cloned from amoeba, and its genetic relationship indicated a separate cluster from myoHr genes that were reported previously (Shin et al., 2001). In another study, trophozoites of *N. fowleri* in cultivating system and in mouse brain tissue infected experimentally with *N. fowleri* were well immunostained, as results of immunohistochemistry of the Nfa1 protein (Cho et al., 2003). The

immunolocalization study of the Nfa1 protein revealed that this protein was abundant in pseudopodia and around food vacuoles of *N. fowleri* trophozoites (Cho et al., 2003). This suggests that the Nfa1 protein is required for food ingestion and amoeba movement.

It was reported that highly virulent amoeba exhibited faster movement than weakly virulent amoeba (Cline et al, 1986). Based on the Nfa1 protein localization on the pseudopodia of *N. fowleri* in this study, the proliferation and the cytotoxicity of *N. fowleri* against CHO target cells were observed by adding an anti-Nfa1 antibody in the co-cultivation system to elucidate the antibody effect. It was shown that anti-Nfa1 antibody inhibited the proliferation of *N. fowleri* trophozoites in a dose-dependent manner. In addition, *N. fowleri* destroyed the CHO cells severely and showed a high cytotoxicity in the LDH release assay, but the cytotoxicity was decreased when an anti-Nfa1 antibody was added in a dose-dependent manner.

As iron-containing heme compounds like myoHr are essential for infectivity, invasion or destruction of the host tissue and anti-bacterial function (Negri et al, 1994; Raner et al., 1997), detailed biological roles of the Nfa1 protein need to be found in future studies. Moreover, further studies need to determine whether the Nfa1 protein is related to in vivo amoebic pathogenicity.

Finally, this study elucidates that the Nfa1 protein is a pathogen-related protein concerned with the pathogenesis of *N. fowleri*, especially with the amoebic proliferation and in vitro cytotoxicity.

## REFERENCES

- Cho MS, Jung SY, Park S, et al. (2003) Immunological characterizations of a cloned 13.1-kilodalton protein from pathogenic *Naegleria fowleri*. *Clin Diagn Lab Immunol* **10**: 954-959.
- Cline M, Carchman R, Marciano-Cabral F (1986) Movement of *Naegleria fowleri* stimulated by mammalian cells in vitro. *J Protozool* **33**: 10-13.
- Culbertson CG (1971) The pathogenicity of soil amoebas. *Annu Rev Microbiol* **25**: 231-254.
- Im K, Shin HJ (2003) *Acanthamoeba sohi*, n. sp., a pathogenic Korean isolate YM-4 from a freshwater fish. *Korean J Parasitol* **41**: 181-188.
- John DT (1982) Primary amebic meningoencephalitis and the biology of *Naegleria fowleri*. *Annu Rev Microbiol* **36**: 101-123.
- Kollars TM Jr., Wilhelm WE (1996) The occurrence of antibodies to *Naegleria* species in wild mammals. *J Parasitol* **82**: 73-77.
- Ma P, Visvesvara GS, Martinez AJ, et al. (1990) *Naegleria* and *Acanthamoeba* infections: Review. *Rev Infect Dis* **12**: 490-513.
- Marciano-Cabral F (1988) Biology of *Naegleria* spp. *Microbiol Rev* **52**: 114-133.
- Negri A, Tedeschi G, Bonomi F, Zhang JH, Kurtz DM Jr. (1994) Amino-acid sequences of the alpha- and beta-subunits of hemerythrin from *Lingula reevii*. *Biochim Biophys Acta* **1208**: 277-285.
- Raner GM, Martins LJ, Ellis WR Jr. (1997) Functional role of leucine-103 in myohemerythrin. *Biochemistry* **36**: 7037-7043.
- Shin HJ, Cho MS, Jung SY, et al. (2001) Molecular cloning and characterization of a gene encoding a 13.1 kDa antigenic protein of *Naegleria fowleri*. *J Eukaryot Microbiol* **48**: 713-717.
- Willaert E (1971) Isolement et culture *in vitro* des amibes de genre *Naegleria*. *Ann Soc Belg Med Trop* **51**: 701-708.