

Effect of Sonicated Extracts of *Enterococcus faecalis* on the Production of Matrix Metalloproteinase-8 by Human Polymorphonuclear Neutrophils

Sung-Kyung Park, Won-Jun Shon, Sung-Sam Lim*

Department of Conservative Dentistry, College of Dentistry, Seoul National University

ABSTRACT

This in vitro study monitored MMP-8 production on PMN by stimulated with the following three groups: Sonicated extracts of *E. faecalis* (SEF), SEF treated with Ca(OH)₂ (12.5mg/ml) for 7 days, and lipopolysaccharides (LPS) of *E. coli*. The level of MMP-8 in each group was immediately measured by ELISA. The data were analyzed with Kruskal-Wallis test and Mann-Whitney U test.

In the SEF group, the level of production of MMP-8 was higher than the negative control group in low concentration (0.05µg/ml) of SEF ($p < 0.05$), but it decreased with an increase in the concentration of SEF ($p < 0.05$). In the case of SEF treated with Ca(OH)₂, all of the MMP levels were higher than negative control group ($p < 0.05$), but no statistical difference was found among the different SEF concentrations ($p > 0.05$). All of the levels in *E. coli* LPS were increased with increasing concentrations ($p < 0.05$).

According to this study we could summarize as follows:

1. MMP-8 was expressed at low level in untreated PMN group and the levels of MMP-8 were upregulated in PMN stimulated by *E. coli* LPS groups.
2. In the SEF groups, the level of production of MMP-8 decreased with an increase in the concentration of SEF ($p < 0.05$). So *E. faecalis* may have suppressive effect on the production of MMP-8 by PMN.
3. In the case of SEF treated with Ca(OH)₂, all of the MMP levels at different SEF concentrations were higher than untreated PMN group ($p < 0.05$), but no statistical difference was found among the different SEF concentrations ($p > 0.05$). [J Kor Acad Cons Dent 30(2):138-144, 2005]

Key words : *E. faecalis*, MMP-8, Calcium hydroxide, PMN leukocyte

- received 2005. 1. 21, revised 2005. 3. 6, accepted 2005. 3. 7 -

I. INTRODUCTION

Bacteria and their by-products are considered to be

the primary etiologic agents of pulpal necrosis and periapical lesions¹⁾. Therefore, endodontic treatment aims at eliminating of microorganisms from the root canal system and preventing of reinfection. However some cases fail even when apparently well treated. In fact, a number of factors associated with failure of endodontic therapy. But, most treatment failures are caused by microorganisms persisting in the apical parts of root canals of obturated teeth. It seems to be bacterial effects in root canal and periapical lesion.

* Corresponding author: **Sung-Sam Lim**

Department of Conservative Dentistry,
College of Dentistry, Seoul National University,
28 YunGun-Dong, ChongNo-Gu Seoul, Korea, 110-749
Tel : 82-2-2072-2656 Fax : 82-2-2072-3859
E-mail : limss@snu.ac.kr

Especially, single species of gram-positive organisms, *Enterococcus faecalis* had been found to be one of predominant bacteria in teeth in which root canal therapy failed²³.

So, the use of intracanal medication to disinfect the root canal system has been advocated³¹. Calcium hydroxide ($\text{Ca}(\text{OH})_2$) has been the intracanal medication of choice. Sjögren et al.⁴ and Byström et al.³ reported that 7-days and 30-days applications of $\text{Ca}(\text{OH})_2$, respectively, eliminated bacteria that survived instrumentation in root canals. However, in case of *E. faecalis*, $\text{Ca}(\text{OH})_2$ dressing is ineffective³¹. Therefore it seems that the frequent isolation of Enterococci in failed endodontic cases puts in question the routine use of $\text{Ca}(\text{OH})_2$ as an intracanal medication⁵¹. Survival of *E. faecalis* in $\text{Ca}(\text{OH})_2$ at high pH was related to a functional proton pump⁷. However few researches reported how these factors affect the antibacterial activity of $\text{Ca}(\text{OH})_2$.

Matrix metalloproteinases (MMPs) form a family of structurally related but genetically distinct endopeptidases expressed at low level in normal tissue, but upregulated during inflammation⁸¹. They are group of zinc-dependent proteinases, secreted or released by host cells (ie, PMN, macrophage, bone cell, fibroblast) as proenzymes. There are multiple members of family, including interstitial collagenase (MMP-1), neutrophil collagenase (MMP-8), stromelysin-1 (MMP-3), stromelysin-2 (MMP-10), gelatinase B (MMP-9). Shin et al.⁹¹ reported that MMPs play an important role in the pulp tissue destruction in inflamed pulp and periapical lesion. MMP-8 of them has been thought to be uniquely produced by developing PMN cells in bone marrow, and that the MMP-8 activity is dependent on the release of the enzyme from PMN cells by degranulation¹⁰¹. However, recent findings demonstrated expression of MMP-8 in mesenchyme derived, non-PMN lineage cells, including dental pulp fibroblasts, odontoblasts, inflamed epithelial cells and plasma cells¹¹⁻¹²¹. In other research, MMP-8 is reported that they have an important role in the degenerations of the extracellular matrix in inflamed pulps and periapical lesions¹³¹.

In this study, the purpose was to examine the levels of production of MMP-8 by PMN stimulated with

Enterococcus faecalis and effect of $\text{Ca}(\text{OH})_2$ on their production.

II . MATERIALS AND METHOD

Preparation of bacterial sonicated extracts

Enterococcus faecalis (ATCC 29212) strains were used in this study. Strains were grown in 1-liter cultures in 85% N_2 -10% H_2 -5% CO_2 chamber for 3 days at 37°C. The medium used was 3.7% brain heart infusion broth supplemented with 0.5% yeast extract, 0.5mg L-cysteine hydrochloride, and 0.5% sodium bicarbonate.

Bacterial fractions were prepared. Briefly, bacterial cell harvest from 1-liter cultures were washed and suspended in 20ml of phosphate-buffered saline (PBS). To minimize the loss of protein, 1mM of Phenylmethylsulfonyl fluoride (PMSF) were added. PMSF act as protease inhibitor at protein extraction. And suspensions of bacterial cell were disrupted by sonication (100W output, Fisher sonicator) on ice for 5min with 30-sec pulses on and 10-sec pulse-off in the presence of glass beads. Sonicated debris was centrifuged at $12,000 \times g$ in a Sorvall RC5C (Sorvall Instruments, DuPont) for 20 min, and membrane fraction was sedimented by ultracentrifugation at $85,000 \times g$ for 60 min at 4°C. To investigate protein profile of sonicated extracts of *E. faecalis* (SEF), supernatant was filtered with 0.22 μm syringe and separated by 10% SDS-PAGE at 100V for 90min, and the protein concentration was determined by Bicinchoninic acid (BCA) protein assay (Pierce, Rockford, IL). The SDS-PAGE reveals that the soluble cytoplasmic protein of *E. faecalis* strain ATCC 29212 is mainly consist of 75kDa and 35-50kDa polypeptide. The pellet was resuspended in 20ml of 10mM Tris buffer (pH 7.4), and stocked in deep-freezer at -20°C.

Sonic extract of *E. faecalis* (SEF) treated with $\text{Ca}(\text{OH})_2$

Each of SEF was suspended to 100 $\mu\text{g}/\text{ml}$ with pyrogen-free water. $\text{Ca}(\text{OH})_2$ powder (Sigma Chemical Co., St Louis, MO, USA) 12.5 mg/ml was added. All samples were vortexed for 20 seconds once a day and incubated at 37°C for 7days. After that, upper part of

solution was ultrafiltered (Centriprep YM-10, 10,000 NMWL, Millipore Co., USA) and stocked in deep-freezer at -20°C.

Preparation of PMN

Using heparinized tubes, Venous blood was obtained from 6 healthy volunteers (26~35years). The erythrocytes of the obtained blood were segregated through 6% dextran (Mwt. 500,000, Sigma Chemical Co., St Louis, MO, USA), the leukocytes were extracted after centrifugation using lymphoprep (Nycomed Pharma AS, Oslo, Norway) and remained erythrocytes were eliminated by carrying out hypotonic lysis. After verification through the Trypan blue dye exclusion test & Giemsa staining, the PMN leukocytes that were obtained by this method proved to have 98% or more vitality and 97% or more purity. One millimeter of the extracted PMN leukocytes were 24 well plate with each well having a cell account of 10×10^4 .

Bacterial stimulation and MMP-8 production

SEF group and SEF treated with Ca(OH)_2 group and *E.coli* LPS were diluted at three concentration. And PMN media were stimulated by each groups and control group was not activated. Final concentrations of each group were $0.05\mu\text{g/ml}$, $0.5\mu\text{g/ml}$ and $5\mu\text{g/ml}$ and incubated at 37°C for 30 minutes.

Enzyme-linked immunosorbent assay (ELISA)

After incubation, the amount of MMP-8 present in the culture supernatants were assayed by Enzyme-linked immunosorbent assay (ELISA, R and D System, Minneapolis, USA) according to the manufacturer's protocols.

Statistical analysis

Data were statistically analyzed using the Man-Whitney rank sum test and Kruskal-wallis test. A value of $P < 0.05$ was considered statistically significant.

III. RESULTS

The results of this study are the mean values of six samples of duplicate cultures and summarized in Table 1, Figure 1 and Figure 2.

In the SEF groups, the level of production of MMP-8 was higher in comparison to the negative control group in low concentration ($0.05\mu\text{g/ml}$) of SEF ($P < 0.05$), but it decreased with an increase in the concentration of SEF ($P < 0.05$). In the case of SEF treated with Ca(OH)_2 , all of the MMP levels at different SEF concentrations were higher than control group ($P < 0.05$), but no statistical difference was found among the different SEF concentrations ($P > 0.05$). All of the levels in *E. coli* LPS were increased with increasing concentrations ($P < 0.05$).

Table 1. Average concentration(±SD) of MMP-8

Group	Mean concentration (±SD, $\mu\text{g/ml}$ protein)		
	$0.05\mu\text{g/ml}$	$0.5\mu\text{g/ml}$	$5\mu\text{g/ml}$
SEF	$2.3209 \pm 0.0398^*$	2.1303 ± 0.1216	$1.9447 \pm 0.1310^*$
SEF treated with Ca(OH)_2	2.8775 ± 0.2100	2.9003 ± 0.0896	2.7745 ± 0.0851
<i>E.coli</i> LPS	$2.4390 \pm 0.0866+$	$2.6975 \pm 0.1650+$	2.7929 ± 0.2193
Untreated PMN	2.2497 ± 0.0041		

SEF = sonicated extract of *Enterococcus faecalis*

+ and * are statically significantly different ($P < 0.05$) between groups

SEF groups and SEF treated with Ca(OH)_2 groups are statically significant different ($P < 0.05$) in comparison with untreated PMN group.




Figure 1. Average concentration of MMP-8 (\pm SD) (ng/mol)




Figure 2. Average concentration of MMP-8 (ng/mol)

IV. DISCUSSION

In recent years, MMPs have gained considerable attention in many studies. MMPs play an important role in the pulp tissue destruction of acute inflamed pulp⁹. The role of MMP-8 synthesized by the cells in the pulp-dentinal complex is not clear. But MMP-8 was reported that they have an important role in the degenerations of the extracellular matrix in inflamed pulps and periapical lesions¹³. Therefore, a hypothesis was proposed that MMP-8 could be present in the periapical lesion, and the enzyme level could be related to the activity of the periapical tissue. And MMPs are endopeptidases expressed at low level in normal tissues, but upregulated during inflammation⁸. So, untreated PMN had some MMP-8 in this study.

In previous studies, sonic extracts from bacteria were studied for their effects on the immune response. For example, sonicated material of *Fusobacterium nucleatum* can evoke a concentration-dependent stimulatory or suppressive effect on the proliferation rate of accessory cell¹⁴. Sonicated extract of *Actinobacillus actinomycetemcomitans* suppressed interleukin-2 production of T-cells¹⁵. Kim et al.¹⁶ showed that sonicated extracts of *Enterococcus faecalis* also suppressed the production of interleukin-2 and interleukin-4 in lymphocytes. So in this study, SBE may have depressive effect to PMN. In positive control group, all of the MMP-8 levels in *E.*

coli LPS were increased with increasing concentrations ($P < 0.05$).

In this study, MMP-8 production level at low level of SEF ($0.05\mu\text{g/ml}$) was seen a little elevation compare to untreated control group ($P < 0.05$). But it decreased with an increase in the concentration of SEF ($P < 0.05$). This phenomenon showed that ability of *E. faecalis* to impair the production of MMP-8 may be related to the suppression of PMN. This fact may be seen another immunosuppressive effect of SEF.

However, there is no study how *E. faecalis* have immunosuppressive effect exactly. But *E. faecalis* is the most prevalent species of enterococci in root canals of failed case¹⁷ and for it to be involved in the pathogenesis and maintenance of apical periodontitis, it must survive in the root filled canals as where the nutrient supply is limited. *E. faecalis* has been shown to have relatively uncommon capacity to survive in root canals as a monoculture without the support of other bacteria¹⁸. *E. faecalis* is also known to survive periods of starvation in water¹⁹ and infected dentin in water for periods of at least 10 days²⁰.

Ca(OH)₂ has been widely used as an intracanal medication to eliminate bacteria and is currently acknowledged to be one of the most effective antimicrobial dressings used during endodontic therapy²¹. Although the exact mechanism of action of Ca(OH)₂ is still unknown, its antimicrobial activity is generally considered to be related to the release of hydroxyl

ions in an aqueous environment, producing a pH of approximately 12.5, even in very dilute mixture. Most endodontic pathogens are unable to survive in this alkaline environment²¹. The mechanism that Ca(OH)₂ uses to eliminate bacteria may include damage to bacteria cytoplasmic membrane by including lipid peroxidation, protein denaturation, and damage to bacterial DNA and by serving as physical barrier that withholds nutrients for bacterial growth and limits space for bacterial multiplication²².

But *E. faecalis* is known to withstand a high pH. This is an identifying characteristic of *E. faecalis*. At pH 11.4 or greater, *E. faecalis* does not survive, yet it can survive at a pH below 11.5 as shown in this and earlier studies³. Because of the buffering effect of dentin²³⁻²⁵, it is unlikely that the high pH of Ca(OH)₂ (> 11.5) is attained within dentinal tubules where *E. faecalis* has the capacity, at least in vitro, to penetrate deeply^{20,26-28}. In radicular dentin, alkalinity may only reach pH 10.3 after dressing the canal with Ca(OH)₂²⁴. An adaptive response in alkaline pH and stress-induced protein synthesis appear to play minor roles in cell survival, a functional proton pump with the capacity to acidify the cytoplasm is critical for survival of *E. faecalis* at high pH⁷.

So considered *E. faecalis* in endodontic treatment, additive medication is recommended. For example, chlorhexidine (CHX) is a broad-spectrum antimicrobial agent that has been reported to be an effective medicament in endodontic treatment therapy²⁹. As a root canal irrigant and intracanal medicament, CHX has an antibacterial efficacy comparable to that of sodium hypochlorite (NaOCl)³⁰⁻³¹. In addition, it is also effective against strains resistant to Ca(OH)₂³². Therefore the use of combination of Ca(OH)₂ and CHX is recommended in endodontic treatment.

In our study, Ca(OH)₂ had some effect to production of MMP-8 by PMN. But comparison to untreated PMN group, this data does not support that Ca(OH)₂ removed immunosuppressive effect of SEF. MMP-8 levels of Ca(OH)₂ treated SEF group had another impression, this may because of the high pH of Ca(OH)₂. Gram positive microorganisms have external membrane structure, cell wall. This cell wall have two important high molecule, peptidoglycan and teichoic acid. In this study, SEF treated with Ca(OH)₂

for 1 week showed different appearance in the production of MMP-8. This result means that Ca(OH)₂ could possibly modified SEF. The high pH of Ca(OH)₂ may modify acidity of teichoic acid. But no statistical difference was found among the different concentrations ($P > 0.05$).

In our study, this result suggested that bacterial protein sonicated extract of *E. faecalis* can inhibit the immune response by PMN and this could be one of possible factors why *E. faecalis* are found frequently in the teeth with failed endodontic treatment. The use of an antimicrobial agent as intracanal medication between appointments may increase the chances for successful endodontic treatment by reducing residual bacteria in root canal system. In endodontic treatment, Ca(OH)₂ may have limited antibacterial effect to *E. faecalis*. So our study suggests that the use of only Ca(OH)₂ in clinical endodontic treatment may not be effective to *E. faecalis*.

V. CONCLUSION

According to this study we could summarize as follows:

1. MMP-8 was expressed at low level in untreated PMN group and the levels of MMP-8 were upregulated in PMN stimulated by *E. coli* LPS groups.
2. In the SEF groups, the level of production of MMP-8 decreased with an increase in the concentration of SEF ($p < 0.05$). So *E. faecalis* may have suppressive effect on the production of MMP-8 by PMN.
3. In the case of SEF treated with Ca(OH)₂, all of the MMP levels at different SEF concentrations were higher than SEF & untreated PMN groups ($p < 0.05$), but no statistical difference was found among the different SEF concentrations ($p > 0.05$).

REFERENCE

1. Kakehashi S, Stanley HR, Fitzgerald RJ. The effects of surgical exposures of dental pulps in germ-free and conventional laboratory rat. *Oral Surg Oral Med Oral Pathol* 20:340-4, 1965.
2. Sundqvist G, Figdor D, Persson S, Sjögren U. Microbiologic analysis of the teeth with failed endodontic treatment and outcome of conservative retreatment.

- Oral Surg Oral Med Oral Pathol Oral Radiol Endo* 85:86-93, 1998.
3. Byström A, Claesson R, Sundqvist G. The antibacterial effect of camphorated paramonochlorophenol, camphorated phenol and calcium hydroxide in the treatment of infected root canals. *Endod Dent Traumatol* 1:170-5, 1985.
 4. Sjögren U, Figdor D, Spangberg L, Sundqvist G. The antimicrobial effect of calcium hydroxide as a short-term intracanal dressing. *Int Endod J* 24:119-25, 1991.
 5. Molander A, Reit C, Dahlen G, Kvist T. Microbiological status of root-filled teeth with apical periodontitis. *Int Endod J* 31:1-7, 1998.
 6. Love RM. *Enterococcus faecalis*—a mechanism for its role in endodontic failure. *Int Endod J* 34:399-405, 2001.
 7. Evans M, Davies JK, Sundqvist G, Figdor D. Mechanisms involved in the resistance of *Enterococcus faecalis* to calcium hydroxide. *Int Endod J* 35:221-8, 2002.
 8. Birkedal-Hansen H. Proteolytic remodelling of extracellular matrix. *Current Opinion in Cell Biology* 7:728-35, 1995.
 9. Shin SJ, Lee JI, Baek SH, Lim SS. Tissue levels of matrix metalloproteinases in pulps and periapical lesions. *J Endod* 28:313-5, 2002.
 10. Mainardi CL, Pourmotabbed TF, Hasty KA. Inflammatory phagocytes and connective tissue degrading metalloproteinases. *Am J Med Sci* 302:171-5, 1991.
 11. Hanemaaijer R, Sorsa T, Konttinen YT, Ding Y, Sutinen M, Visser H, van Hinsbergh VW, Helaakoski T, Kainulainen T, Ronka H, Tschesche H, Salo T. Matrix metalloproteinase-8 is expressed in rheumatoid synovial fibroblasts and endothelial cells. Regulation by tumor necrosis factor- α and doxycycline. *J Biol Chem* 272:31504-9, 1997.
 12. Palosaari H, Wahlgren J, Larmas M, Ronka H, Sorsa T, Salo T, Tjaderhane L. The expression of MMP-8 in human odontoblasts and dental pulp cells is down-regulated by TGF- β 1. *J Dent Res* 79:77-84, 2000.
 13. Wahlgren J, Salo T, Teronen O, Luoto H, Sorsa T, Tjaderhane L. Matrix metalloproteinase-8 (MMP-8) in pulpal and periapical inflammation and periapical root-canal exudates. *Int Endod J* 35:897-904, 2002.
 14. Yoshida H, Jontell M, Sundqvist G, Bergenholtz G. Effect of sonicated material from *Fusobacterium nucleatum* on the functional capacity of accessory cells derived from dental pulp. *Oral Microbiol Immunol* 10:208-12, 1995.
 15. Kurita-Ochiai T, Ochiai K, Ikeda T. Immunosuppressive effect induced by *Actinobacillus actinomycetemcomitans*: effect on immunoglobulin production and lymphokine synthesis. *Oral Microbiol Immunol* 7(6):338-43, 1992.
 16. Son HH, Lim S, Shon W, Kim HS, Lee W. Effects of sonicated *Enterococcus faecalis* extracts on Interleukin-2 and Interleukin-4. *J Endod* 30(10):701-3, 2004.
 17. Peciuliene V, Reynaud AH, Balciuniene I, Haapasalo M. Isolation of yeasts and enteric bacteria in root-filled teeth with chronic apical periodontitis. *Int Endod J* 34:429-34, 2001.
 18. Fabricius L, Dahlen G, Holm SE, Moller AJ. Influence of combinations of oral bacteria on periapical tissues of monkeys. *Scand J Dent Res* 90:200-6, 1982.
 19. Hartke A, Giard JC, Laplace JM, Auffray Y. Survival of *Enterococcus faecalis* in an oligotrophic microcosm: changes in morphology, development of general stress resistance, and analysis of protein synthesis. *Appl Environ Microbiol* 64:4238-45, 1998.
 20. Orstavik D, Haapasalo M. Disinfection by endodontic irrigants and dressings of experimentally infected dentinal tubules. *Endod Dent Traumatol* 6:142-9, 1990.
 21. Heithersay GS. Calcium hydroxide in the treatment of pulpless teeth with associated pathology. *J Br Endod Soc* 8:74-93, 1975.
 22. Siqueira Junior JF, Lopes HP. Mechanisms of antimicrobial activity of calcium hydroxide: a critical review. *Int Endod J* 32:361-9, 1999.
 23. Wang JD, Hume WR. Diffusion of hydrogen ion and hydroxyl ion from various sources through dentine. *Int Endod J* 21:17-26, 1988.
 24. Nerwich A, Figdor D, Messer HH. pH changes in root dentin over a 4-week period following root canal dressing with calcium hydroxide. *J Endod* 19:302-6, 1993.
 25. Haapasalo HK, Siren EK, Waltimo TM, Orstavik D, Haapasalo MP. Inactivation of local root canal medications by dentine: an in vitro study. *Int Endod J* 33:126-31, 2000.
 26. Haapasalo M, Orstavik D. In vitro infection and disinfection of dentinal tubules. *J Dent Res* 66:1375-9, 1987.
 27. Safavi KE, Spangberg LS, Langeland K. Root canal dentinal tubule disinfection. *J Endod* 16:207-10, 1990.
 28. Vahdaty A, Pitt Ford TR, Wilson RF. Efficacy of chlorhexidine in disinfecting dentinal tubules in vitro. *Endod Dent Traumatol* 9:243-8, 1993.
 29. Delany GM, Patterson SS, Miller CH, Newton CW. The effect of chlorhexidine gluconate irrigation on the root canal flora of freshly extracted necrotic teeth. *Oral Surg Oral Med Oral Pathol* 53:518-23, 1982.
 30. Ohara P, Torabinejad M, Kettering JD. Antibacterial effects of various endodontic irrigants on selected anaerobic bacteria. *Endod Dent Traumatol* 9:95-100, 1993.
 31. Jeansonne MJ, White RRA. Comparison of 2.0% chlorhexidine gluconate and 5.25% sodium hypochlorite as antimicrobial endodontic irrigants. *J Endod* 20:276-8, 1994.
 32. Siqueira Junior JF, Batista MM, Fraga RC, de Uzeda M. Antibacterial effects of endodontic irrigants on black-pigmented gram-negative anaerobes and facultative bacteria. *J Endod* 24:414-6, 1998.

국문초록

ENTEROCOCCUS FAECALIS 추출물이 다형핵 백혈구의 METALLOPROTEINASE-8 분비에 미치는 영향에 관한 연구

박성경 · 손원준 · 임성삼*

서울대학교 치과대학 치과보존학교실

*Enterococcus faecalis*의 추출물을 성인의 말초혈액으로부터 얻은 인간의 백혈구를 이용해 Matrix Metalloproteinase-8 (MMP-8)의 생산에 미치는 영향에 대해서 알아보았다.

*Enterococcus faecalis*를 배양한 뒤 초음파 분쇄를 하여 추출물을 얻어냈다. 이중 일부는 적절한 농도의 $\text{Ca}(\text{OH})_2$ 를 처리하였다. 이후 *E. faecalis* 추출물군, *E. faecalis* 추출물에 $\text{Ca}(\text{OH})_2$ 를 처리한 군으로 분리하여 다형핵백혈구를 자극하여 MMP-8의 생산량을 측정하였다.

MMP-8는 처리하지 않은 PMN group에서도 나타났고, PMN을 *E. coli* LPS로 자극시킨 경우 증가하였다 ($p < 0.05$). SEF group 들에서는 MMP-8의 생성량이 SEF의 농도가 증가함에 따라 감소하였다 ($p < 0.05$). 따라서 *Enterococcus faecalis*는 PMN의 MMP-8 생산 억제 효과를 가진다고 볼 수 있다. $\text{Ca}(\text{OH})_2$ 로 처리한 SEF 군의 경우 각기 다른 SEF 농도에서도 모두 MMP-8 생성량이 아무 처리도 하지 않은 PMN 군보다 많았다 ($p < 0.05$). 하지만 각 군들 사이에 통계적 차이는 없었다 ($p > 0.05$).

주요어 : *E. faecalis*, MMP-8, 수산화칼슘, 다형핵백혈구